



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



9.75.
a. 8.



E. BIBL. RADCL.


12.
13. 3.
10.
27/2
18
12.
13. 3.
14.
69.2.





600027048R

19176 d 27/2



FLORA
BOREALI-AMERICANA.

TO THE

MEMBERS OF THE BOARD OF DIRECTORS

FLORA

BOREALI-AMERICANA,

SISTENS CHARACTERES PLANTARUM
quas in America septentrionali collegit et detexit

ANDREAS MICHAUX,
Instituti Gallici Scientiarum, necnon Societatis Agriculturæ
Caroliniensis socius.

TABULIS ÆNEIS 51 ORNATA.

TOMUS SECUNDUS.

TYPIS CAROLI CRAPELET.

PARISIIS & ARGENTORATI,
Apud fratres LEVRAULT.

ANNO XI — 1803.

1900

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX AND TILDEN FOUNDATIONS

1195 FIFTH AVENUE

NEW YORK

1900

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX AND TILDEN FOUNDATIONS

1195 FIFTH AVENUE

NEW YORK

1900

~~~~~

# FLORA

## BOREALI-AMERICANA.

---

### CLASSIS XIV.

### DIDYNAMIA.

### GYMNOSPERMIA.

#### TEUCRIUM. L.

*Corollae* labium superius (nullum) ultra basim  
2-partitum, divaricatum ubi stamina.

CANADENSE. T. caule herbaceo, erecto : foliis lanceolatis, serratis, subtus canescentibus :  
L. spica conferta, multiflora, longa.

*Obs.* Variat foliis ovato-lanceolatis. *T. virginicum* videtur hujusce varietas.

*HAB.* in Virginia, Pennsylvania, Canada; Carolina superiore.

#### HYSSOPUS. L.

*Corollae* labium inferius lacinula intermedia crenata. *Stamina* recta, distantia.

NEPETOIDES. H. caule altissimo, acute quadrangulo : foliis ovalibus : spica longa, confertiflora : corolla lutescente.

*HAB.* juxta flumen *Ohio* et in Kentucky.

2 DIDYNAMIA. GYMNOSPERMIA.

NEPETA. L.

*Corollae* labium inf. lacinula intermedia crenata :  
*faux* margine reflexo. *Stamina* approximata.

CATARIA. N. floribus spicatis : verticillis subpedicellatis : foliis petiolatis, cordatis, dentato-serratis.

MENTHA. L.

*Cor.* subæqualis, quadrifida, lacinia latiore emarginata. *Stamina* erecta, distantia.

BOREALIS. M. assurgens, humilis, pubescens : foliis petiolatis, ovali-lanceolatis, utrinque acutissimis : floribus verticillatis : staminibus exertis.

*HAB.* juxta amnes ad sinuum *Hudsonis* defluentes.

TENUIS. M. erecta, glabriuscula : foliis lanceolato-ovalibus, subsessilibus : spica gracili, verticillis minimis interrupta : staminibus non exertis.

*M. viridis.* WALT.

*HAB.* in humidis Carolinæ et Georgiæ.

ISANTHUS. L.

*Ires*, æqualis. *Aster*, flos : corolla subregularis.

Genus Labiatum. Herba, foliis basinerviis; floribus axillaribus nec verticillatis.

CAL. patentiuscule campanulatus, semi5-fidus : laciniis subæqualibus, lanceolatis; duabus inferioribus (præsertim elapsa corolla) magis conniventibus.

COR. calyci subæqualis, subregularis : tubus rectiusculus, anguste cylindraceus : limbus subrotatus, 5-partitus; laciniis conformibus, ovalibus; intermedia inferiore vix longiore et basi bimaculata.

STAM. subæqualia, corolla paulo breviora, distincte erecta : antheræ obliquatæ, inferne bipartitæ; loculis ovatis, ad angulum fere rectum divergentibus.

PIST. ovarium 4-fidum : stylus longitudine staminum, superne recurvus : stigmata duo, recurvo-potentissima, lineari-ligulata; inferiore paulo promissiore.

FRUCT. globuliquatuor, calycispatuli tubum adimplentes, obovati, reticulato-rugosi, basibus tantum connexi, monospermi.



4 DIDYNAMIA. GYMNOSPERMIA.

CÆRULEUS. I. subviscido-puberulus : foliis ovali-lanceolatis, utrinque acutis, trinerviis :  
*tab. 30.* pedunculis 1-2-floris.

*Obs.* Herba annua, erecta, habitu fere SATUREIÆ  
*hortensis*. Corolla dilute cœrulea.

*HAB.* in cretaceis Virginia Carolina Kentucky.

L A M I U M. L.

*Corollae* labium superius integrum, fornicatum ;  
labium inferius 2-lobum ; faux utrinque mar-  
gine dentata.

HISPIDULUM. L. caule hispidulo : foliis longe  
petiolatis, lato-cordatis, parce puben-  
tibus : foliorum floralium axillis uniflo-  
ris : corolla majuscula, alba.

*HAB.* in sylvis Tennessee opacis.

S T A C H Y S. L.

*Corollae* labium superius fornicatum ; lab. in-  
ferius lateribus reflexum : lacinia intermedia  
majore, emarginata. *Stamina* deflorata versus  
latera reflexa.

HYSSOPIFOLIA. S. glabriuscula, gracilis, erecta :  
foliis sessilibus, lineari-lanceolatis, ra-  
riter subdentatis : verticillis spicae sub-  
quadrifloris.

*HAB.* in Carolina.



P. J. Redoute Del.

Plén. Sc.

**ISANTHUS** *caeruleus.*



**DIDYNAMIA. GYMNOSPERMIA. 5**

**ASPERA.** S. caulis. erectis, retrorsum hispidissimis : foliis subpetiolatis, lanceolatis, argute serratis : spica verticillis subsexfloris : calycibus rigide spinescentibus.

**Obs.** Affinis *S. palustri*.

**HAB.** in campestribus Carolinæ.

**CLINOPODIUM. L.**

*Involucrum* multisetum, verticillo subjectum.

**VULGARE.** C. capitulis subrotundis, hispidis,  
**L.** bracteis setaceis.

**HAB.** in montibus Pensylvaniæ et in Canada.

**BRACHYSTEMUM.**

*Brachys*, brevis. *Στημων*, stamen : *stamina brevia*.

Labiatae perennes : folia puncticulosa : verticilli confertiflori seu capitula ; bracteolis ciliatis.

**CAL.** tubulosus ; dentibus quinque, brevibus, subæqualibus, erectis, acutis : fauce nuda.

**COR.** tubus longitudine calycis, gracilis : labium superius breve, porrectum, leviter emarginatum : labium inferius multoties amplius, patens, obtuse trilobum ; intermedio productiore, subligulato-oblongo.

6 DIDYNAMIA. GYMNOSPERMIA.

STAM. filamentis brevissimis inclusa, sub-  
æqualia.

SEM. cylindraceo-oblonga.

VERTICILLATUM. B. foliis ovali-lanceolatis, integris : verticillis infra terminalem paucis,  
*tab. 31.* compactis.

*HAB.* in montibus Pennsylvania, adusque Carolinam superiorem.

MUTICUM. B. foliis lanceolato-ovalibus, leviter  
*tab. 32.* rariterque dentatis, glabellis : capitulis terminalibus; bracteolis lanceolatis, non acuminatis.

*HAB.* in Carolina superiore.

VIRGINICUM. B. caule stricte erecto, pube asperiuscula : foliis lanceolatis vel linearibus, integerrimis, glabellis : capitulis subfasciculatis.

*THYMUS virginicus.* LINN.

*HAB.* in Pennsylvania, Virginia, regione Illinoensi et in montibus Carolinae.



*P. J. Redouté Del.*

*Plu. 50.*

**BRACHYSTEMUM** *verticillatum*.







P. J. Redoute Del.

Plac. Sc.

**BRACHYSTEMUM** *muticum*.



PYCNANTHEMUM.

*Herbes*, densus. *Arbores*, flos : *flores conferti*.

Herbæ perennes seu suffrutices. Folia punctata. Capitula subsimplicia seu composita, multibracteata.

CAL. tubulosus, multistriatus, 5-fidus : dentibus erectis, semilanceolatis seu subulatis.

COR. tubus longitudine calycis : labium superius subrecurvo - porrectum, oblongiusculum, modice convexum, apice rotundatum, subintegrum : labium inferius multo majus, recurvo - patens, subcanaliculatum, trifidum ; laciniis lateralibus subsemi-ellipticis, intermedia longiore pauloque latiore, nonnihil repandula.

STAM. exerta, distantia ; duo labio superiori longitudine subæqualia, duo conspicue breviora : antheræ loculis subparallelis, &c.

Obs. Genus affine SATUREIÆ.

INCANUM. P. caule pubescente : foliis petiolatis, ovalibus, in acumen acutissimum sensim angustatis, serratis, molliter incano-tomentosis ; capitulis fasciculato-

8 DIDYNAMIA. GYMNOSPERMIA.

compositis, terminalibus lateralibusque;  
bracteolis setaceis.

CLINOPODIUM *incanum*. LINN.

ORIGANUM *incanum*. WALT.

HAB. in Virginia et Carolina.

ARISTATUM. P. subcandicans : foliis lanceolato-  
ovalibus, subserratis, breviter petiola-  
tab. 33. tis : verticillis paucis capituloque termi-  
nali sessilibus; bracteolis linearibus,  
aristato-acuminatis.

NEPETA *virginica*. LINN.

HAB. a Marylandia ad Carolinam superiorem.

MONTANUM. P. foliis ovali-lanceolatis, serratis :  
capitulo sessili : calycibus approximatis,  
erectis, breviter dentatis.

Obs. Caulis partimque aliæ partes solito purpureæ.


Folia subsessilia.

HAB. in altis montibus Carolinæ.

MONARDELLA. P. hirsutulum : foliis abrupte pe-  
tiolatis, subcordato-ovalibus, serratis :  
tab. 34. capitulo bracteis, omnibus coloratis, in-  
volucrato; bracteolis ciliatis : calycibus  
summitate barbatis.

Obs. Habitus MONARDÆ *fistulosæ*. Binæ ultimæ  
species vix hujus generis ! Calyce diverso ; stigma-  
tis lacinia altera brevissima.

HAB. in altis montibus Carolinæ.




P. J. Redouté Del.

Plée. Sc.

**PYCNANTHEMUM** *aristatum.*







P. J. Redouté Del.

Plac. Sc.

**PYCNANTHEMUM** *Monardella* ?.



H Y P T I S. JACQ.

*Cal.* turbinatus, 5-fidus. *Cor.* infundibuliformis, fauce ampliata, limbo patens. *Stam.* in labium inferius declinata.

CAPITATA. H. foliis inferne cuneatim angustatis:

*Jacq.* capitulis pedunculatis, involucratis, lateralibus terminalibusque.

*HAB.* in Carolina inferiore.

T H Y M U S. L.

*Calycis* bilabiati faux villis clausa.

CAROLINIANUS. T. erecto - cespitosus : foliis glabris, ovalibus, subintegris : verticillis e binis fasciculis substipitatis : calycibus sub3-fido-bilabiatis.

THYMERA caroliniana. WALT.

*HAB.* in Carolina et Georgia, secus amnem Savannah, ad urbem Augusta.

10 DIDYNAMIA. GYMNOSPERMIA.

DRACOCEPHALUM. L.

*Corollae* faux inflata : labium superius concavum.

VIRGINIANUM. D. glabellum : caulibus simplicibus : foliis lanceolatis , serratis : spica aphylla : corolla purpurea.

*Obs.* Variat foliis linearibus, lanceolatis et ovali-lanceolatis.

PRASII tres species *Walterianae* hujus sunt varietates.

*HAB.* in montibus Carolinæ, Virginiae, &c.

TRICHOSTEMA. L.

*Corollae* labium superius falcatum. *Stam.* longissima.

DICHOTOMA. T. foliis subrhomboideo-lanceolatis vel linearibus : ramulis floriferis bifurcatis , staminibus longissimis.

*HAB.* a Pennsylvania ad Carolinam.

DIDYNAMIA. GYMNOSPERMIA. II

BRUNELLA. L.

*Filamenta* bifurca : altero apice antherifera. *Stigma* bifidum.

VULGARIS. B. foliis oblongo-ovalibus vel sublan-  
L. ceolatis , omnibus indivisis , dentatis.

HAB. a Canada ad Carolinam.

SCUTELLARIA. L.

*Calyx* ore integro : post florescentiam clauso ,  
operculato.

LATERIFLORA. S. ramosissima , glabriuscula : fo-  
L. liis longe petiolatis , dentatis ; caulinis  
subcordatis : rameis , ovali-lanceolatis :  
floribus minimis.

OBS. Flores cœrulei.

HAB. in Canada , montibus *Alléghanis* et in Ken-  
tucky.

PILOSA. S. caulibus foliisque pilosissimis : foliis  
subrhomhoideo-ovalibus , crenatis : flo-  
ribus laxiuscule spicatis ; bracteis par-  
vulis.

SCUTELLARIA *caroliniana*? WALT.

HAB. in Carolina et Georgia.

PARVULA. S. pusilla : dense pubescens : foliis ova-



14 DIDYNAMIA. *ANGIOSPERMIA.*

RIGENS. V. caulibus rigide erectis : foliis ob-  
ovalibus , subtomentoso-hirsutissimis :  
spicis imbricatis.

*OBS.* Spicæ strictæ, longæ, pubentes. Corolla majus-  
cula, coerulea. Proxima *V. carolinianæ*.

*HAB.* in regione Illinoensi.

HASTATA. V. erecta ; foliis lanceolatis , sensim  
L. angustatis, acutissimis, inciso-serratis :  
spicis paniculatis , linearibus , subim-  
bricatis.

*OBS.* Folia primaria solito aurito-trifida.

*HAB.* in Nova Anglia, lacu *Ontario*, regione Illi-  
noensi, Carolina superiore, &c.

ANGUSTIFOLIA. V. humilis erecta : foliis lineari-  
lanceolatis , rariter subserratis : spicis  
filiformibus.

*HAB.* in Tennassée et in comitatu *Carlisle* Pennsyl-  
vaniæ.

CAROLINIANA. V. erecta : foliis oblongo-obova-  
L. libus, obtusis : spicis filiformibus, lon-  
gissimis , distinctifloris.

*OBS.* Corolla dilute rubens.

*HAB.* in Carolina.

SPURIA. V. caule decumbente , ramosissimo ,  
L. divaricato : foliis ovalibus, incisis : spicis

**DIDYNAMIA. ANGIOSPERMIA. 15**

filiformibus : bracteis calyces superantibus.

*Obs.* Corolla subcœrulea.

*HAB.* in regione Illinoensi.

**URTICIFOLIA.** V. erecta : foliis ovalibus seu ovali-

*L.* lanceolatis, acutis : paniculis divaricatis ; e spicis filiformibus, distinctifloris.

*Obs.* Corolla alba.

*HAB.* in Virginia, Carolina.

**L I P P I A. L.**

*Cal.* 4 - dentatus, subrotundus, erectus, compresso-membranaceus. *Caps.* 1-locul. 2-valvis, 2-sperma.

**NODIFLORA.** *L?* repens, lævis : foliis cuneato-obovalibus : pedunculis solitariis, elongatis.

*VERBENA nodiflora.* LINN.

*BLAIRIA nodiflora.* GÆRT.

*HAB.* ad littora Carolinæ et ad ripas fluminis *Ohio*.

**LANCEOLATA.** *L?* repens, lævis : foliis lineari-lanceolatis, argute serratis : pedunculis solitariis, elongatis.

*HAB.* in Carolina, juxta amniculum *Ashley*.

16 DIDYNAMIA. *ANGIOSPERMIA.*

EUPHRASIA. *L.*

*Cal.* 4-fidus, cylindricus. *Caps.* 2-locularis, ovato-oblonga. *Antheræ* inferiores altero lobo basi spinosæ.

OFFICINALIS. *E. foliis ovalibus, lineatis, argute dentatis.*

MELAMPYRUM. *L.*

*Cal.* 4-fidus. *Corollæ* labium superius compressum, margine replicato. *Caps.* 2-locularis, obliqua, hinc dehiscens. *Semina* 2, gibba.

AMERICANUM. *M. humile, gracile; foliis lineari-lanceolatis; superiorum basi parce setaceo-dentata; paribus omnibus distinctis et distanter floridis.*

*HAB.* a sinu *Hudsonis* ad montosam Carolinam.

DIDYNAMIA. *ANGIOSPERMIA*. 17

R H I N A N T H U S. *L.*

*Cal.* 4-fidus, ventricosus. *Caps.* 2-locularis, obtusa, compressa.

CRISTA-GALLI. *R.* corollis labio superiore compresso, brevior.

*Obs.* Europæo simillimus; parvitate tantum discrepans.

*HAB.* in Canada.

B A R T S I A. *L.*

*Cal.* 2-lobus, emarginatus, coloratus. *Corolla* minus ipso calyce colorata: labio superiore longiore. *Caps.* 2-locularis.

PALLIDA? *B.* foliis alternis, linearibus, indivisis; superioribus lanceolatis; floralibus subovalibus, summitate subdentatis: spica terminali, hirsuta.

*HAB.* in Canada.

COCCINEA. *B.* foliis alternis, linearibus, rariter inciso-dentatis vel laciniatis: bracteis dilatatis, coloratis, solito trifidis.

Var. *α. coccinea*: bracteis superne coccineis.

— *β. pallens*: bracteis lutescentibus. Rarior.

*HAB.* a Nova Anglia ad Floridam.

18 DIDYNAMIA. *ANGIOSPERMIA*.

PEDICULARIS. *L.*

*Cal.* 5-fidus. *Caps.* 2-locularis, mucronata, obliqua. *Semina* tunicata.

LANCEOLATA. *P. simplex* : foliis lanceolatis, incisodentatis; spica aphylla : calycibus glabris : capsulis brevibus.

*HAB.* in regione Illinoensi.

GLADIATA. *P. simplex* : spica foliosa, alterniflora : capsula in mucronem gladiatum ipsa longiorem protracta : floribus lutescentibus.

*HAB.* in Pennsylvania.

CANADENSIS. *P. caule simplici, debili* : capitulo *L.* basi frondoso : calycibus deorsum truncatis.

*HAB.* a Canada ad Carolinam.

BUCKNERA. *L.*

*Cal.* 5-dentatus. *Corollae* limbus 5-fidus, æqualis : lobis cordatis. *Caps.* 2-locularis.

AMERICANA. *B. caule simplici* : foliis lineari-lanceolatis vel obovali-lanceolatis, rariter subdentatis, asperis, trinerviis.

*HAB.* in Carolina.

SCHWALBEA.

*Cal.* 4-fidus : lobo superiore minimo ; infimo maximo, emarginato.

AMERICANA. SCHWALBEA.

*L.*

*HAB.* in Carolina inferiore.

GERARDIA. *L.*

*Cal.* 5-fidus. *Cor.* bilabiata : labio inferiore 5-partito : lobis emarginatis : medio 2-partito.

*Caps.* 2-ocularis, dehiscens.

PURPUREA. *G.* caule opposite ramosissimo : foliis

*L.* angusto-linearibus : floribus sparsis, purpureis, subsessilibus.

*HAB.* a Nova Anglia ad Carolinam.

FLAVA. *G.* caulibus simplicibus : foliis lan-

*L.* ceolatis, integris vel dentatis ; inferioribus solito subpinnatifido-incisis : corollis amplissimis, luteis.

*HAB.* a Nova Anglia ad Floridam.

PEDICULARIA. *G.* caule brachiatim paniculato :

*L.* foliis oblongis, duplicatim inciso-serratis, subpinnatifidis : calycis laciniis foliaceis, incisis.

*HAB.* a Pennsylvania ad Carolinam.

22 DIDYNAMIA. *ANGIOSPERMIA*.

*CAPRARIA*. L.

*Cal.* 5-partitus. *Cor.* campanulata, 5-fida, acuta.

*Caps.* 2-valvis, 2-locularis, polysperma.

**MULTIFIDA.** C? annua, erecta, viscidulo-pubens :

*tab.* 35. foliis terno-verticillatis, multifidis : pedicellis solitariis.

*HAB.* in ripis arenosis fluminum amniculorumque, in *Tennessee* et *Illinoensi* regione.

*MONNIEBA*. BROWN.

→ *GRATIOLA*. L.

*Cal.* 2-bracteatus. *Stam.* vix fauce exerta. *Cor.* subæqualiter 5-fida. *Caps.* dissepimento valvis parallelo.

**ROTUNDIFOLIA.** *M.* minutim pubens : foliis sub-ovali-orbiculatis, multinerviis : pedunculis passim oppositis, folia subæquantibus.

*HAB.* in regione *Illinoensi*.

**AMPLEXICAULIS.** *M.* caulibus lanatis : foliis breviter cordato-ovalibus, amplexicaulibus, integris, obtusis : pedunculis folio brevioribus.

*HAB.* in fossis, stagnis *Carolinæ*.

**CUNEIFOLIA.** *M.* glaberrima : foliis cuneato-ob-



P. J. Redouté Del.

Plu. Sc.

CAPRARIA *multifida*.





**DIDYNAMIA. ANGIOSPERMIA. 23**

ovalibus, superne obsolete subcrenatis:  
pedunculis folia subæquantibus.

*HAB.* in locis mari inundatis Carolinæ inferioris.

**MIMULUS. L.**

*Cal.* 4-dentatus, prismaticus. *Cor.* ringens: labio-  
superiore lateribus replicato. *Caps.* 2-locu-  
laris, polysperma.

**RINGENS.** M. erectus, foliis oblongis, linearibus,  
L. sessilibus.

*HAB.* a Pennsylvania ad Carolinam.

**RUELLIA. L.**

*Cal.* 5-partitus. *Cor.* subcampanulata. *Stam.*  
per paria approximata. *Capsula* dentibus elas-  
ticis dissiliens.

**HUMISTRATA.** R. glabriuscula: caule diffuse ra-  
moso, humistrato et radicante: foliis in  
petiolum longiuscule angustatis, ovali-  
bus, obtusis: floribus subsessilibus: cap-  
sulis linearibus.

*HAB.* ad fines Georgiæ et Floridæ.

**OBLONGIFOLIA.** R. assurgens, tota minutim den-  
seque pubens: foliis erectis, subsessili-

24 DIDYNAMIA. *ANGIOSPERMIA*.

bus, obovali-oblongis : floribus subso-  
litariis.

*HAB.* in Georgia.

**STREPENS.** *R. erecta vel assurgens, hirsuta : foliis*  
*L.* in petiolum angustatis vel ovali-lanceo-  
latis : fasciculis axillaribus, subsessilibus,  
subtrifloris : calycibus hispidis.

*HAB.* a Virginia ad Floridam.

CHELONE. *L.*

*Cal.* 5-partitus, 3-bracteatus. Filamentum quin-  
tum sterile, cæteris brevius. *Caps.* 2-locularis,  
2-valvis. *Semina* plurima, margine membra-  
nacea.

**GLABRA.** *C. glabella : foliis ovalibus vel lanceo-*  
*L.* latis, inæqualiter serratis : floribus spi-  
catis.

*Var. α. purpurea.*

— *β. alba.*

*HAB. α.* in rivulis altissimorum montium Carolinae.

— *β.* a Pennsylvania ad Carolinam, juxta rivulos.

BIGNONIA. L.

*Cal.* 5-fidus, cyathiformis. *Cor.* fauce campanulata, 5-fida, subtus ventricosa. *Siliqua* 2-locularis. *Semina* membranaceo-alata.

CATALPA. B. arborea : foliis simplicibus, ternis,  
L. cordatis : panicula laxissima ; floribus  
diandris : capsula gracili, longa, tereti.

*HAB.* ad ripas fluviorum *Ohio* et *Mississippi*.

CAPREOLATA. B. petiolis brevibus, cirrhosis, bi-  
L. foliolatis : foliolis glabris, oblongo-ovalibus,  
basi subcordatis : pedunculis axillaribus,  
aggregatis : capsula lineari, plana.

*HAB.* a Virginia ad Floridam.

RADICANS. B. caule scandente, geniculis strigillatim  
L. radicanibus : foliis pinnatis ; foliolis serratis,  
acuminatis : floribus fasciculato-terminalibus.

*HAB.* a Pennsylvania ad Floridam.

OROBANCHE. L.

*Cal.* 2-fidus. *Corolla* ringens. *Caps.* 1-locularis,  
2-valvis, polysperma. *Glandula* sub basi ovarii.

VIRGINIANA. O. caule ramoso : floribus distanter  
L. alternis, parvulis : corollis deciduis,  
quadridentatis.

HAB. a Canada ad Georgiam ; in radice FAGI nec aliæ  
plantæ : notissima sub nomine *Cancer root*.

AMERICANA. O. scapo humili, squamis ovali-lan-  
L. ceolatis, imbricatis oblecto : spica glabra :  
corollis recurvatis : staminibus exertis.

HAB. a Pennsylvania ad Carolinam.

UNIFLORA. O. scapis, omitta basi, nudis, uniflo-  
ris : calyce ebracteato : corolla recurvata.

HAB. a Canada ad Virginiam.

TETRADYNAMIA. *SILICULOSA*. 27

CLASSIS XV.

TETRADYNAMIA.

*SILICULOSA*.

COCHLEARIA. L.

*Silicula* emarginata, turgida, scabra : valvulis gibbis, obtusis.

HUMIFUSA. C? caule humifuso, pubente : foliis pinnatifidis ; pinnulis incis : spicis lateralibus ; siliculis didymis, subrugosis.

BISCUTELLA *apetala*. WALT.

LEPIDIUM *didymum*. LINN.

*Obs.* Forsan in posterum una cum CORONOPo, genus constitutura : diversa tamen defectu corollæ et staminibus perfectis tantum 2, filamentis 4 exantheratis.

*HAB.* in cultis, campestribus, ruderalis Carolinæ.

LEPIDIUM. L.

*Silicula* emarginata, cordata, polysperma : valvulis carinatis, contrariis.

VIRGINICUM. L. foliis radicalibus pinnatifidis : caulinis lineari-lanceolatis, plus minus in-

30 TETRADYNAMIA. *siliquosa*.

ROTUNDIFOLIA. C. caulibus simpliciusculis, debiliter procumbentibus: foliis omnibus simplicibus, suborbiculatis, integriusculis.

*HAB.* in rivulis altissimorum montium Carolinæ.

DENTARIA. L.

*Siliqua* elastice dissiliens valvulis revolutis. *Stigma* emarginatum. *Calyx* longitudinaliter connivens.

CONCATENATA. D. radice tuberibus oblongo-ovoides, angusto nexu concatenatis: caulibus ternate triphyllis; foliis subtrifoliolatis, foliolis incisis, oblongis.

Affinis *D. enneaphyllae*. LINN. Flores cœrulescentes.

*HAB.* in Nova Anglia et altissimis montibus Carolinæ.

DIPHYLLA. D. radice dentata: caulibus approximatis, diphyllis; foliis trifoliolatis, oblongis, inæqualiter incisis; floribus flavescentibus.

*Obs.* Radices ab indigenis loco SINAPIS, pro ciborum condimentis exsiccantur.

*HAB.* in sylvis vastis Tennassée et montibus altissimis Carolinæ.

**TETRADYNAMIA. SILIQUOSA. 31**

**HESPERIS. L.**

*Petala* oblique flexa. *Glandula* intra stamina breviora. *Siliqua* stricta. *Stigma* basi bifurca apice connivente. *Cal.* clausus.

**PINNATIFIDA.** H. foliis argute serratis; superioribus lanceolatis, inferioribus inferne pinnatifidis: siliquis breviter pedunculatis, breviusculis.

*HAB.* in humidis sylvarum Tennassée.

**ARABIS. L.**

*Glandulae* nectariferae 4, singulae intra calycis foliola, squamæ instar reflexæ.

**FALCATA.** A. foliis lanceolatis, utrinque angustatis, obsolete subdentatis, sessilibus; siliquis pendulis, ancipitibus, falcatis.

*HAB.* in rupestribus Canadae et Novæ Angliæ ad Virginiam.

**ERYSIMUM. L.**

*Siliqua* columnaris, exacte tetraedra. *Cal.* clausus.

**OFFICINALE.** E. siliquis spicæ adpressis, foliis runcinatis.

*HAB.* in Virginia.



32 TETRADYNAMIA. *SILIQVOSA*.

CLEOME. *L.*

*Glandulae* nectariferæ 3, ad singulum sinum calycis singulæ, excepto infimo. *Petala* omnia adscendentia. *Siliqua* 1-locularis, 2-valvis.

DODECANDRA. *C. floribus* dodecandris : foliis ternatis.  
*L.*

---

CLASSIS XVI.

MONADELPHIA.

TRIANDRIA.

SISYRINCHIUM. L.

Monogyna. *Spatha* 2-phylla. *Petala* 6, plana.  
*Caps.* 3-locularis, infera.

BERMUDIANA. *S. foliis gramineis : seapo dilatato-  
L. ancipite : involucris muticis, totidem  
ramulos terminantibus.*

*Obs. S. anceps* CAVAN. videtur varietas, superne 2-3-  
ramulosa.

*HAB.* a Pennsylvania ad Carolinam.

MUCRONATUM. *S. foliis angustissimis : scapis seta-  
ceis, simplicissimis : gluma terminali  
unica, bivalvi, colorata; valvæ exterioris  
mucrone recto, floribus longiore.*

*Obs. Foliis ima parte glumisque amethystæis.*

*HAB.* in Pennsylvania.

*PENTANDRIA.*

*ILLECEBRUM. L.*

*Cal.* 5-phyllus, cartilagineus. *Cor.* nulla. *Stigma* simplex. *Caps.* 5-valvis, 1-sperma.

POLYGONOIDES. *I.* caulibus hirsutis : foliis ovalibus : capitulis alternatim axillaribus , arcte sessilibus.

*I. achyrantha. WALT.*

*HAB.* in viis urbis *Charlstown.*

VERMICULATUM. *I.* glabrum : foliis subtereti-linearibus , carnosius : capitulo sessiliter terminali.

*HAB.* in maritimis *Florida.*

*ERYTHROHIZA.*

Perennis ; acaulis , radicalifolia ; in scapo nudo minutim spiciflora : squamæ gemmales plurimæ basim scapi obvestientes.

*CAL.* subcampanulatus , 5-partitus , erecto-appressus ; laciniis oblongis.

*COR.* sub5-petala : petala subspathulata , inferne erecta , superne extra calycem patentiuscula.

**MONADELPHIA. PENTANDRIA. 35**

Basibus, mediante imissimo staminum tubo, quasi connexa; una cum isto decidunt.

**STAM.** 5, 1-delpha; petalis dimidio breviora; tubus cylindraceus, margine rotundus; filamentis 5 setaceis, castratis; 5 alternis brevioribus crassioribusque antheriferis: antheræ subrotundæ, introrsum 2-loculares;

Una cum corolla fundo calycis, ad contactum ovarii inserta.

**PIST.** ovarium latiuscula basi affixum; subrotundum, summitate attenuatum quasi in brevissimum crassumque stylum: stigma subovato-capitatum, sulcato-trifidum.

**CAPS.** calycē appresso paulo longior; brevivoidea; 3-ocularis, superne trivalvis; valvis medio septiferis.

**SEM.** plurima, columellæ crassiusculæ adnexa; inæqualiter angulosa, hinc attenuata: tegumine rugelloso, nucleo ampliore.

**OBS.** Interdum quarta pars in pistillo et fructu additur.

Genus *Ericaceum*.

**ROTUNDIFOLIA.** E. glaberrima: foliis orbiculato-

36 MONADELPHIA. PENTANDRIA.

reniformibus, dentatis : spica longis-  
tab. 36. sima.

*GALAX aphylla*. LINN. An *Pyröla floribus albis*,  
*spicatis* ; *caule aphylo* ; *folio subrotundo*, *ser-*  
*rato*, *pediculo longissimo insidenti*. CLAYT.  
n° 31.

*Obs.* *Radix repens*, saturate rubra, fere rubiacea.  
Folia radicalia, promisse petiolata. Scapus basi squa-  
mis imbricatis stipatus, plantagineus, pedalis et  
ultra : spica longa : floribus albis, pusillis, copiosis,  
brevissime pedicellatis.


*HAB.* in excelsis Carolinæ montibus. Maio florens.

L I N U M. L.

*Cal.* 5-phyllus. *Petala* quinque. *Caps.* 5-valvis,  
10-ocularis. *Sem.* solitaria.

VIRGINIANUM. L. foliis lineari-lanceolatis, sparsis,  
L. rarius oppositis : summo caule panicu-  
lato; ramulis remote alternifloris : caly-  
cibus acutis; petalis luteis.

*HAB.* a Pennsylvania ad Carolinam.



P. J. Redoute Del.

Plat. Sc.

**ERYTHRORHIZA** *rotundifolia*.



*PASSIFLORA*. *L.*

*Trigyna*. *Cal.* 5-phyllus. *Petala* 5. *Nectarium*-  
corona. *Bacca* pedicellata.

*INCARNATA*. *P.* foliis ad ipsam paginæ basim bi-  
*L.* glandulosis, tripartitis, serrulatis : co-  
ronæ filamentis variegatis : bracteis par-  
vulis, brevi-ovalibus, crenulatis : bacca  
ovata.

*HAB.* a Virginia ad Floridam.

*EUTEA*. *P.* caule, petiolis foliisque subtus pi-  
*L.* losis : foliis abbreviato-cordatis, trilobis ;  
lobis sinubusque obtusis ; margine inte-  
gerrimo : glandulis nullis : pedunculis  
plerumque geminis, unifloris.

*OBS.* Flos parvulus, lutescens : fructus pusillus, glô-  
bosus. In regionibus meridionalibus folia profun-  
dius lobata et solito maculata : in septentrionalibus  
brevissime rotundato-lobata.

*HAB.* in Virginia et regione Illinoensi, usque ad  
Floridam.



*DECANDRIA.*

*GERANIUM. L.*

Monogyna. *Stigmata* quinque. *Fructus* rostratus, 5-coccus.

CAROLINIANUM. *G. annuum*, diffusum : foliis oppositis, 5-partito-multifidis : pedunculis bifloris, subfasciculatis ; fructiferis erectis.

*Obs.* Calyces lanceolato-ovales, aristati. Petala emarginata : fructus hirsuti.

*HAB.* in Carolina utraque. ♂

MACULATUM. *G. perenne*, retrorsum pubens : caulibus erectis, opposite diphyllis : foliis 3-5-partitis : pedunculis elongatis, bifloris : petalis obovalibus.

*Obs.* Calyces pilosi, lato-ovales, abrupte aristati. Petala non emarginata.

*HAB.* a Canada ad Carolinam et in regione Illinoensi. ♀.

*OXALIS. L.*

*Cal.* 5-phyllus. *Petala* unguibus connexa. *Caps.* angulis dehiscens, 5-gona.

ACETOSELLA. *O. radice* dentata : acaulis, tenella : foliis trifoliatis, foliolis dilatato-obcor-

**MONADELPHIA. DECANDRIA. 39**

datis, lobis rotundatis : scapo radicali,  
1-floro : corolla albido-carnea.

*HAB.* in Americæ borealis sylvis, a Quebec ad sinum  
*Hudsonis.*

**VIOACEA.** O. scapo radicali, umbellifero : pedi-

*L.* cellis subviscido-pubentibus : calycis laciniis apice callo croceo : corolla violacea.

*HAB.* a Nova Anglia ad Carolinam.

**CORNICULATA.** β. O. tota conspicue pubescens :

*L.* caule ramoso, diffuso seu procumbente : pedunculis umbelliferis ; pedicellis longiusculis : floribus luteis.

*Obs.* Flores majusculi, amplitudine *Acetosellæ*, etiam paulo majores. An inde distincta ab Europæa species?

*HAB.* in cultis, a Canada ad Carolinam.

**STRICTA.** O. tota hirsutula : caulibus stricte erectis :

*L.* tis : pedunculis elongatis ; umbella composita.

*HAB.* in Virginia.

*P O L Y A N D R I A.*

*H A L E S I A. L.*

*Cal.* 4-dentatus, superus *Cor.* 4-fida. *Nux* quadrangularis, 2-sperma.

N°. A foliis nulla recte eruitur specierum differentia, neque a glandulis petiolaribus.

PARVIFLORA. *H.* fructu minore, deorsum promisse angustato, inde quasi clavato, inæqualiter subtetraptero : floribus parvulis.

*HAB.* in Florida, circa *Matança.*

TETRAPTERA. *H.* fructu subæqualiter rhomboideo-  
- *L.* tetragonoptero : floribus majoribus.

*HAB.* in sylvis umbrosis et rivulosis Carolinae.

DIPTERA. *H.* fructu subelliptico, quasi diptero.  
*L.*

*HAB.* in Georgiae umbrosis-rivulosis.

MONADELPHIA. POLYANDRIA. 41

STYRAX. L.

*Cal.* 1-phyllus, 5-dentatus. *Cor.* infundibuliformis, 5-fida. *Ovarium* polyspermum. *Drupa* matura 1-2-sperma.

GRANDIFLORUM. *S.* foliis lato-obovalibus, acuminatis, solito denticulatis, subtus tomentosis: racemis multifloris, elongatis.

*S. officinale*? WALT. *Car.* 140.

*HAB.* in sylvis Carolinæ.

PULVERULENTUM. *S.* foliis parvulis, subsessilibus, ovalibus, non acuminatis, subtus quasi pulverulento-tomentosis: floribus axillaribus et subterno-terminalibus, brevissime pedicellatis.

*Obs.* Affine *S. grandiflora*.

*HAB.* in sylvis Carolinæ.

GLABRUM. *S.* foliis parvulis, ovali-lanceolatis, utrinque acutis, glabellis.

*STYRAX americanum*. RICHARD. *Dissert. ms.* in academia scientiarum Parisina lecta anno 1781: in qua, characterem generis reformatum proponens, ovarium semiinferum et polyspermum statuit.

CAVAN. *Dissert.* vi. p. 50. t. 188. f. 1.

*S. laevigatum*. Hort. Kew. 11. 75.

*S. laeve*. WALT. *Car.* 140.

*Obs.* Numerus staminum variat 6-10.

*HAB.* in uliginosis Carolinæ et Georgiæ.

*POLYANDRIA*.

*HALESIA*. L.

*Cal.* 4-dentatus, superus *Cor.* 4-fida. *Nux* quadrangularis, 2-sperma.

N°. A foliis nulla recte eruitur specierum differentia, neque a glandulis petiolaribus.

PARVIFLORA. H. fructu minore, deorsum promisse angustato, inde quasi clavato, inæqualiter subtetraptero : floribus parvulis.

*HAB.* in Florida, circa *Matança*.

TETRAPTERA. H. fructu subæqualiter rhomboideo-  
L. tetragonoptero : floribus majoribus.

*HAB.* in sylvis umbrosis et rivulosis Carolinae.

DIPTERA. H. fructu subelliptico, quasi diptero.  
L.

*HAB.* in Georgiae umbrosis-rivulosis.

MONADELPHIA. POLYANDRIA. 41

STYRAX. L.

*Cal.* 1-phyllus, 5-dentatus. *Cor.* infundibuliformis, 5-fida. *Ovarium* polyspermum. *Drupa* matura 1-2-sperma.

GRANDIFLORUM. *S.* foliis lato-obovalibus, acuminatis, solito denticulatis, subtus tomentosis: racemis multifloris, elongatis.

*S. officinale*? WALT. *Car.* 140.

*HAB.* in sylvis Carolinæ.

PULVERULENTUM. *S.* foliis parvulis, subsessilibus, ovalibus, non acuminatis, subtus, quasi pulverulento-tomentosis: floribus axillaribus et subterno-terminalibus, brevissime pedicellatis.

*OBS.* Affine *S. grandiflora*.

*HAB.* in sylvis Carolinæ.

GLABRUM. *S.* foliis parvulis, ovali-lanceolatis, utrinque acutis, glabellis.

*STYRAX americanum*. RICHARD. *Dissert. mss.* in academia scientiarum Parisina lecta anno 1781: in qua, characterem generis reformatum proponens, ovarium semiinferum et polyspermum statuit.

CAVAN. *Dissert. vi.* p. 50. t. 188. f. 1.

*S. laevigatum*. Hort. Kew. 11. 75.

*S. laevis*. WALT. *Car.* 140.

*OBS.* Numerus staminum variat 6-10.

*HAB.* in uliginosis Carolinæ et Georgiæ.

44 MONADELPHIA. *POLYANDRIA.*

libus vel oblongis : floribus pusillis ;  
breviuscule pedicellatis.

*HAB.* in maritimis Carolinæ.

ALCÆOIDES. *S.* erecta, herbacea : foliis inferioribus triangulato-cordatis, incis; superioribus palmato-multifidis : corymbo terminali : calycibus hispidis.

*OBS.* Habitus quodammodo *MALVÆ Alceæ* vel *moschatae*.

*HAB.* in glareosis Kentucky et Tennassée.

M A L V A. *L.*

*Cal.* duplex : exterior 3-phyllus. Capsulæ plures, evalves, 1-spermæ.

CAROLINIANA. *M.* caule repente : foliis lobato-multifidis : pedunculis solitariis, longis : corollis obsolete rubris : fructibus hirsutis, aristatis.

*HAB.* in Carolina et Virginia.

HIBISCUS. L.

*Cal.* duplex : exterior polyphyllus. *Caps.* 5-locularis, polysperma.

MANIHOT. H. caule petiolisque inermibus : foliis  
L. lineari-palmatis : pedunculis hispidis :  
calyce exteriori 1-phylo ; interiore hinc  
lacero : corolla amplissima , sulphurea :  
capsula hirsutissima , pyramidata , acu-  
minata.

*HAB.* in paludosis juxta *Mississipi*.

SCABER. H. caule scabro : foliis asperis , basi  
quasi truncatis , circumscriptione subro-  
tundis ; superioribus palmatis , lobis su-  
perne dilatatis , crenatis : floribus sub-  
sessilibus ; calycibus hispidissimis , exte-  
riore furcellato.

*OBS.* Folia quasi *URENÆ*. Calyces rigidi ; nervis sub-  
aculeatis. Corolla amplissima , sulphurea , fundo  
atro-purpurea. Capsula ovoideo-pentagona.

*H. aculeatus ? WALT.*

*HAB.* in maritimis Carolinæ et Floridæ.

HASTATUS. H. glaberrimus : foliis hastatis , ser-  
rulatis : corolla tubulato-campanulata ,  
carnea , majuscula : capsula glabra ,



46 MONADELPHIA. POLYANDRIA.

oblongo-ovoidea, acuminata : seminibus holosericeis.

*H. virginicus.* WALT.

*HAB.* ad ripas fluviorum *Ohio*, *Mississipi* et am-  
nium *Carolinæ*.

VIRGINICUS. *H.* universe tomentosus et asper :

- L.* foliis cordatis, superne angustatis, in-  
ferne subangulosis : ramulis quandoque  
subracemifloris : floribus parvulis ; co-  
rolla rosea, patula ; genitalibus decli-  
natis : capsula depresso-rotunda ; lo-  
culis submonospermis.

*H. clypeatus?* WALT.

*HAB.* in paludosis maritimis *Virginiae* et *Carolinæ*.

GRANDIFLORUS. *H.* foliis utrinque eximie tomen-  
tosis, subtus incanis, subcoriaceis, am-  
plis, quasi triangulato-cordatis, tri-  
lobis : corolla omnium amplissima : cap-  
sula subtomentoso-hirsutissima, subtrun-  
cata.

*Obs.* Affinis *H. clypeato* : a quo discrepat præprimis  
corolla multoties ampliore, regulari, dilute carnea :  
foliis crassius tomentosus vel holosericeis. Corollæ  
fundus ruber ; stamina stigmataque lutea.

*HAB.* in maritimis *Georgiæ* et *Floridæ* et in regione  
*Natchez* ad *Mississipi*.

MONADELPHIA. *POLYANDRIA*. 47

**SPECIOSUS.** H. caule lævi, foliis calycibusque glabris : foliis profunde palmatis; laciniis longo-lanceolatis, serratis : corolla ampla, purpurea, patula : capsula glabra, pentagono-ovoidea, subacuta; seminibus minutim tomentosis.

*H. coccineus.* BART. WALT.

*HAB.* in Floridæ scirposis.

**MOSCHEUTOS.** H. foliis ovalibus, acuminatis, dentatis, subtus incano-tomentosis; nonnullis subtricuspidatis : pedunculis quasi petiolis innatis : corolla amplissima : capsula extus glabra.

*Obs.* Corolla subcampanulata, albida; fando purpureo : staminibus pistillisque albis.

*ALCEA indica ulmifolia* &c. PLUCK. t. VI. *fig.* 3.

*HAB.* a Nova Angliæ ad Carolinam.

**CLASSIS XVII.**

**DI A D E L P H I A.**

**P E N T A N D R I A.**

**P E T A L O S T E M U M.**

**DALEA. JUSS.**

Herbæ perennes. Stipulæ setacæ : folia imparipinnata, glanduloso-punctata. Spica seu capitulum ; floribus confertis et arcte sessilibus ; bracteis totidem squamiformibus bracteolisque binis singulos calyces stipantibus.

**CAL.** inferne turbinatus ; striatus ; glandulosus ; incurvus ; 5-fidus seu 5-partitus : laciniis subulatis , conniventibus.

**COR.** petala 5 , vage exerta , filamentose unguiculata : quatuor filamentis staminum interjecta , limbo oblongo-ovali ; quintum superius , vexilli sedem occupans , a cæteris segregatum , imo calyci insertum , limbo cordato et conduplicato-canaliculato.

**STAM.** 5 : filamenta setacea , recta , una cum unguiculis subconsimilibus petalo-



*POLYANDRIA*.

*HALESIA*. L.

*Cal.* 4-dentatus, superus *Cor.* 4-fida. *Nux* quadrangularis, 2-sperma.

N°. A foliis nulla recte eruitur specierum differentia, neque a glandulis petiolaribus.

PARVIFLORA. *H.* fructu minore, deorsum promisse angustato, inde quasi clavato, inæqualiter subtetraptero : floribus parvulis.

*HAB.* in Florida, circa *Matança*.

TETRAPTERA. *H.* fructu subæqualiter rhomboideo-  
- L. tetragonoptero : floribus majoribus.

*HAB.* in sylvis umbrosis et rivulosis Carolinae.

DIPTERA. *H.* fructu subelliptico, quasi diptero.  
L.

*HAB.* in Georgiae umbrosis-rivulosis.

MONADELPHIA. POLYANDRIA. 41

STYRAX. L.

*Cal.* 1-phyllus, 5-dentatus. *Cor.* infundibuliformis, 5-fida. *Ovarium* polyspermum. *Drupa* matura 1-2-sperma.

GRANDIFLORUM. *S.* foliis lato-obovalibus, acuminatis, solito denticulatis, subtus tomentosis: racemismultifloris, elongatis.

*S. officinale?* WALT. *Car.* 140.

*HAB.* in sylvis Carolinæ.

PULVERULENTUM. *S.* foliis parvulis, subsessilibus, ovalibus, non acuminatis, subtus quasi pulverulento-tomentosis: floribus axillaribus et subterno-terminalibus, brevissime pedicellatis.

*Obs.* Affine *S. grandiflora*.

*HAB.* in sylvis Carolinæ.

GLABRUM. *S.* foliis parvulis, ovali-lanceolatis, utrinque acutis, glabellis.

*STYRAX americanum.* RICHARD. *Dissert. mss.* in academia scientiarum Parisina lecta anno 1781: in qua, characterem generis reformatum proponens, ovarium semiinferum et polyspermum statuit.

CAVAN. *Dissert.* VI. p. 50. t. 188. f. 1.

*S. laevigatum.* Hort. Kew. 11. 75.

*S. laeve.* WALT. *Car.* 140.

*Obs.* Numerus staminum variat 6-10.

*HAB.* in uliginosis Carolinæ et Georgiæ.

## O C T A N D R I A.

## P O L Y G A L A. L.

*Cal.* 5-phyllus; foliolis duobus alæformibus, coloratis. *Legumen* obcordatum, biloculare.

INCARNATA. P. caule herbaceo, simplici: foliis

L. rariter sparsis, subulatis: spicis ovali-oblongis: corollis cristatis; tubo gracili, longissime exerto.

*HAB.* in Virginia, Carolina.

SETACEA. P. caule setaceo, aphylo, simplici, summitate subramoso: floribus minutis, incarnatis, imberbibus, dense spicatis.

*Obs.* Caules erecti, angulosi, summitate in paucos ramulos simplices, setaceos, elongatos divisi; quasi aphylli: setulae brevissimæ, acutæ, rariter sparsæ.

*HAB.* in Carolina septentrionali.

SANGUINEA. P. caule summitate ramoso: foliis

L. alternis, linearibus: spicis capitatis; floribus imberbibus, roseis.

*Obs.* Caulis interdum simplicissimus, 1-stachyus. Rami hujus apicem superantes. Cur Linnæus *sanguineam* dixerit nescio.

*HAB.* in Virginia et Carolina.

CRUCIATA. P. caule erecto, ramoso, subalato-

L. anguloso: foliis quaterno-verticillatis,

**DIADELPHIA. OCTANDRIA. 53**

lanceolatis : capitulis solitarie terminabilibus ; sessilibus ; floribus imberbibus , viridulo-carneis.

*HAB.* a Nova Anglia ad Virginiam.

**VERTICILLATA.** P. caule erecto , ramoso : foliis verticillatis , linearibus : spicis setaceis , pedunculatis , oblongis ; floribus albidis , imberbibus , proxime sed distincte alternis.

*HAB.* a Pensylvania ad Carolinam.

**SENEGA.** P. caulibus erectis , simplicissimis , foliosis : foliis alternis , lanceolatis : spica oblonga ; floribus imberbibus , alternis.

Var. *α. albida* : foliis lanceolatis , interdum etiam ovalibus : spica minus laxa ; floribus albidis , subsessilibus.

— *β. rosea* : glabella vel pubens ; foliis lineari-lanceolatis : spica laxiuscule alterniflora ; floribus roseis.

*HAB. α.* in Canada et in montibus *Alleghanis*.

*β.* in Carolina et Georgia.

**UNIFLORA.** P. herbacea , pumila : foliis latis , ovalibus , in petiolum angustatis : floribus imberbibus , solitarie sparsis , pedicellatis.

*HAB.* in finitimis Canadae , ad Lacum *Ontario*.



54     **DIADELPHIA. OCTANDRIA.**

**LUTEA.**     **P.** foliis radicalibus infimisque spatulatis ; cæteris lanceolatis : capitulo promisse pedunculato ; floribus imberbibus, luteis.

Var. *α. elatior* : caule subvirgatim elatiore.

— *β. nana* : subacaulis ; foliis omnibus spatulatis ; capitulo majore.

**HAB. α.** a Marylandia ad Floridam.

— *β.* in pratis udis Carolinæ.

**CORYMBOSA.**     **P.** caule inferne simplicissimo, summitate ramosa corymbose multiflora : floribus luteis, imberbibus : foliis linearibus.

*P. cymosa.* **WALT.**

**HAB.** in paludosis, a Carolina ad Floridam.

**D. E C A N D R I A.**

**C R O T A L A R I A. L.**

*Legumen* turgidum, inflatum, pedicellatum. *Filamenta* connata cum fissura dorsali.

**SAGITTALIS.** C. hirsuta : foliis simplicibus : caule  
**L.** stipulis decurrentibus alato : pedunculis  
 elongatis, subtrifloris : legumine obtu-  
 sissimo.

Var. *α. linearis* : foliis linearibus, caule  
 erecto.

— *β. oblonga* : foliis ovali-oblongis : caule  
 erecto.

— *γ. ovalis* : foliis subelliptico-ovalibus,  
 caule procumbente.

**HAB.** in Virginia et Carolina.

**L U P I N U S. L.**

**Cal.** 2-labiatus. *Antheræ* 5 oblongæ; 5 subro-  
 tundæ. *Legumen* coriaceum.

**PERENNIS.** L. radice perenni, repente : caule fo-  
**L.** liisque glabriusculis : spica alterniflora  
 nec verticillata; floribus cœruleis : calyce  
 laciniis lateralibus nullis : seminibus va-  
 riegatis.

**HAB.** a Canada ad Floridam, in locis a mare remotis.

56      **DIADELPHIA. DECANDRIA.**

**PILOSUS.**    **L.** perennis, vilosissimus : foliis sim-

**L.**    plicibus, oblongis : spica longa, inordinate alterniflora : calyce absque laciniis lateralibus : seminibus variegatis.

**HAB.** a Carolina ad Floridam.

**D A L E A.** **L.** *Hort. Cliff.*

Leguminosa, herbacea : folia glanduloso-puncticulosa, impari-pinnata : flores spicati, bracteis unifloris.

**CAL.** glandulosus ; subsemi 5-fidus ; laciniis subæqualibus, subulatis ; inferne subovoideo-tubulatus.

**COR.** insolite papilionacea : vexillum calyce paulo longius ; lamina oblonga, compresso-canaliculata, porrecta ; ungue hujus longitudine, posticam calycis parietem premente : petala quatuor, bina alarum binaque carinæ loco, consimilia, supra medietatem vaginæ staminalis inserta, unguiculis brevibus, laminis oblongis, latera staminum prementibus, vexillo paulo brevioribus.

**STAM.** vagina staminalis exterum calycis latus premens, angusto-longa, superius margine fissa, summitate 10-fida ; filamentis erectis, petala subæquantibus ; antheris muricatis.





P. J. Redouté Del.

Phé Sc.

DALEA *Linnaei*.

**DIÆDELPHIA DECANDRIA. 57**

**PIST.** ovarium brevi-stipitatum, ovoideum, superne barbatum : stylus glaber, longitudine staminum : stigma obliquum, ambitu minutissimis glandulis quasi fimbriatum.

**FRUCT.** calyx turbinato-obovatus et connivens includit legumen breviter turbinato-ovatum, modice compressum, stylo oblique aristatum, superne barbatum, tenuissime membranaceum; hinc sutura modice dehiscens.

**SEM. 1**, loculo subconforme, obovato-reniforme.

**LINNÆI. D A L E A.**

*tab. 38.*

**P S O R A L E A. L.**

*Calyx* punctis callosis adpersus, longitudine *Leguminis* 1-spermi.

**CANESCENS.** P. tota canescens, foliis breviter petiolatis, trifoliatis : spicis laxifloris; floribus pedicellatis; calycibus villosissimis.

**OBS.** Habitus quodammodo *CYTISI Cajan.* : folia, bracteæ et calyces pellucido-punctati : legumina in acumen ensiforme rectumque desinentia.

**HAB.** in Carolina et Georgia.

58      **DIADELPHIA. DECANDRIA.**

**LUPINELLUS.** P. foliis quino-digitatis; foliolis angustissime linearibus : spica pauciflora; floribus parvulis : leguminibus crassiusculis, ovoideis, uncinato-mucronatis, nervoso-rugosis.

*HAB.* in Carolinæ locis aridis.

**MELILOTOIDES.** P. herbacea, parce minuteque pubescens : foliis lanceolato-trifoliatis : spica oblonga; bracteis latis, acuminatis : leguminibus abbreviato-rotundatis, nervoso-rugosissimis.

*TRIFOLIUM psoralioides. WALT.*

*OBS.* Flores cœrulei.

*HAB.* in Carolina et Florida.

**TRIFOLIUM. L.**

*Flores* subcapitati. *Legumen* vix calyce longius, non dehiscens, deciduum.

**CAROLINIANUM.** T. pusillum, erectum : foliolis rotundato-obcordatis, omisso margine glabris : capitulis pedunculatis, reflexis, paucifloris; corollis vix exertis, albis : leguminibus subtrispermis.

*OBS.* Perpusillum, modice pubescens : capitula longiuscule pedunculata. Calyx pubens, ultra medietatem 5-fidus; laciniis linearibus, rectis.

*HAB.* in Carolina, circa *Charlstown*.

**DIADELPHIA. DECANDRIA. 59**

**REFLEXUM.** T. procumbens, molliter pubens : ca-

**L.** pitulis multifloris ; floribus pedicellatis,  
demum omnibus reflexis : calycibus 5-par-  
titis : leguminibus sub3-spermis.

*Obs.* Foliola obovalia ; ima obcordata. Stipulæ oblique  
cordatæ. Capitula subrotunda ; floribus majusculis.

*HAB.* in Kentucky, Tennassée et montosa Carolina.

**REPENS.** T. capitulis umbellaribus , legumini-

**L.** bus tetraspermis , caule repente.

*HAB.* in Canada.

**AGRARIUM.** T. spicis ovalibus , imbricatis : vexillis

**L.** deflexis , persistentibus ; calycibus nudis ;  
caule erecto.

*Obs.* Nostrate humilior et pubentior.

*HAB.* in Pennsylvania.

**ARVENSE.** T. spicis villosis , ovalibus ; dentibus

**L.** calycinis setaceis , villosis , æqualibus.

*Obs.* Alæ intus macula sanguinea.

*HAB.* in Virginia.



60      **DIADELPHIA DECANDRIA.**

**MEDICAGO. L.**

*Legumen compressum, cochleatum. Carina corollæ a vexillo deflectens.*

**LUPULINA.** M. spicis ovalibus, leguminibus reniformibus, monospermis; caulibus procumbentibus.

*HAB.* in maritimis Carolinæ, juxta *Charlstown* et in Canada.

**PHASEOLUS. L.**

*Carina cum staminibus styloque spiraliter tortis.*

**TRILOBUS.** P. pubescens : foliolis inferioribus rhomboideo-ovalibus ; superioribus trilobatis : capitulis longe pedunculatis : leguminibus linearibus.

*HAB.* in Carolina.

**HELVOLUS.** P. volubilis, gracilis, hirtellus : foliis oblongo-ovalibus, subdeltoideis, obtusiusculis ; capitulis promissa pedunculatis : leguminibus tereti-linearibus.

*HAB.* a Pennsylvania ad Carolinam. 74.

**PANICULATUS.** P. perennis, volubilis, totus pubescens : foliolis lato-ovalibus impari subcordato : panicula pyramidatim elon-

DIADELPHIA. DECANDRIA. 61

gata, multiflora : leguminibus compressis, lunato-falcatis.

*Obs.* Planta more HEDYSARI tenacissima. Panicula interdum pedalis. Flores paulo minores, quam in *P. vulgari* obsolete purpureo-violacei : alæ porrectæ, vexillo longiores : carina brevissima, contorta, pallide-virens. Legumen subbiunciale : semina compresso-reniformia, nigerrima.

*HAB.* in regione Illinoensi.

ERYTHRINA. L.

*Cal.* 2-lobatus. *Cor.* vexillum longissimum, lanceblatum.

HERBACEA. *E. pumila* : foliis trifoliatis; foliolis subhastatis : spica longissima; floribus linearibus, coccineis.

*HAB.* in Carolina et Florida.

GALACTIA. BROWN.

*Cal.* 4-dentatus, 2-bracteatus. *Cor.* 5-petala, petalis omnibus oblongis; vexillo cæteris latiore, incumbente. *Stigma* obtusum. *Legumen* teres. *Semina* subrotunda.

MOLLIS. *G. densa* mollique pube vestita et subcanescens : foliolis ovalibus : leguminibus villosis.

*HAB.* in Carolina inferiore.

62      **DIADELPHIA. DECANDRIA.**

**GLABELLA.** *G. glabriuscula* : foliolis oblongo-ovalibus, utrinque subemarginatis, obtusis : calycibus glabris.

*ERVUM? volubile. WALT.*

*Obs.* Radix perpendicularis, fusiformis.

*HAB.* in Carolina et Georgia.

**CLITORIA. L.**

*Cor.* supinata : vexillo maximo, patente, alas obumbrante.

**VIRGINIANA.** *C. trifoliata* : calyce bracteis longiore,

*L.* 5-partito ; laciniis subulatis, incurvis, divergentibus : legumine quasi ensiformi.

*Obs.* Valde affinis *C. brasiliense* ; cujus calyx decurtatus intra bractea delitescit.

*HAB.* in Virginia et Florida.

**MARIANA.** *C. trifoliata* : calyce bracteis lineari-

*L.* lanceolatis multoties majore, oblongo-tubuloso, 5-fido : legumine faciebus convexis toruloso : seminibus glutinosis.

*HAB.* a Virginia ad Carolinam.

GLYCINE. L.

*Cal.* 2-labiatus: *Corollae* carina apice vexillum reflectens.

FRUTESCENS. G? sarmentis lignosis : foliis impari-  
L. pinnatis, pubentibus : leguminibus coriaceis.

*Anonymos frutescens.* WALT. Car. 186.

HAB. in Virginia, Carolina, regione Illinoensi.

APIOS. G. radice tuberosa : glabella : foliis pin-  
L. natis ; foliolis 5-7, ovali-lanceolatis, superne angustatis : spicis confertifloris.

Obs. Radix edulis.

HAB. a Virginia ad Carolinam et in regione Illinoensi.

TOMENTOSA. G. caule anguloso : foliis trifoliatis,  
L. rarius simplicibus, nervosis, rugosis, plerumque subrotundis : spicis subsessilibus, brevibus : leguminibus compresso-ovoideis, 2-spermis.

Obs. Caulis confertim pubens. Folia leguminaque subtomentosa.

Var. *α. erecta* : caule erecto ; tomentosior, foliolis sæpe oblongo-ovalibus.

TRIFOLIUM *erectum.* WALT.

— *β. volubilis* : caule volubili ; foliis imis simplicibus, cæteris trifoliatis.

— *γ. monophylla* : caulibus brevissimis ; foliis

DIADELPHIA. *DECANDRIA*.

omnibus unifoliatis sive simplicibus, subreniformi-rotundatis.

*TRIFOLIUM simplicifolium*. WALT.

*HAB.* in Virginia et Carolina.

MONOICA. *G.* caule gracili, retrorsum piloso:

*L.* foliis trifoliatis, tenuissime membranaceis: spicis pendulis; pedunculis squamosis.

*Obs.* Flores sexu variat.

*HAB.* in umbrosis regionis Illinoensis et Carolinae.

AMORPHA. *L.*

*Corollae* vexillum ovatum, concavum. *Alae* o.  
*Carina* o.

FRUTICOSA. *A.* arborescens: foliis petiolatis: ca-

*L.* lycibus nudiusculis, pedicellatis.

*HAB.* in Carolina et Florida.

PUMILA. *A.* humilis, fruticosa: foliis sessilibus:  
spicis incanis, tomentosis.

*Obs.* Flores subsessiles.

*A. herbacea*. WALT.

*HAB.* in Carolina et Georgia.

ROBINIA. L.

*Cal.* 4-fidus. *Legumen* gibbum, elongatum.

PSEUD-ACACIA. R. stipulis spinescentibus : foliis

L. impari-pinnatis : racemis cernuis seu pendulis ; calycis dentibus muticis.

*OBS.* Flores albi.

*HAB.* a Canada ad Carolinam.

HISPIDA. R. caule subinermi, hispidissimo : foliis

L. impari-pinnatis ; foliolis rotundato-ovalibus, mucronatis : racemis axillaribus : calycibus acuminatis.

*OBS.* Flores rosei.

*HAB.* in altissimis montibus Carolinæ.

VISCOSA. R. ramis, petiolis pedunculisque glanduloso-viscosis : foliis impari-pinnatis :

racemis axillaribus, confertifloris ; calycibus acuminatis.

*OBS.* Flores roseo-albi.

*R. montana.* BART.

*HAB.* in altissimis montibus Carolinæ.

## LATHYRUS. L.

*Stylus planus, supra villosus, superne latior. Calycis laciniae superiores duo breviores.*

**PALUSTRIS.** L. caule alato : stipulis lineari-lanceo-  
**L.** latis : petiolis apteris ; foliolis sex, lanceolatis : pedunculis sub3-floris.

*Obs.* Flores purpurei.

*HAB.* in septentrionalibus Canadæ.

## ASTRAGALUS. L.

*Legumen biloculare, gibbum.*

**GLABER.** A. caule erecto, sulcato - anguloso,  
 \* glabro : foliolis lanceolato - oblongis ;  
 spicis longe pedunculatis ; floribus exalbidis, laxiusculis ; leguminibus distantibus, incurvis, teretibus.

*Obs.* Foliola subciliata : legumina glabra, oblonga, tumida, juxta suturas depressa.

*HAB.* in sabulosis aridis Georgiæ.

**CAROLINIANUS.** A. elatius, caulescens, erectus :  
**L.** foliolis ovali - oblongis : spica floribus laxiusculis, pallidis ; leguminibus incumbenti-erectis, oblongo - teretibus, glabris.

*HAB.* in altis montibus Carolinæ.

**SECUNDUS.** A. caulescens, procumbens : spicis pe-

**DIADELPHIA. DECANDRIA. 67**

dunculosus ; leguminibus secundis, pendulis, nigricantibus.

*Oss.* Minutim pubescens, foliola ovalia : flores purpurascentes : legumina recta, utrinque longiuscule acuminata.

*HAB.* in septentrionalibus Canadæ.

**VILLOSUS.** A. subacaulis, pilosissimus : foliolis ovalibus : pedunculis folia subæquantibus, brachystachyis : leguminibus incano-villosissimis, assurgentibus, oblongis.

*Oss.* Pumilus, procumbens : foliola rotundato-obtusa seu etiam emarginata : pedunculi subradicales, 3-4-unciales : spicæ paucifloræ ; floribus pallidis : calycis dentibus subulatis.

*HAB.* in Georgia.

**GALEGA. L.**

*Cal.* dentibus subulatis, subæqualibus. *Legumen* striis obliquis, seminibus interjectis.

**VIRGINIANA.** G. herbacea, erecta, tota candicanti-

**L.** villosa : foliolis oblongis : racemo terminali, brevi, subsessili : leguminibus non sulcatis, subfalcatis, molliter villosis.

*Oss.* Affinis præsertim habitu *G. caribææ* !

*HAB.* a Canada ad Floridam.

**VILLOSA.** G. prostrata, villosissima : foliolis cu-



68     **DIADELPHIA. DECAENDRIA.**

neato-obovalibus : spicis suboppositifoliis , pedunculatis , paucifloris : leguminibus rectiusculis , oblongis.

**OBS.** Variat villis albis et rufidulis.

**HAB.** a Carolina ad Floridam.

**HISPIDULA.** *G. gracilis* , humifusa , pubescens : foliolis parvulis , oblongo-ovalibus : pedunculis elongatis , subtrifloris : leguminibus oblongis , falcatis , hispidulis.

**OBS.** Affinis *G. littorali*; diversa vero florum raritate et leguminum forma hirsutietque. Huc forsitan *Onobrychidis* species foliis amplis , glabris , &c. **CLAYT.**

*F. virg.*

**HAB.** in Virginia, Carolina et Georgia.

**INDIGO FERA. L.**

*Cal.* patens. *Corollae* carina utrinque calcari subulato , patulo ! *Legumen* lineare.

**CAROLINIANA.** *I. caule herbaceo* , erecto : foliis

**WALT.** pinnatis; foliolis obovalibus : spicis foliolongioribus , pedunculatis; floribus distantibus : leguminibus pendulis subdispermis , reticulato-venosis. .

**HAB.** a Carolina ad Floridam.

V I C I A. L.

*Stigma latere inferiore transverse barbatum.*

SATIVA ? V. foliolis retusis : floribus binis, subsessilibus : leguminibus erectis, subteretilinearibus, glabris.

*V. sativa. WAT.*

*HAB.* in Carolina inferiore.

CRACCA. V. stipulis integris : foliolis lanceolatis, oblongis, pubescentibus : spicis multifloris, retrorsum subimbricatis : floribus coeruleis.

*HAB.* in Canada.

PARVIFLORA. V. petiolis multifoliatis ; foliolis lineari-lanceolatis, obtusis : spicis multifloris ; floribus parvis, albis : calyce brevissime obtuseque dentato.

*HAB.* in montibus altis Carolinae et Virginiae.

**LESPEDEZA.**

Caulis fruticulosus : folia trifoliata aut raro simplicia.

CAL. 5-partitus ; laciniis subæqualibus ,  
lineari-lanceolatis seu subulatis.

COR. }  
STAM. } **HEDYSARI.**

PIST. ovarium breviter stipitatum ; compresso-subovatum : stylus capillaris : stigma conoideo-capitatum.

LEG. pusillum , lenticulare , monospermum.

**D. LESPEDEZ**, gubernator Floridæ , erga me peregrinatorem officiosissimus.

**SESSILIFLORA.** L. erecta : foliolis oblongis : fasciculis florum sessilibus , numerosis : leguminibus calyce minuto subnudatis , acutis.

**HEDYSARUM junceum.** WALT.

**MEDICAGO virginica.** LINN.

**OBS.** Variat foliolis latiuscule oblongo-ellipticis et sublinearibus tuncque **HEDYSARO junceo** congeneri subsimilis.

**HAB.** in Virginia et Carolina.

**PROCUMBENS.** L. procumbens , gracilis , universe





P. J. Redouté Del.

Plée Sc.

**LESPEDeza** *procumbens*.





P. J. Redoute Del.

Plée Sc.

LESPEDeza *polystachya*.

**DIADELPHIA. DECANDRIA. 7<sup>a</sup>**

pubens : foliolis ovalibus : pedunculis setaceis, distincte pauciterque spicifloris :  
*tab. 39.* leguminibus calyce minuto nudatis.

*Obs.* Flores purpurei.

Affinis *HEDYSARO violaceo*.

*HAB.* in Virginia et Carolina.

**CAPITATA.** L. erecta : foliis subsessilibus ; foliolis oblongis : capitulis sessiliter conglomerato-terminalibus : legumine intra calycem multo majorem recondito.

*HAB.* in Virginia et Carolina.

**POLYSTACHYA.** L. erecta, villosissima : foliolis rotundato-ovalibus : spicis pedunculatis :  
*tab. 40.* corolla calyci subæquali : legumine calycem subæquante.

*HEDYSARUM hirtum. LINN.*

*HAB.* in Carolina et Georgia.

**HEDYSARUM. L.**

*Cor.* carina transverse obtusa. *Legumen* articulis 1-spermis.

**NUDIFLORUM.** H. ovali-trifoliatum : pedunculis  
L. subradicalibus, supra cauliculos procumbentes assurgentibus, nudis, paniculatis : leguminibus hinc profundissime incisis, articulis subtriangulis.

*Obs.* Foliola lato-ovalia, acuminata, subtus subalbi-



**DIADELPHIA. DECANDRIA.**

cantia, pube marginata. Datur varietas minor, foliolis suborbiculatis; quam in Kentucky legi.

*HAB.* a Nova Anglia ad Carolinam.

**ACUMINATUM.** H. erectum, simplex, summitate frondosum : foliis trifoliatis, ovalibus, longe acuminatis; impari rotundato-subrhombeo : panicula terminali longissime pedunculata.

*OBS.* Proximum *H. nudifloro*.

*HAB.* a Nova Anglia ad Carolinam.

**LINEATUM.** H. caule repente, viridi-lineato : foliis subsessilibus, trifoliatis, suborbiculatis : racemis elongatis, laxe parvifloris : leguminibus articulis lenticularibus.

*OBS.* Variat foliolis ovalibus.

*HAB.* in Carolina.

**ROTUNDIFOLIUM.** H. caule prostrato petiolisque hirsutis : stipulis rotundato-cordatis, reflexis : foliis trifoliatis, orbiculatis, utrinque pilosis : racemis paucifloris : leguminibus articulis subrhomboideis.

*OBS.* Racemi axillares et paniculato-terminales; floribus paucioribus, pedicellatis, purpureis. Inferior racemorum pars solito alternatim stipulosa.

*HAB.* in Carolina.

**VIRIDIFLORUM.** H. erectum, puberulum : ramis  
*L.* virgatim et composite racemifloris : foliis

**DIADÉLPHIA. DECANDRIA. 73**

trifoliatis , oblongo - ovalibus , sursum angustatis ; stipulis cordatis : leguminibus tenacissimis ; articulis ovalibus.

*Obs.* Canles erecti, superne laxè ramosissimi ; minuto pubentes simulque pilis longioribus hirti. Foliola acuta, subtus reticulato-venosissima. Flores purpurascens, marcescendo subvirides. Leguminum articuli majusculi.

*HAB.* in Virginia et Carolina.

**BRACTEOSUM.** *H.* erectum , glabrum : foliis trifoliatis , oblongo - ovalibus , acuminatis : stipulis subulatis : racemo terminali, sparsifloro : leguminibus articulis sub-ovalibus.

*Obs.* Majusculum. Racemus pedunculatus ; inferno solito ramulis simplicibus, steriliter bracteatis ; bracteis majusculis, ovalibus ; longius acuminatis, multistriatis, glabris. Flores pedicellati, sæpe gemini, purpurei.

*HAB.* in montosis Virginiae et Carolinae.

**GLABELLUM.** *H.* glabriusculum , erectum , foliosum : foliis trifoliatis, apprimè ovalibus, utrinque obtusis, subtus subglaucis : stipulis minutis, subulatis : racemo laxè paucifloro : leguminibus articulis triangulo-subrhombeis.

*Obs.* Flores longiuscule pedicellati, plerumque gemini, purpurei. Affine *H. canescenti*.

*HAB.* in Carolina inferiore.

74     **DIADELPHIA. DECANDRIA.**

**PANICULATUM.** *H. stans* : foliis trifoliatis , lineari-  
**L.**     ribus , glabris : panicula sparsiflora :  
         leguminibus tenacissimis ; articulis sub-  
         ovali-lenticularibus.

**OBS.** Caulis minutissime puberulus. Folia breviter petiolata. Stipulae setaceae.

**HAB.** in Carolina inferiore.

**ALPINUM** : americanum. *H. erectum* : foliis pinnatis :  
         stipulis subvaginantibus : leguminibus  
         patulis , glabris , ex utraque sutura articu-  
         latis ; articulis brevi-ovalibus.

*H. virginicum* ? LINN.

**OBS.** Foliola ovali-oblonga , pilosa. Legumina multo  
         minora quam in Europæo , licet flores magnitudine  
         non cedant.

**HAB.** in borealibus Canadæ et in cataractis montium  
         *Alléghanis*.

**ÆSCHYNOMENE. L.**

**Cal.** bilabiatus. *Legumen* articulis truncatis , monospermis.

**OBS.** Genus in posterum dividendum in duo  
         genera distinctissima , *SESBAN* scilicet et  
         *ÆSCHYNOMENE*.

**VISCIDULA.** *Æ. caule* prostrato , gracili ; foliolis  
         7-9 , obovalibus : pedunculis subbifloris :

**DIADELPHIA. DECANDRIA. 75**

legumine pubente, profunda incisura articulato.

*Obs.* Caulis brevissima, densa viscidulaque pube vestitus.

*HAB.* in arenosis insulæ *Cumberland* et *Floridæ*.

**PLATYCARPA.** *Æ.* foliolis passim ciliatis : legumine lato-lanceolato, longius stipitato, subdispermo.

*Obs.* Certo *SESBANIS* species; licet leg. 2  $\frac{1}{2}$  unc. longo et 6-7 lin. lato multum discrepat. Affinis *Æ. virgata*, *CAVAN. Icon. tab. 293*, et *ROBINIA vesicaria*, *JAC. Misc. 105*.

*HAB.* in humidis *Carolinæ*, usque ad *Floridam*. ☉.

**STYLOSANTHES. SWARTZ.**

*Cal.* tubulosus, longissimus, corollifer. *Ovarium* infra corollam. *Legumen* 2-articulatum, hamatum.

**HISPIDA.** *S.* procumbens : stipulis foliisque at saltem superioribus hispidis : capitulis 2-3-floris.

*TRIFOLIUM biflorum. LINN.*

Var.  $\alpha$ . *nudiuscula* ; caule foliisque inferioribus glabellis.

—  $\beta$ . *hispidissima* : universe hispidissima.

*HAB.* in *Virginia* et *Carolina*.

## ZORNIA. GMEL.

*Cal.* campanulatus, 2-labiatus. *Cor.* inféra. *Vexillum* cordatum, revolutum. *Antheræ* alternæ oblongæ, alternæ globosæ. *Legumen* articulatum, hispidum.

TETRAPHYLLA. *Z.* foliis digitato-quadrifoliatis :  
*tab. 41.* bracteis suborbiculatis.

*Z. bracteata.* GMEL. *Syst.* 1096.

*Obs.* Congener HEDYSARUM *diphyllum.* LINN.

*HAB.* in Carolina inferiore.

---



P. J. Redouté Del.

Plac. Sc.

*ZORNIA tetraphylla.*



CLASSIS XVIII.

POLYADELPHIA.

POLYANDRIA.

ASCYRUM. L.

*Cal.* 4-phyllus. *Petala* 4. *Filamenta* multa, in 4  
phalanges digesta.

PUMILUM. *A. humillimum* : foliis parvulis , ova-  
libus , obtusis : pedicellis longis , re-  
flexis : floribus 1-2-gynis.

*HAB.* in Georgia.

MULTICAULE. *A. multicaule*, erectiusculum : foliis  
sublanceolato-oblongis : floribus subses-  
silibus , digynis.

*HAB.* in Virginia, Carolina.

STANS. *A. caule erecto*, inferne simplici , an-  
cipite : foliis oblongo-ovalibus : floribus  
breviter pedicellatis , trigynis.

*A. hypericoides* ? WALT.

*HAB.* in Carolina.

AMPLEXICAULE. *A. caule erecto*, superne dichotome  
paniculato : foliis amplexicaulibus,  
cordatis : floribus trigynis.

*HAB.* in Florida.



HYPERICUM. L.

*Cal.* 5 - partitus. *Petala* 5. *Filamenta* multa,  
in 5 phalanges basi connata. *Capsula*.

**GLAUCUM.** H. caule fruticoso, cylindrico : foliis  
cordatis, amplexicaulibus, lævigatis,  
glaucis : panicula divaricato-dichotoma,  
foliosa.

*Oss.* Folia sinu profundo cordata, coriacea. Flores ma-  
jusculi; staminibus numerosissimis, petala sub-  
æquantibus.

*HAB.* in Florida.

**NUDIFLORUM.** H. frutescens, elatius, glabrum :  
caule alato-4-gono : foliis ovali-ōblon-  
gis : panicula pedunculata, dichotoma,  
aphylla; floribus monogynis.

*Oss.* Folia obtusiuscula; calyx post florationem pa-  
tulus : capsula ovoidea, stylo tripartibili.

*HAB.* in Carolina.

**SPHÆROCARPUM.** H. herbaceum, glaberrimum,  
erectum : foliis oblongis : panicula nuda,  
dichotoma; dichotomiis omnibus 1-flo-  
ris : stylo unico, demum tripartibili :  
capsula globosa.

*HAB.* in Kentucky.

**ANGULOSUM.** H. herbaceum, erectum, quadran-  
gulum : foliis lanceolato-ovalibus, acutis;

POLYADELPHIA. *POLYANDRIA*. 79

panicula dichotoma , distanter alterniflora : calyce inferne anguloso.

*H. denticulatum* ? WALT.

*Obs.* *HYPERICO canadensi* paulo majus. Folia arcte sessilia , erectiuscula , 5-nervia. Flores pusilli ; calyce subcampanulato ; foliolis oblongis , inferne prominenti-carinatis.

*HAB.* in paludosis Carolinæ.

**SAROTHTA.** *H. pusillum* , erectum , setaceo-ramosissimum , quasi aphyllum : floribus in ramulis solitarie alternis , 5-10-andris , 3-stylis.

*SAROTHTA gentianoides*. LINN.

*HYPERICUM nudicaule*. WALT.

*Obs.* Folia minuta , appressa , pellucido-punctata ( congenerum more ). Capsula oblonga , 1-locularis ; juxta suturas valvarum oligosperma.

*HAB.* in apricis New-Yorck , Virginiae , Carolinæ , &c.

**CANADENSE.** *H. pusillum* , erectum , 4-gulum , parviflorum : foliis lineari-lanceolatis : paniculæ ramis primariis oppositis , secundariis dichotomis : capsula longa , conoidea.

*HAB.* a sinu *Hudsonis* ad Pensylvaniam et in excelsis montibus Carolinæ.

**QUINQUENERVIUM.** *H. gracile* , decumbens , quadrangulum , glabrum : foliis subamplexicaulibus , oblongo-ovalibus , obtusis , 5-

80 POLYADELPHIA. *POLYANDRIA*.

nerviis : paniculæ multifloræ ramis terminalibus subquaternis.

*HAB.* in graminosis palustribus Carolinæ.

**SIMPLEX.** *H.* annuum, minute lanulosum : caulibus simplicissimis, erectis : foliis oblongis : floribus paucis, trigynis, terminalibus.

*ASCYRUM villosum.* LINN.

*HYPERICUM pilosum.* WALT.

*OBS.* Parvulum, lana rufescente vestitum. Folia sessilia, subappressa, erecta, canaliculata.

*HAB.* in Carolina inferiore.

**MACULATUM.** *H.* herbaceum, erectum, teres, glabrum : foliis floribusque nigro-punctatis : foliis amplexicaulibus, oblongo-ovalibus, obtusis : panicula brachiata, fasciculato-multiflora.

*OBS.* Affine *H. montano*.

*HAB.* in Virginia et in montibus Carolinæ.

**FASCICULATUM.** *H.* fruticosum, erectum, opposite ramosum, glabrum : foliis linearibus, obtusis, convexis, passim fasciculatis ; calycibus foliiformibus.

*OBS.* Flores in ramulis lateralibus foliose congesti, Affine *H. prolifico*.

*HAB.* in Carolina.

**POLYADELPHIA. POLYANDRIA. 8t**

**VIRGINICUM.** H. herbaceum, erectum, glabrum:

**L.** foliis amplexicaulibus, cordato-oblongis, obtusissimis: fasciculis pedunculatis, paucifloris.

**OBS.** Flores triadelpho-enneandri.

**HAB.** in Virginia et Carolina superiore.

**AXILLARE.** H. herbaceum, erectum, glabrum, oppositè ramosum: foliis petiolatis, oblongo-obovalibus, rotundato-obtusis: floribus oppositè axillaribus, subsessilibus, subternis: capsula oblonga.

*H. campanulatum?* WALT.

**HAB.** in aquis stagnantibus Carolinæ.

**FRONDOSUM.** H. frutescens, glaberrimum, ramis ancipitibus: foliis ovali-oblongis, obtusis, basi angustatis: floribus majusculis, subsolitarie terminalibus: calyce foliaceo, petalis subæquali.

**OBS.** Affine *H. Ascyro*. Calycis foliola admodum inæqualia, foliiformia, petalis latiora etiamque interdum longiora.

**HAB.** in rupibus, ad flumen *Tennassée*.

**PROCUMBENS.** H. procumbens, glaberrimum, ramis ancipitibus, summitate paucifloris: foliis linearibus, enerviis; patulis: calyce foliaceo: staminibus corolla brevioribus.

**OBS.** Calyx amplitudine corollæ. Stylus tripartibilis.

**HAB.** in aridis collibus Kentucky.

82 POLYADELPHIA. *POLYANDRIA*.

**MACROCARPUM.** *H. majusculum*, glabrum ; caule ramisque stricte erectis : foliis amplexicaulibus, oblongo-ovalibus, acutis : summitatibus paucifloris : stylis quinque : capsula ovata, maxima.

*Obs.* Perenne : caulis firmus, erectus, 2-pedalis et ultra ; quadrangulatus ; ramis oppositis, strictis, erectis. Corolla calyce multoties amplior ; petalis subrotundo-obovalibus.

*HAB.* in Canada, circa *Montréal*.

---

CLASSIS XIX.

SYNGENESIA.

*POLYGAMIA ÆQUALIS*.

§. I. *LIGULATÆ*.

*PRENANTHES*. *L.*

*Receptaculum* nudum. *Cal.* calyculatus. *Pappus* simplex, subsessilis. *Flosculi* simplici serie.

*ALBA.* *L.* *P.* foliis subhastato-angulosis vel lobatis : panicula laxa, e fascioulis terminalibus, cernuis : calycibus 8-fidis, 9-10-floris, floribus albidis.

*HAB.* in Virginia et Carolina.

*VIRGATA.* *P.* caule simplici, in spicam virgatam longissimam desinente; floribus subsecundis, purpurascentibus, pendulis : calycibus glabris, 8-fidis, 12-floris; foliis lyrato-sinuatis.

*PRENANTHES autumnalis*. GRON. WALT.

*HAB.* in Virginia et Carolina.

*ASPERA.* *P.* caule simplici foliisque asperrimis : foliis omnibus indivisis, ovali-lanceola-

84 SYNGENESIA. *POLYG. ÆQUALIS*:

tis ; racemo longo , fasciculis subsessilibus , erectis , hirsutis.

Affinis *P. racemosae*.

*HAB.* in pratis regionis Illinoensis.

**RACEMOSA.** *P.* caule simplici ; foliis omnibus indivisis , lævibus ; radicalibus ovali-lanceolatis ; caulinis semiamplexicaulibus : racemo oblongo , hirsutissimo , fasciculis cernuis : calycibus 8-9-partitis ; 9-12-floris.

*HAB.* in septentrionalibus Canadæ.

**CREPIDINEA.** *P.* foliis lato-lanceolatis , inferne in petiolum angustatis , inæqualiter subangulato-dentatis : panicula fasciculis terminalibus , paucifloris , nutantibus : calycibus hirsutis , 10-12-fidis , calyculatis , sub20-floris.

*HAB.* in regione Illinoensi et in excelsis montibus Carolinæ.

SYNGENESIA. *POLYG. ÆQUALIS.* 85

LACTUCA. *L.*

*Recept.* nudum. *Cal.* imbricatus, cylindricus, margine membranaceo. *Pappus* simplex, stipitatus. *Sem.* lævia.

LONGIFOLIA. *L.* foliis inermibus, amplexicaulibus, sursum angustatis, plerumque indivisis : panicula aphylla, corymbosa.

*Obs.* Folia interdum versus medietatem laciniis 2-4-subruncinata, margine integerrima. Panicula ramosissima; floribus omnibus pedunculatis, lutescentibus.

*HAB.* in Carolina superiore.

GRAMINIFOLIA. *L.* caule erecto, simplici : foliis inermibus, plerisque indivisis, basi simplicibus, longissime linearibus : panicula aphylla, laxa; ramis rarifloris; floribus omnibus pedunculatis.

*Obs.* Folia radicalia paucioribus laciniis subruncinata.

*HAB.* in Carolina inferiore.

SONCHUS. *L.*

*Recept.* nudum. *Cal.* imbricatus, ventricosus. *Pappus* pilosus.

FLORIDANUS. *S.* lævis glaberque : foliis lyrato-subhastatis seu ovali-lanceolatis, utrinque acuminatis : panicula oblonga; pe-



86 SYNGENESIA. *POLYG. ÆQUALIS.*

dunculis squamulosis; floribus parvulis,  
cœruleis.

*HAB.* in utraque Carolina.

HIERACIUM. *L.*

*Recept.* nudum. *Cal.* imbricatus, ovatus. *Pappus*  
simplex, sessilis.

CANADENSE. *H.* caule erecto, simplici, folioso :  
foliis lanceolatis, acutissimis, semiam-  
plexicaulibus, rariter et longiuscule ser-  
ratis; hirsutis : corymbo paucifloro : pe-  
dunculis longis calycibusque hispidulis.

*Obs.* Affine *H. sabauda*.

*HAB.* in Canada.

SCABRUM. *H.* caule erecto, elatiore, folioso, his-  
pido, puncticulis fuscis scaberrimo :  
foliis oblongo-obovalibus, integris, hir-  
sutis : panicula multiflora calycibusque  
glanduloso-hispidulis.

*HAB.* in septentrionalibus Canadæ et in excelsis mon-  
tibus Carolinæ.

PANICULATUM. *H.* caule erecto, folioso, inferne  
*L.* densis longisque villis lanoso : foliis lan-  
ceolatis, rariter denticulatis, glabellis :  
panicula glabra, composita, subcapil-  
lari, floribus minutis.

*HAB.* in montibus Carolinæ.

SYNGENESIA. *POLYG. ÆQUALIS*. 87

GRONOVII. H. caule simplici foliisque pilosissi-

*L.* mis : foliis oblongo-ovalibus , integris :  
panicula parviflora calycibusque glandu-  
loso-hispidulis.

Var. *α. nudicaule* : caule subunifoliato : panicula  
subfastigiata.

— *β. foliosum* : caule parce folioso : panicula  
oblonga.

*HAB.* *β.* in Virginia et Carolina.

— *α.* in Canada et Pennsylvania.

HYOSERIS. *L.*

*Recept.* nudum. *Cal.* subæqualis. *Pappus* pilosus  
calyculatusque.

MONTANA. H. glaberrima , procumbens : foliis  
lanceolatis , integerrimis : scapo unifloro :  
pappo duplici ; paleolis minutis exteriori-  
bus pilisque paucis.

*HAB.* in altissimis montibus Carolinae septentrionalis.

ANGUSTIFOLIA. H. foliis lineari-lanceolatis , sen-  
sim acutissimis , glabellis ; dentibus ra-  
rioribus , exertis : scapo unifloro.

Affinis *H. majori*. *WALT.*

*HAB.* in Carolina.

AMPLEXICAULIS. H. erecta , glabra , glaucescens :  
foliis radicalibus oblongo-ovalibus , sub-  
integris : scapo summitate bracteis am-

88 SYNGENESIA. *POLYG. ÆQUALIS.*

plexicaulibus vestito et inde subtrifloro;  
pedicellis elongatis.

*Obs.* Umbellula sæpius triflora, interdum 5-flora.

*H. biflora?* WALT.

*HAB.* in Pensylvania, *Tennassée* et Carolina.

VIRGINICA. *H. pusilla*; foliis lyratis, glaucis:

*L.* scapo 1-floro: calyce 10-phylo: pappo  
duplici, membranaceo, vix piloso.

*Obs.* Habitus parvuli *Taraxaci*.

*HAB.* a Canada ad Floridam.

LEONTODON. *L.*

*Recept.* nudum. *Cal.* imbricatus, squamis laxius-  
culis. *Pappus* plumosus.

TARAXACUM. *L.* calyce squamis inferne reflexis:

*L.* foliis runcinatis, denticulatis, lævibus.

*HAB.* ad ripas rivulorum ascendente mari inundatas,  
juxta *Charlstown*.

SCORZONERA. L.

*Recept.* nudum. *Cal.* imbricatus squamis margine scariosis. *Pappus* plumosus.

PINNATIFIDA. S. caule erecto, 3-floro; pedunculis elongatis : foliis lanceolatis, acutissimis, acute pinnatifidis; interdum rariter dentatis.

*HAB.* in ruderatis Carolinæ.

§. II. FLOSCULOSÆ.

CIRSIIUM. T. HALL. GÆRT.

*Pappus* plumosus : cætera CARDUI.

*Obs.* Huc CARDUI et CNICI LINN. plures species.

REPANDUM. C. foliis amplexicaulibus, angustoblongis, levissime obtuseque sinuosis, spinulis crebris, lanuginosis : ramis unifloris, foliosis : calycis squamis lanceolatis, erectis, quasi aristatis.

CARDUUS *virgipianus.* WALT.

*HAB.* in Carolina.

MUTICUM. C. elatius : foliis omnibus pinnatifidis; laciniis sublanceolatis, acutis : ramis in summitate pluribus, nudiusculis, uni-

90 SYNGENESIA. *POLYG. ÆQUALIS.*

floris : calyce globoso , squamis muticis.

*Obs.* Folia subtus subtomentoso-lanuginosa , margine spinulosa. Calyx communis lanuginosus.

*HAB.* in altissimis montibus Carolinæ.

**VIRGINIANUM.** *C. simpliciusculum* : foliis lanceolatis , subintegris seu sinuato - acutangulis , margine spinulosis , subtus canotomentosis : floribus solitariis : calyce globoso ; squamis spinula acuminatis.

*CARDUUS virginianus.* JACQ. *Obs.* IV. 13. t. 99.

*HAB.* in Carolina.

**HORRIDULUM.** *C. elatius* : foliis sessilibus , pinnatifidis , acute subdivisis , valide spinosis : involucro terminali , unifloro , polyphyllo , quasi pectinatim spinosissimo.

*Obs.* Affine *CNICO spinosissimo.*

Calyx oblongus ; squamis etiam summitate acutissima et inermi crectis.

*HAB.* in pascuis ruderalisque Carolinæ.

SYNGENESIA. *POLYG. ÆQUALIS.* 91

LIATRIS. SCHREB.

*Anonymos.* WALT. *Car.* 196.

*Pappus* plumosus : cætera CARDUI.

*Obs.* Omnium flores purpurei.

\* *Spicatae seu racemosae : bulbosae.*

MACROSTACHYA. L. caule simplici, elato : foliis longo-linearibus, nitidis, inferne ciliatis : spica longissima, floribus sessilibus : calycibus squamis appressis.

*SERRATULA spicata.* LINN.

*HAB.* in Virginia et Carolina.

PYCNOSTACHYA. L. caule simplici, hirsuto : foliis strictis, angusto-linearibus, pubescentibus : spica longa ; floribus confertim sessilibus : calycibus superne squarrosis.

*HAB.* in pratis Illinoensibus.

ELEGANS. L. caule simplici : foliis subfalcato-linearibus : spica subfoliosa ; floribus breviter pedicellatis : calycum squamis intimis in ligulam petaloideam protractis.

*STÆHELINA elegans.* WALT.

*SERRATULA speciosa.* AIT.

*HAB.* in Carolina et Florida.

92 SYNGENESIA. *POLYG. ÆQUALIS*.

**SQUARROSA.** L. foliis longissime linearibus, margine asperiusculis : spica foliosa ; floribus magnis, paucis : calyce squarroso ; squamis superne foliaceis , lanceolatis , patulis.

*SERRULATA squarrosa.* LINN.

*Obs.* Hujus et *L. elegantis* pappus manifestius plumosus.

*HAB.* in Virginia , Kentucky et Carolina.

**SQUARRULOSA.** L. foliis lanceolatis , margine asperis ; imis multoties majoribus : spica longa ; floribus alternis , stipitatis , distantibus : calycibus inferne squarrosis.

*Obs.* Videtur *SERRATULA scariosa* L. non autem video cur , Pluknetio duce , scariosam dixerit ! Interdum flores sessiles. Calycum squamæ spathulatæ.

**ASPERA.** L. foliis lineari-lanceolatis , asperimis : spica floribus distincte alternis , sessilibus , solitariis : calycibus majusculis , breviusculis ; squamis rotundatis.

*HAB.* in regione Illinoensi.

**SPHEROIDEA.** L. foliis lævibus ; inferioribus lato-lanceolatis ; superioribus lanceolato-linearibus : spica floribus majusculis , solitariis , alternis , stipitatis : calycibus subglobosis ; squamis ovalibus , erectis.

*HAB.* in pratis Illinoensibus et in excelsis montibus Carolinæ.

**SYNGENESIA. *POLYG. ÆQUALIS.* 93**

**CYLINDRACEA.** *L. gracilis*, tota hirsutula : foliis gramineis : spica rariflora : calycibus subsessilibus, cylindraceis, paucifloris; squamis apice rotundatis, abrupte mucronatis.

*HAB.* in pratis sylvisque Illinoensibus.

**\*\* *Paniculatae seu corymbosae : non bulbosae.***

**PANICULATA.** *L. caule*, panicula calycibusque hirsutulis : foliis radicalibus imisque plantagineo-lanceolatis; caulinis parvulis, sessilibus : panicula multifasciculata : calycibus sub5-floris.

*HAB.* a Virginia ad Floridam.

**ODORATISSIMA.** *L. glaberrima* : foliis radicalibus plantagineo-oblongis; caulinis semiamplexicaulibus : corymbo composito, laxo, parvifloro, calycibus sub8-floris.

*HAB.* a Carolina ad Floridam.

**TOMENTOSA.** *L. caule* simplicissimo foliisque cuneato-lanceolatis, hirsutis : corymbo paucifloro, depresso, divaricato; ramis 1-2-floris : calycibus tomentosis; squamis ovalibus, subacutis.

*HAB.* in Carolina septentrionali.

**BELLIDIFOLIA.** *L. pumila*, glabella : foliis cuneato-ob lanceolatis : corymbo inæquali; flori-



94 SYNGENESIA. *POLYG. ÆQUALIS.*

bus omnibus longiuscule pedicellatis ;  
squamis oblongo-obovalibus , obtusis.

*HAB.* in sabulosis aridis Carolinæ septentrionalis.

VERNONIA. SCHREB.

*CHRYSOCOMA.* WALT. *Car.* 196.

*Cal. comm.* imbricatus. *Recept.* nudum , cavo-  
punctatum. *Flosculi* omnes hermaphroditi :  
*Stigma* bifidum. *Pappus* setaceo-pilosus.

*Obs.* In omnibus speciebus hic memoratis flores vio-  
laceo-purpurei.

OLIGOPHYLLA. V. caule simplici, subnudo seu re-  
mote oligophyllo : foliis serratis ; radica-  
libus oblongo-ovalibus ; caulinis lanceo-  
latis : corymbo paniculato.

Var. *α. verna* : utroque flore pedunculato.

— *β. autumnalis* : altero flore sessili.


*HAB.* in Carolina.

ANGUSTIFOLIA. V. foliis crebris , longe angustequae  
linearibus , subintegris : corymbo sub-  
umbellato : calycibus squamis rigidule  
mucronatis.

*HAB.* in aridis apricis sylvarum Carolinæ.

FASCICULATA. V. foliis longo-linearibus , rariter  
serratis : floribus corymbosis , erecto-ap-





P. J. Redoute Del.

Plée. Sc.

**SPARGANOPHORUS** *verticillatus*.

**SYNGENESIA. POLYG. ÆQUALIS. 95**

proximatis : calycibus ovoideis, lævibus ;  
squamis muticis.

*HAB.* in pratis Illinoensibus.

**PRÆALTA.** V. caule altissimo, anguloso, dense pubente : foliis crebris, lanceolatis, acute serratis : calycibus squamis muticis.

*SERRATULA praealta.* LINN.

*HAB.* in Nova Anglia.

**NOVEBORACENSIS.** V. altissima, foliis crebris, longolanceolatis, leviter serratis : calycibus squamis aristatis.

*SERRATULA noveboracensis.* LINN.

*HAB.* in Virginia et Carolina.

**SPARGANOPHORUS. GÆRT.**

*Cal. comm.* subglobosus, imbricatus squamis inæqualibus, apice recurvato-patulis. *Recept.* nudum. *Semen* coronatum cupula subcartilaginea, integerrima, nitida. *Flosculi* omnes androgyni, uniformes.

**VERTICILLATUS.** S. foliis setaceo-linearibus, verticillatis : caulibus summitate subunifloris : pappo campanulato, 5-dentato.

*ETHULIA uniflora?* WALT.

*HAB.* in inundatis, a Carolina ad Floridam.

*CACALIA. L.*

*Recept.* nudum. *Pappus* pilosus. *Cal.* cylindricus, oblongus, basi tantum subcalyculatus.

**ATRIPLICIFOLIA.** *C.* herbacea, elatior, glaberrima,  
*L.* glauca : foliis imis lato-cordatis, sinuato-angulosis ; caulinis subdeltoideis : calycibus 5-floris.

*Obs.* Aliam in regione Illinoensi vidi speciem huic proximam, nullo modo glaucam, &c.

*HAB.* a Pennsylvania ad Carolinam.

**SUAVEOLENS.** *C.* herbacea, elatior, glabella : foliis acute hastato-triangulatis, basi abrupte cuneatim angustatis, serratis : corymbo erecto, multifloro.

*Obs.* Flores albid.

*HAB.* ad ripas fluminis *Ohio.*

SYNGENESIA. *POLYG. ÆQUALIS.* 97.

EUPATORIUM. *L.*

*Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus, oblongus. *Stylus* semibifidus, longus.

SCANDENS. *E.* caule volubili : foliis subhastato-  
*L.* cordatis, acuminatis, repandis dentatisve : corymbis parvulis, fasciculatis, calycibus 4-floris.

*Obs.* Flores purpurascentes.

*HAB.* in humidis, a Canada ad Carolinam.

ALTISSIMUM. *E.* foliis sessilibus, lanceolatis, sub-  
*L.* integris, trinerviis : panicula foliosa ; floribus fasciculatis : calycibus candidi-pubentibus, 5-floris.

*Obs.* Flores albi.

*HAB.* in Pennsylvania.

LINEARIFOLIUM. *E.* parvifolium, pubens : foliis  
*WALT.* sessilibus, lineari-lanceolatis, subintegris, subtus punctatis : calycibus hirsutulis, subacutis, 5-floris.

*Obs.* Caulis corymbique pubentes. Folia plerumque integra, nonnunquam dentata; modo lanceolata; modo angusto-linearia et passim fasciculata. Flores albi : calycis squamæ lineares, subacutæ, atomis quadrangularibus adpersæ.

*HAB.* in inundatis æstate exsiccabilibus Carolinæ.

98 SYNGENESIA. *POLYG. ÆQUALIS.*

SESSILIFOLIUM. E. gracile , glabellum : foliis sub-

- L. amplexanti-sessilibus , longo-lanceolatis ,  
subæqualiter serratis : calycibus 5-floris.

*Obs.* Flores albi.

*HAB.* in occidentalibus montibus *Allèghanis.*

ROTUNDIFOLIUM. E. foliis sessilibus , breviter et

- L. subrotunde cordato-ovoideis , crebro et  
subæqualiter dentatis , supra asperis : co-  
rymbo fastigiato : calycibus acuminatis ,  
5-floris.

*Obs.* Flores albi.

*HAB.* in Carolina.

VERBENÆFOLIUM. E. foliis sessilibus , sublanceo-  
lato-ovalibus , acutis , obtusiuscule in-  
ciso-dentatis , pube asperiusculis : caly-  
cibus pubentibus , muticis , 5-floris.

*Obs.* Folia erecta , e latiore basi sensim acuta nec acu-  
minata ; superioribus *VERBENÆ officinalis* subsi-  
milia. Corymbus compositus , pubens ; floribus par-  
vulis , albis , glomerato-fasciculatis.

*HAB.* in humidis Carolinæ.

GLANDULOSUM. E. hirsutulum : foliis sessilibus ,  
longo - lanceolatis , serratis , asperis ,  
punctatis : calycibus 5 - floris , creber-  
rime glanduloso-punctatis , lineari-subu-  
latis , superne coloratis.

*Obs.* Herba habitusque fere *VERBENÆ bonariensis.*

*HAB.* in aridis sylvarum Carolinæ.

SYNGENESIA. *POLYG. ÆQUALIS*. 99

**FALCATUM.** *E. glabellum* : foliis 4 - verticillatis ,  
ovalilanceolatis , utrinque acuminatis ,  
subfalcatis , rariter serratis : calycibus  
5-floris , gradatim imbricatis , squamis  
extimis brevibus , ovalibus.

*OBS.* Affine *E. albo* ; præaltum : panicula longius  
brachiata , multicorymbosa. Flores albi.

*HAB.* ad ripas fluviorum *Ohio* et *Scioto*.

**PURPUREUM.** *E. altissimum* ; caule glabro : foliis  
3-4-verticillatis , lato-lanceolatis , utrin-  
que subæqualiter angustatis , rugosis ,  
serratis : calycibus sub8-floris.

*OBS.* Caulis glauco-purpureus. Flores purpurei ; ca-  
lyces 6-9-flori.

*HAB.* in *Canada*.

**MACULATUM.** *E. altissimum* ; caule pulverulento ,  
*L.* saturatius lineolato : foliis 3-5-verticil-  
latis , petiolatis , ovalibus , acuminatis ,  
serratis , rugosis , basi subtrinerviis : ca-  
lycibus sub8-floris.

*OBS.* Valde affine *E. purpureo*.

*HAB.* in *Canada* , montibus *Alléghanis* aliisque regio-  
nibus occidentalibus.

**CONNATUM.** *E. caule lanuloso* : foliis connatis ,  
oblongis , sursum angustatis , serratis ,  
rugosis , subtus subtomentosis : calyci-  
bus 10-floris et ultra : floribus albis.

*E. perfoliatum*. LINN.

*HAB.* a *Canada* ad *Floridam*.



100 SYNGENESIA. *POLYG. ÆQUALIS.*

SEROTINUM. E. caule pulverulento : foliis longiuscule petiolatis, subdeltoideo-lanceolatis, rariter serratis : calycibus sub12-floris, tomentulosis, obtusis.

*OBS.* Flores albi.

*HAB.* in scirpetis Carolinæ maritimis.

URTICÆFOLIUM. E. caule lævi, glabro : foliis longe  
L. petiolatis, subcordato-ovalibus, acuminatis, dentatis, glabellis : calycibus simplicibus, sub20-floris.

*OBS.* Flosculi 10-20, candidi : calyce simplici, pariter ac nonnullæ aliæ species, ab EUPATORIIS veris aliquantulum recedere videtur!

*E. ageratoides* GÆRTN. videtur differre calyce imbricato.

*HAB.* in maritimis, a Canada ad Pensylvaniam et in occidentalibus montium *Alléghanis* usque ad Kentucky.

AROMATICUM. E. caule asperiusculo : foliis brevibus,  
L. petiolatis, subcordato-ovalibus, non acuminatis, asperiusculis, rariter et sæpius obtuse dentatis : calycibus sub20-floris, simplicibus.

*OBS.* Flores albi.

*HAB.* a Virginia ad Floridam.

COELESTINUM. E. foliis petiolatis, subcordato-deltoideis, obtuse dentatis, rugosis : corymbulis subumbellatis : calycibus bre-

SYNGENESIA. *POLYG. ÆQUALIS*: 101  
vibus, 30-36-floris : floribus azureo-cœ-  
ruleis.

*OBS.* Datur varietas minor, foliis profundius incis; ;  
quæ forsan *E. incarnatum*. WALTERI.

*HAB.* in sylvis Carolinæ.

C R I T O N I A. GÆRT.

*Cal. comm.* oblongus, imbricatus squamis inæqua-  
libus. *Recept.* nudum. *Pappus* plumosus. *Flos-*  
*culi* omnes androgyni, 5-fidi; tubo antherali  
integerrimo; *stylo* profunde bifido.

KUHNIÆ. C. pubens : foliis lineari-lanceolatis,  
GÆRT. plerisque integris, subtus punctatis : flo-  
ribus paniculatis.

C H R Y S O C O M A. L.

*Recept.* nudum. *Pappus* simplex. *Cal.* hemi-  
sphæricus, imbricatus. *Stylus* vix flosculis lon-  
gior.

NUDATA. C. foliis radicalibus spathulato-lanceo-  
latis; caulinis linearibus, rariter sparsis :  
caule subnudo : corymbo composito,  
elegantè fastigiato : calycibus oblongis,  
3-4-floris.

*OBS.* Flores lutei.

*HAB.* in humidis Carolinæ.

CAPILLACEA. C? foliis capillaceis, multipartitis,

102 SYNGENESIA. *POLYG. ÆQUALIS.*

glabellis : floribus minimis , copiosissimis , composite paniculatis : calycibus 5-floris.

*E. fœniculoides ? WALT.*

*OBS.* Hanc et sequentem ad *CHRYSOCOMAM* refero ; licet genus forsan diversum constituere debeant. Utriusque perennis folia alterna , paucipartita et simplicia , in axillis ramuliferis quasi fasciculata : florescentia *ERIGERI canadensis*. Calyx communis angustatus , imbricatus ; squamulis paucis , linearibus ; extimis brevioribus : flosculi lutei 5 : stigmata longissima , sursum gracilescentia : pappus pilosus , receptaculum nudum , favosulum.

*HAB.* in pascuis , juxta *Charlstown* et in Florida.

*CORONOPIFOLIA.* C? foliis lineari - multipartitis , pubescentibus.

*E. compositifolium ? WALT.*

*OBS.* Habitus florescentiaque præcedentis. Flores subduplo majores : calycibus pubescentibus.

*HAB.* in aridis sylvarum Carolinae.

**SYNGENESIA. POLYG. ÆQUALIS. 103**

**HYMENOPAPPUS. L'HÉRIT.**

**Herba alternifolia; corymbo floribus distinctis.**

**CALYX COMMUNIS** involucriformis, duplici serie subro-phyllus : foliolis consimilibus, laxiuscule patulis, rotundato-ovalibus, membranaceis, coloratis.

**FLOSCULI** plurimi, ovariis densati, corollis discreti; omnes conformes, hermaphroditi-fertiles : congerie depresso-globosa.

**RECEPTACULUM** plano-convexiusculum, nudum.

**FLOSCULORUM**

**COROLLA** infundibuliformis, recta : tubus cylindricus, basi dilatatus, longitudine ovarii : limbus campanulatus, quinquefidus, reflexus.

**STAM.** in tubum coalita; tubus exertus, prismaticus; denticulis conniventibus, brevissimis, obtusis; basi simplici.

**PIST.** ovarium obconicum, obscure trigonum, pubens; denticulis plurimis, erectis apice quasi fimbriatum : stylus longitudine staminum : stigmata duo, exerta, revoluta, lineari-oblonga, obtusa.

**SEMEN** acute obconicum, hirsutulum : brevissime coronatum paleolis plurimis, scarioso - membranaceis, subovalibus,

104 SYNGENESIA. *POLYG. ÆQUALIS.*

- obtusis, subtilissime denticulatis; unica serie ambientibus areolam verticis radiato-substriatam, medio protuberantem, centro minute umbilicatam.

SCABIOSÆUS. *H. candicanti-lanuginosus*: foliis pro-  
*L'HÉRIT.* funde pinnatifidis.

*ROTHIA caroliniensis. Journ. d'Hist. nat. n° 1.*  
cum icone.

*Obs.* Flores majusculi, albi. Interdum folia infima  
opposita.

*HAB.* in Carolina.

P E R S O O N I A.


Herbæ: caulis ramive in pedunculum uniflorum,  
nudum desinentes: folia alterna, integra; nervis  
longitudinalibus: habitus quodammodo PHY-  
TEUMÆ.

CALYX COMM. simplex, polyphyllus: foliolis  
sublanceolatis, erectis subduplici serie  
incumbentibus.

RECEPT. COMM. paleis longitudine calycis,  
angustis, erectis, virescentibus.

FLOSCULI plurimi, calyce longiores, erecti;  
corollulis non congestis, sed quasi dis-  
tincte parallelis: omnes conformes et  
hermaphroditi fertiles.





P. J. Redoute Del.

Plac. Sc.

PERSOONIA *latifolia.*

SYNGENESIA. *POLYG. ÆQUALIS.* 105

Singulorum

COR. recta, infundibuliformis : tubo gracili, longissimo : limbo profunde 5-partito, laciniis linearibus.

STAM. vaginula antherica fere longitudine limbi, angusta : filamenta brevissima.

PIST. ovarium semini subconforme : stylus capillaris : stigmata duo, longiuscule angustequeligulata, solito partim exerta, revoluta.

SEM. oblongo-ovatum, sulcato-striatum : pappus pileis quinque membranaceis, contiguis, acuminatis.

LANCEOLATA. P. caule simplici, inferne folioso, superne nudato : foliis longo-lanceolatis : calycis foliolis obtusis : paleis spathulatis.

*ATHANASIA obovata?* WALT.

*HAB.* in montosis Carolinæ.

LATIFOLIA. P. caule simplici : foliis lanceolato-tab. 43. ovalibus, acuminatis, infimis vaginantibus : calycis foliolis acutis : paleis angusto linearibus.

*ATHANASIA trinervia?* WALT.

*OBS.* Pappus subaristatim acuminatus.

*HAB.* in montosis Carolinæ.



106 SYNGENESIA. *POLYG. ÆQUALIS.*

ANGUSTIFOLIA. P. caule ramoso : foliis infimis angusto-lanceolatis ; cæteris et ramorum angustissime linearibus ; calycis foliolis rigidis , superne angustatis , acutissimis : paleis setaceis.

*HAB.* in territorio Tennasséc.

MELANANTHERA.

*Μελανς , Ανθηρα* : *antherae nigricantes.*

Herbæ : folia opposita : flores subterminales ; pedunculis unifloris , sæpe geminis.

CAL. COMM. primum concavus demumque planus , quasi duplici serie oligophyllus ; foliolis ovalibus , planiusculis , appressis , subæqualibus.

FLOS UNIVERSALIS depresso-globosus vel demum hemisphæricus : flosculis numerosis , omnibus conformibus , hermaphroditis ; centralibus nonnullis solito abortientibus.

RECEPTACULUM convexiusculum , paleaceum : paleis firmis , carinatis , singulos flosculos inferne amplexantibus.

Singulorum flosculorum

COR. candida , infundibuliformis : tubus brevis : limbus tubo multo crassior et longior , tubulosus , quinquefidus ; laciniis conniventibus , lanceolatis.

STAM. vaginula antherica , nigricans , inclusa.

**SYNGENESIA. POLYG. ÆQUALIS. 107**

**PIST.** stylus breviter exertus : stigmata alba, revoluta, superne angustata.

**FRUCT.** capitulum hemisphæricum, calyce vix a paleis supra semina rigide eminentibus distincto : horum unumquodque glabrum, turbinatum, sub 4 - gonum, quasi truncatum ; umbilico angusto, margine annulatim prominulo et setulas paucissimas, numero et longitudine variabiles, inermes, deciduas emittente.

**HASTATA.** M. foliis hastato-trilobis : paleis receptaculi lanceolatis, acuminatis.

**BIDENS nivea. LINN.**

**HAB.** in Carolina.

**OBS.** Planta Antillana quam SWARTZIUS in *Observ.* p. 296 descripsit et cui referendum est synonymon VALLANTII, distincta est species subsequente caractere designanda.

*M. deltoidea* foliis omnibus indivisis, subcordato-deltoideis : paleis receptaculi obtusis.

**CALEA aspera. JACQ. Am. piet.**

*POLYGAMIA SUPERFLUA.*

§. I. PILOSO-PAPPOSÆ.

A S T E R. L.

*Recept.* nudum. *Pappus* simplex. *Cor.* radii plures 10. *Cal.* imbricati squamæ inferiores patulæ.

\* *Radio albo : ligulis 5.*

SOLIDAGINEUS. A? caule simplici : foliis linearib lanceolatis , obtusiusculis , integris , sub3-nerviis ; margine aspero : corymbo fastigiato ; floribus sessilibus , aggregatis , albis , 5-radiatis.

CONYZA *linifolia.* LINN.

*Obs.* Habitus fere SOLIDAGINIS *lanceolatae.* Calyces oblongi , pallido-exalbidi , imbricati ; squamis obtusis , viridibus. Radii quinque , longi.

*HAB.* in Virginia et Carolina.

MARILANDICUS. A? caule simplici : foliis ovali-lanceolatis , superne subserratis ; summis integris : corymbo fastigiato.

CONYZA *asteroides.* LINN.

*Obs.* Rami paucissimi , simplices , summitate subcapituliflori. Flores albi , 5-radiati.

*HAB.* a Pennsylvania ad Virginiam et in montosis Carolinæ.

SYNGENESIA. POLYG. SUPERFL. 109

TORTIFOLIUS. A? foliis parvulis, cuneato-obovatis, integris, tortuose patulis : corymbo composito, subfastigiato.

*CONYZA bifoliata?* WALT.

*OBS.* Flores albi, 5-radiati.

*HAB.* in Carolina inferiore.

**\*\* Radio albo : ligulis pluribus.**

INFIRMUS. A. caule gracili, subflexuoso : foliis subrhomboideis, ovali-lanceolatis, utrinque acuminatis, integris : panicula divergenti-dichotoma ; nudiuscula, pauciflora, floribus omnibus pedunculatis, albis, 8-radiatis.

*A. caule infirmo, foliis ovatis, &c.* GRONOVII.

PLUKN. t. 79. f. 1.

*HAB.* a Canada ad Carolinam, per tractus montium.

ACUMINATUS. A. caule simplici, flexuoso, anguloso : foliis omnibus conformibus, majusculis, ovali-lanceolatis, utrinque subæqualiter angustatis, longissime acuminatis, inæqualiter inciso-serratis : panicula divaricato-dichotoma.

*OBS.* Flores omnes pedunculati, albi.

*HAB.* in Canada.

AMYGDALINUS. A. caule præalto, summitate tantum

*LAM.* ramoso : foliis consimilibus, lanceolatis,

110 SYNGENESIA. *POLYG. SUPERFL.*

integerrimis : corymbo universali partialibusque fastigiatis; floribus albis.

*A. umbellatus. Hort. Kew.*

*HAB.* in Canada et excelsis montibus Carolinæ septentrionalis.

\*\*\* *Foliis integris.*

UNIFLORUS. *A.* caulibus simplicissimis, in pedunculum nudiusculum, 1-florum desinentibus : foliis lanceolatis, integerrimis : calycibus simpliciusculis, subulatis.

*HAB.* in vastis sphagnosis, juxta lacus Mistassinos.

RIGIDUS. *A.* humilis : caulibus erectis, simplicissimis, summitate subsessilibus, paucifloris : foliis linearibus, integerrimis, rigidulis, margine ciliatis.

*HAB.* in aridis Carolinæ.

LINARIIFOLIUS. *A.* decumbens : quasi fruticulosus, minutissime subalbido-pubescens : foliis linearibus, integerrimis, asperis : ramis aggregatis, unifloris : calycibus appressis.

*Obs.* Folia parva, creberrima, carinata, rigida. Rami elongati.

*HAB.* in aridis, a Pennsylvania ad Carolinam; copiose in sabulosis apricis Carolinæ inferioris. *W.*

SYNGENESIA. *POLYG. SUPERFL.* III

GRANDIFLORUS. A. hispidus : foliis oblongis, in-

L. tегris, reflexis : ramulis foliosis, unifloris : calyce polyphyllo, squarroso.

HAB. in montibus Virginiae et Carolinae septentrionalis.

SUBULATUS. A. glaberrimus, parviflorus ; caule paniculato ; ramis multifloris : foliis lineari-subulatis, integerrimis ; calycibus cylindraceis : ligulis radii minutis.

HAB. in maritimis paludosis Pennsylvaniae et Carolinae.

CONCOLOR. A. caulibus virgatis : foliis ovali-lanceolatis,

L. ceolatis, integerrimis : racemo subspicato : calycum squamis linearibus, appresso-imbricatis.

HAB. in sylvis maritimis, a Virginia ad Floridam.

ARGENTEUS. A. caule gracili, decumbente, laxo ramoso ; ramis ramulisve subunifloris : foliis lanceolatis, integerrimis, sericeo-tomentosis, incanis : calycibus subsquarrosis.

Obs. Calycis squamæ foliaceæ, lanceolatae.

HAB. in rupibus ripariis fluminis *Missouri*.

CAROLINIANUS. A. caule frutescente, subscandente,

WALT. ramosissimo : foliis amplexicanti-sessilibus, ovali-lanceolatis, integris : ra-

112 SYNGENESIA. *POLYG. SUPERFL.*

mulis summitate paucifloris : calycibus squarrosis.

*HAB.* in dumosis palustribus, a Carolina inferiore ad Floridam.

SQUARROSUS. A. caule sparsim ramoso, asperius-

*WALT.* culo ; foliis minutissimis , creberrimis , rigide recurvo-reflexis , subcordatis , integris : panicula incompta ; ramulis subunifloris : calycibus subsquarrosis.

*HAB.* in apricis sylvarum Carolinæ inferioris.

CORIDIFOLIUS. A. caule divaricatum ramosissimo :

foliis integerrimis ; caulinis linearibus ; rameis minutis, creberrimis : floribus parvis , laxè paniculatis : calycis squamis appresso-imbricatis , obtusis.

*HAB.* in aridis sylvarum Carolinæ inferioris.

SPARSIFLORUS. A. glabellus : foliis linearibus , integris , reflexis : caule tenui , ramosissimo ; ramis ramulisque patulis , setaceis , 1-floris : calycis squamis appressis.

*HAB.* in Carolina inferiore.

SURCULOSUS. A. humilis : radice repente : caulibus debilibus : simplicibus ; foliis longo-lanceolatis , glabriusculis ; radicalibus spatulatis : corymbo terminali , incompte

**SYNGENESIA. POLYG. SUPERFL. 113**

rarifloro : calycis squamis lineari-oblongis , obtusiusculis.

*HAB.* in sylvia Carolinæ septentrionalis comitatus  
*Burke.*

**DIVERSIFOLIUS.** A. caule aspero - pubente : foliis subintegris, undulatis; inferioribus alato-petiolatis, cordato-ovalibus; superioribus sessilibus, ovali - lanceolatis : panicula laxa , ramis gracilibus , microphyllis.

*Obs.* Flores parvi , omnes longiuscule pedunculati.

*HAB.* in Carolina.

**VILLOSUS.** A. undique villosissimus : foliis linearibus , integerrimis : panicula subsolidaginea , ramis subrecurvo-patulis , racemifloris; floribus sursum secundis : calycibus subglobosis , arcte imbricatis.

*HAB.* in pratis Illinoensibus.

**ERICOIDES.** A. hispidulo-pubescens : caule firmo,  
*L.* ramosissimo : foliis linearibus, integerrimis , pilo acuminatis , sub5-nerviis : panicula ramis recurvo-foliosis; racematum multifloris : floribus pusillis, subalbidis.

*HAB.* in Virginia et regione Illinoensi.

**NOVÆ ANGLIÆ.** A. caule hirsutissimo : foliis amplexicaulibus, cordato-lanceolatis, integris : corymbo subfastigiato : calycibus



114 SYNGENESIA. *POLYG. SUPERFL.*

squarrosus ; foliolis angustissime linearibus.

*HAB.* a Canada ad Virginiam et in regione Illinoensi.

**AMPLEXICAULIS.** A. foliis quasi perfoliato-amplexicaulibus , subcordato-oblongis , inferne non angustatis , integris ; panícula laxa , rariflora.

*HAB.* in Carolinæ inferioris dumosis.

\*\*\*\* *Foliis serratis.*

**MACROPHYLLUS.** A. universe pubescens : foliis radicalibus imisque amplissime cordatis , petiolatis ; omnibus argute serratis , asperis : corymbo floribus majusculis.

*HAB.* in Canada.

**CORDIFOLIUS.** A. caule flexuoso : foliis inferioribus petiolatis , cordatis ; omnibus argute serratis , acuminatis , lævibus : paniculæ floribus divergentibus , omnibus pedunculatis.

*HAB.* a Canada ad Floridam , per tractus montium et in occidentalibus *Alléghanis*.

**BIFLORUS.** A. humilis : caulibus simplicissimis , gracilibus , summitate pedunculis binis , (rarius unico) unifloris : foliis lato-lanceolatis , acutissimis , rariuscule serratis ;

SYNGENESIA. POLYG. SUPERFL. 115

calycis appresso-imbricati squamis lanceolatis.

*HAB.* in septentrionalibus Canadæ, juxta lacus amnesque in sinum *Hudsonis* defluentes. *W.*

TRADESCANTI. *A.* caule puberulo, ab imo ramoso;

*L.* ramis gracilibus, virgatis, patulis, simplicibus, foliose racemifloris; floribus parvis, subsessilibus: calycibus erectis; radiis brevibus, subalbidis: foliis lineari-lanceolatis.

*A. vimineus.* *LAM. Dict.* n° 32.

*HAB.* a Pennsylvania ad Carolinam, per tractus montium.

PUNICEUS. *A.* caule anguloso, hispidō, paniculatum ramoso;

*L.* ramis laxè paucifloris; floribus distinctis, nudiuscule pedunculatis, majusculis: calyce laxo, subuniformiter foliaceo; foliis subamplexicaulibus, lato-lanceolatis, inferne angustatis, rariuscule serratis.

*A. hispidus.* *LAM. Dict.* n° 33.

*A. amœnus.* ——— n° 29.

*HAB.* a sinu *Hudsonis* ad Pennsylvaniam et in excelsis montibus Carolinæ.

SOLIDAGO. L.

*Recept.* nudum. *Pappus* simplex. *Cor.* radii circiter 5. *Cal.* squamæ imbricatæ, clausæ.

PAUCIFLOSCULOSA. *S.* suffruticosa, glabra : foliis lanceolatis, obtusis, enerviis : panicula composite multiflora, fasciculis erectis : calycibus angusto-oblongis, 5-floris; radio unico.

*HAB.* in sabulosis aridis Carolinæ.

LANCEOLATA. *S.* foliis lanceolato-linearibus, integris, trinerviis : corymbo fastigiato; ramulis capitulifloris : ligulis vix manifestis.

Var. *α. major* : foliis rarioribus, latiuscule linearibus : quæ *CHRYSOOMA graminifolia*. LINN.

— *β. minor* : foliis crebrioribus, anguste linearibus; axillis foliosis : subglutinosa.

*HAB. α.* in Canada.

— *β.* in pascuis circa *Charlstown*.

BICOLOR. *S.* caule pubescente : foliis ovali-lanceolatis, subintegris, subalbicantibus, brevi densaque pube asperis : racemis virgatis, interruptis; floribus albidis.

*HAB.* in montibus Carolinæ et in Canada.

SYNGENESIA. *POLYG. SUPERFL.* 117

**ASPERA.** S. caule pubescente : foliis ovali-lanceolatis , serratis , rugosissimis , hirsutie brevi asperrimis : panicula subsecunda , racemis recurvatis.

*Obs.* Folia plantæ sylvestris plerumque lanceolata , utrinque acuta.

*HAB.* in Canada.

**GLOMERATA.** S. caule humili , simplicissimo : foliis glabris , longe lateque lanceolatis , serratis : racemo simplici , e glomerulis axillaribus ; superioribus capitatis congestis : calycibus turgidis ; multifloris.

*HAB.* in montibus Carolinae.

**VIRGATA.** S. glabra : caule virgatim simplicissimo , superne ob foliorum parvitatem quasi nudo : foliis subcuneato-lanceolatis , obtusis , integris , appresso-erectis : racemo simplici , subspiciformi , secundo.

*HAB.* in humidis sylvarum Carolinae inferioris.

**RETRORSA.** S. foliis subamplexanti - sessilibus , lanceolatis , integris , margine asperis . omnibus ab ipsa basi reflexis : panicula racemis recurvatis.

*Obs.* Affinis *S. odoræ*.

*HAB.* in Carolina inferiore.

118 SYNGENESIA. *POLYG. SUPERFL.*

ODORA. S. caule minutim pubescente : foliis

*Ait.* crebris, pusillis, patulis, lineari-lanceolatis, subintegris, margine nervoque asperrimis : panicula subsecunda ; racemis gracilibus, recurvatis.

*HAB.* in Carolina inferiore.

ALTISSIMA. S. caule hirsuto : foliis crebris, sub-

*Ait.* amplexanti-sessilibus, breviuscule lanceolatis, rugosis : panicula racemis recurvatis, subsecunda.

*HAB.* a Pennsylvania ad Carolinam, in locis a mare remotis.

FLEXICAULIS. S. caule glabro, anguloso, flexuoso :

*L.* foliis ovalibus, utrinque acuminatis, argute crebroque serratis, glabris : glomerulis racemulisve axillaribus.

Variat 1°. caule simplici ; glomerulis vel racemis simplicibus, axillaribus.

2°. caule ramoso ; racemis prioris in ramos foliosos et racemifloros conversis.

3°. foliis etiam inferioribus longo-lanceolatis.

*HAB.* in Canada.

RIGIDA. S. undique minuta pube asperrima :

*L.* foliis superioribus semiamplexicaulibus, subcordato-ovalibus : corymbo ramis sub-

**SYNGENESIA. POLYG. SUPERFL. 119.**

fastigiatis , summitate tantum fasciculato-floriferis.

**HAB.** a Pennsylvania ad Carolinam , per tractus montium et in occidentalibus *Alleghanis*.

**SEMPERVIRENS.** S. foliis lanceolatis , lævibus , integerrimis ; axillis solito foliosis seu ramuliferis : panicula racemis rectis.

**Obs.** Foliorum margo modo asper , modo lævis. Mœ panicula oblonga , conferta ; racemis rectis , spiciformibus.

**HAB.** in Canada.

**SENECIO. L.**

**Recept.** nudum. *Pappus simplex. Cal.* cylindricus , calyculatus : squamis apice sphacelatis.

**CINERACIFOLIUS.** S. annuus : foliis amplexicaulibus , oblongis , acutis , inæqualiter acuteque serrato-laceris : calycibus oblongis , lævibus , basi multisetis : radiis nullis.

**HAB.** in Carolina.

**TOMENTOSUS.** S. totus incano - tomentosus : caule simplici : foliis petiolatis , ovali-lanceolatis , subintegris : corymbo subumbellato : radiis parvulis.

**Obs.** proxime affinis *CINERARIÆ integrifoliae* , Jacq.

**HAB.** in Carolinæ loco dicto *Flat-Roc*.

120 SYNGENESIA. POLYG. SUPERFL.

**LYRATUS.** S. annuus, glabellus, simplex : foliis pinnatifido - lyratis ; lobis distantibus , subrotundatis , repando-angulosis vel obsolete dentatis : corymbo composito ; pedunculis summis subumbellatis ; floribus multiradiatis.

*HAB.* in nemoribus Carolinæ.

**PAUPERCULUS.** S. pumilus : caule simplicissimo , rigide erecto , subnudo : foliis omnibus lanceolatis , nonnullis radicalibus subintegris , cæteris rariter inciso-dentatis serratisve : corymbo paucifloro ; calycibus glabellis ; radiis parvulis.

*Obs.* Flores magnitudine *Jacobeae*.

*HAB.* in Canada, juxta lacus.

**AUREUS.** S. caule simplicissimo , remote oligophyllo : foliis denticulatis ; radicalibus cordatis , rotundato-obtusis ; caulinis petiolo pinnatifidis : corymbo pedunculis unifloris subumbellato.

*HAB.* a Canada ad Virginiam et in excelsis montibus Carolinæ.

SYNGENESIA. POLYG. SUPERFL. 121

TUSSILAGO. L.

*Recept. nudum. Pappus simplex. Cal. squamæ æquales, discum æquantes, submembranaceæ.*

**INTEGRIFOLIA.** T. foliis oblongo-ovalibus, integerrimis . scapo nudissimo, nutante, unifloro : ligulis radiantibus femineis ; flosculis extimis filiformibus, femineis; intimis infundibuliformibus, submasculis.

*PERDICUM semiflosculare. WALT.*

*HAB.* a Carolina ad Floridam, in sylvis humidis.

DORONICUM. L.

*Recept. nudum. Pappus simplex. Calycis squamæ duplicis ordinis æquales, disco longiores. Sem. radii nuda pappoque destituta.*

**NUDICAULE.** D. foliis decussatim oppositis, oblongo-ovalibus, subintegris : caule subaphyllo, summitate in nonnullos pedunculos 1-flo-ros diviso : seminibus radii papposis.

*Obs.* Caulis 1-2-pedalis, duobus tantum foliorum parvulorum paribus remotissimis solito vestitus. Calycis foliola lanceolata. Videtur *DORONICUM foliis plantagineis*, &c. CLAYT. *Fl. Virg.* Phrasis GRO-NOVII non quadrat.

*HAB.* in umbrosis sylvarum, a Virginia ad Floridam.



I N U L A. L.

*Recept.* nudum. *Pappus simplex.* *Antherae* basi in setas duas desinentes.

MARIANA. I. pilosa : caulibus simplicibus : foliis sessilibus, subcuneato-lanceolatis, subserratis : corymbo simplici, subumbelliformi : pedunculis calycibusque minutim confertimque glanduloso-pubescentibus.

*HAB.* a Virginia ad Carolinam.

GOSSYPINA. I. tota lanugine gossypina mollissima candicans : foliis sessilibus, subspathulato-oblongis, rotundato-obtusis, integris : corymbo subfastigiato.

*HAB.* in maritimis Carolinæ et Floridæ.

GRAMINIFOLIA. I. argenteo-sericea : caule simplici : foliis longissime linearibus, integerrimis, longitudinaliter nervosis : corymbo laxo, composito ; floribus omnibus pedunculatis : calycibus turbinatis ; squamis numerosis, acutissimis, superne glandulosis.

*ERIGERON glandulorum?* WALT.

*HAB.* a Carolina ad Floridam, frequens.

SYNGENESIA. POLYG. SUPERFL. 123

ERIGERON. L.

*Recept.* nudum. *Pappus* pilosus. *Cor.* radii lineares, angustissimæ.

CANADENSE. E. annuum : caule hispido, paniculato latius ramosissimo; ramis quasi racemose multifloris : foliis lanceolato-linearibus, ciliatis : calycibus cylindricis, subimbriatis : radiis multiplici serie confertis, brevissimis, albis.

*HAB.* a Canada ad Floridam.

DIVARICATUM. E. annuum, pusillum, divaricataramosissimum, subfastigiato-paniculatum, multiflorum : foliis subulatis : floribus brevissime radiatis; corollulis disci 4-fidis.

*Obs.* Affine *E. canadensi*.

*HAB.* in pratensibus Illinoensibus, prope *Kaskaskia*.

HYSSOPIFOLIUM. E. caule ramis sterilibus paniculato; paucis in pedunculos nudos, unifloros protractis : foliis linearibus, integris, glabellis, ciliatis : radiis calyce cylindraceo multo longioribus.

*HAB.* ad sinum *Hudsonis* et juxta lacum *Mistassins*.

PHILADELPHICUM? E. puberulum : foliis cuneato-oblongis, rariter inciso-dentatis : caule debili, simplici; superne ramis raris,

224 SYNGENESIA. *POLYG. SUPERFL.*

paucifloris : radiis angustissimis, cœrulescentibus, calyce hemisphærico duplo longioribus.

*HAB.* in Carolinae campestribus.

**PULCHELLUM.** *E. hirsutum, subincanum* : foliis radicalibus oblongo-obovalibus sive cuneato-oblongis, subintegris; caulinis paucis, distantibus, semiamplexicaulibus : floribus majusculis; radiis numerosis, elongatis, subcœrulescenti-albidis.

*E. caule simplicissimo, saepius bifloro, foliis caulino semiamplexicauli.* GRONOV. *Fl. Virg.* 122.

*Obs.* Caulis simplicissimus, summitate divisus in pedunculos paucos, unifloros.

*HAB.* in Canada, Pensylvania et montibus Carolinae.

**NUDICAULE.** *E. glabellum* : foliis ovali-lanceolatis, acutis, subdentatis : caule simplicissimo, subaphyllo, elongato, summitate corymbo paucifloro; floribus pedunculatis : radiis longitudine calycis, subalbidis.

*HAB.* in udis herbosis Carolinae.

BACCHARIS. L.

*Recept.* nudum. *Pappus* pilosus. *Cal.* imbricatus, cylindricus. *Flosculi* feminei hermaphroditis immixti.

*Obs.* Species omnes veræ hujus generis dioicæ fruticant.

ANGUSTIFOLIA. B. glaberrima, paniculato-ramosissima : foliis angusto-linearibus, integris : panicula composita, multiflora : calycibus minusculis, sub20-floris.

*Obs.* Folia primaria interdum rarissime dentata.

*HAB.* in scirpetis maritimis, a Carolina ad Floridam.

SESSILIFLORA. B. foliis lævibus, cuneato-obovalibus, superne dentatis : floribus in ramulis per glomerulos axillares, sessiles, remotos dispositis : calycum squamis superne rufis.

*Obs.* B. *halimifoliae* affinis, sed diversa.

*HAB.* in sylvis maritimis Carolinæ.

HALIMIFOLIA. B. foliis rhomboideo-obovalibus, L. superne inciso-dentatis : panicula composita, foliosa ; fasciculis pedunculatis.

*Obs.* Quasi subalbido-pulverulenta.

*HAB.* in maritimis, a Virginia ad Floridam.

126 SYNGENESIA. POLYG. SUPERFL.

C O N Y Z A. L.

*Recept.* nudum. *Pappus simplex*. *Cal.* imbricatus, subrotundus. *Cor.* radii trifidæ.

MARILANDICA. *C. puberula*, erecta, herbacea : foliis sessilibus, lato-lanceolatis, acutissimis, acute serratis : panicula e corymbulis ramulos fastigiatim terminantibus, purpureis.

DILL. *Elth.* t. 88. f. 104. nondum perfecte florida.

BACCHARIS *fætida*? WALT.

ERIGERON *camphoratum*? LINN.

HAB. a Pensylvania ad Carolinam.

AMPLEXICAULIS. *C. puberula*, subglutinosa, erecta, herbacea : foliis amplexicaulibus, cordato-oblongis, serratis : paniculæ corymbulis capitato-glomeratis.

HAB. in humidis Carolinæ.

PYCNOSTACHYA. *C. caule alato* : foliis lanceolatis, subtus tomentosis, subintegris ; superioribus sublinearibus : spica confertim et continue capitata.

GNAPHALIUM *undulatum*. WALT.

Obs. Ab antillana *C. virgata* LINN. distincta species ; quæ forsân genus CONYZÆ et GNAPHALIO intermedium constituet.

HAB. a Carolina ad Floridam.

**G N A P H A L I U M. L.**

*Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus : squamis marginalibus rotundatis, scariosis, coloratis.

**MARGARITACEUM.** G. herbaceum : foliis lineari-lanceolatis, sensim angustatis, acutis, corymbo composito, fastigiato ; floribus pedicellatis, albis.

*HAB.* a Canada ad Virginiam.

**POLYCEPHALUM.** G. herbaceum, erectum : foliis lineari-lanceolatis : caule summitate composita paniculato ; glomerulis numerosis, pallidis.

*Obs.* *G. obtusifolium*, LINN. omisso synonymo *DILENII* ; qui plantam cultura mutatam tradidit.

*HAB.* in sylvis Carolinæ et Virginiae.

**PURPUREUM.** G. herbaceum : caulibus simplicibus :  
*L.* foliis spathulato-lanceolatis : spica interrupta ; glomerulis omnibus sessilibus.

*Obs.* Habitus *G. sylvatici*.

*HAB.* a Pennsylvania ad Carolinam.

**ULIGINOSUM.** G. annuum, humile, diffuse ramulosum : foliis linearibus : capitulis foliosis, sessilibus.

*HAB.* a Canada ad Pensylvaniam.

128 SYNGENESIA. POLYG. SUPERFL.

DIOICUM : *β. plantaginifolium*. G. perenne : stolonibus procumbentibus : foliis spathulato-obovalibus , mucronatis : caule simplicissimo : corymbo terminali simplici ; floribus dioicis.

*G. plantaginifolium*. LINN.

OBS. Ab Europæo nonnisi latioribus foliis differt. Interdum legi prorsus priori simile.

Variat caule breviori , unifloro ; flore manifeste majore.

HAB. a Canada ad Carolinam et in occidentalibus *Alléghanis* montibus.

§. II. NON PAPPOSE.

ARTEMESIA. L.

*Recept.* subvillosum vel nudiusculum. *Pappus* nullus. *Cal.* imbricatus squamis rotundatis , conniventibus. *Cor.* radii nullæ.

VULGARIS. A. erecta : foliis pinnatifidis ; subtus  
L. tomentosis ; laciniis latis , incisis ; lacinulis lanceolatis : floribus sessilibus , ovoideocylindraceis.

1 HAB. in septentrionalibus Canadæ.

CANADENSIS. A. subdecumbens , parce pubescens : foliis planis , lineari-pinnatifidis : ramu-

**SYNGENESIA. POLYG. SUPERFL. 129**

lis spicifloris ; floribus majusculis , sub-hemisphæricis : calycibus scariosis.

*Obs.* Calycum squamæ subrotundo-ovales : receptaculum glabrum.

*HAB.* in arena mobili secus lacus , ad sinum *Hudsonis*.

**LAUDATA.** A. erecta , glabra : foliis subsetaceo-pinnatifidis ; laciniis convexis : caule primariisque ramis in racemum longissimum , strictum dispositis ; ramulis confertis ; floribus omnibus pedicellatis , subgloboso-ovoideis.

*HAB.* ad ripas sabulosas fluminis *Missouri*.

**E C L I P T A. L.**

*Recept.* paleaceum. *Pappus* nullus. *Corollulae* disci 4-fidæ.

*Obs.* Species hujus generis , per Asiam , Africam et Americam dispersæ , adhucdum obscuræ. Speciem olim legi ad *Euphratem* , quæ forsân *MICRELIIUM astéroides* FORSKALII.

**PROCUMBENS.** E. procumbens assurgensve : foliis longo-lanceolatis , inferne angustatis , rariter subserratis : pedunculis longiusculis : calycum foliolis acute lanceolatis : flosculis quadrifidis.

*HAB.* in Carolina.



130 SYNGENESIA. POLYC. SUPERFL.

BRACHYPODA. E. divaricato-prostrata : foliis lanceolatis, planis, rarissime leviterque subserratis : pedunculis solitariis geminisque, brevibus : calycum foliolis ovalilanceolatis : flosculis quinquefidis.

AMELLUS? *carolinianus*. WALT.

HAB. in Carolina.

BUPHTHALMUM. L.

*Recept.* paleaceum. *Pappus* margo obsoletus.  
*Seminum* latera, præsertim radii, marginata.  
*Stigma* floscul. hermaphroditicorum indivisum.

HELIANTHOIDES. B. caule herbaceo, lævi, rore  
L. glauco consperso : foliis oppositis, petiolatis, subcordato-ovalibus, argute serratis : calycum foliolis oblongis, obtusiusculis : disco convexo : seminibus nudissimis.

*Obs.* Forsan huc referenda *RUDBECKIA oppositifolia*, LINN.

HAB. in montibus Carolinæ et in regione Illinoensi.

FRUTESCENS. B. fruticosum : foliis oppositis, cuneato-lanceolatis, incanis : petiolis bidentatis.

HAB. in maritimis, a Carolina ad Floridam.

SYNGENESIA. POLYG. SUPERFL. 131

BELLIS. L.

*Recept.* nudum, conicum. *Pappus* nullus. *Cal.* hemisphæricus : squamis æqualibus. *Sem.* obovata.

INTEGRIFOLIA. B. caulescens, divaricato-ramosa : foliis integerrimis ; inferioribus obovatis ; supremis lanceolatis : foliolis calycum acutissimis, pilo acuminatis.

*HAB.* ad ripas rivulorum et in collibus umbrosis Tenassée.

SPILOANTHUS. L.

*Recept.* paleaceum. *Sem.* apice 2-setosa, setis minimis, deciduis. *Cal.* duplici serie polyphylus, subæqualis, hemisphæricus, squamis non divaricatis.

REPENS. S. caule repente : foliis ovali-lanceolatis, subintegris : pedunculis solitarie terminalibus, longissimis : floribus 10-radiatis : seminibus apice nudis.

*ANTHEMIS repens.* WALT.

*Obs.* Hæc et plures aliæ species Indiæ utriusque forsân in posterum constituent genus distinctum a SPILOANTHO, præsertim seminibus nudis.

*HAB.* in Carolinæ inundatis, ad fluvium *Santee*.

132 SYNGENESIA. *POLYG. SUPERFL.*

T A G E T E S. *L.*

*Recept.* nudum. *Pappus* aristis 5, erectis. *Cal.*  
1 - phyllus, 5 - dentatus, tubulosus. *Flosculi*  
radii quatuor, persistentes.

**PAPPOSA.** T. foliis anguste duplicato - pinnatifidis : pedunculis brevissimis, foliosis : calycibus involucello calyculatis : radiis subnullis : seminibus pappo fasciculatim piloso.

*HAB.* in locis alluvionibus fluvii *Missouri* quotannis inundatis etiamque in ripariis sabulosis *Mississippi*.

B O L T O N I A. *SCHREB.*

*Cal. comm.* convexus, imbricatus. *Recept.* nudum. *Flores* radiati. *Semina* plano-compressa, membranaceo-marginata. *Pappus* setulis minutis, pluribus; duabus oppositis interdum elongatis.

**GLASTIFOLIA.** B. floribus breviter pedunculatis : seminibus obcordatis, conspicue alatis, puberulis; aristis pappi duabus ipsorum longitudine.

*HAB.* in regione Illinoensi.

**ASTEROIDES.** B. floribus longe pedunculatis : seminibus ovalibus, glabris, submuticis.

*Obs.* Hædæ species habitu, foliatione coloreque florum omnino conveniunt.

SYNGENESIA. *POLYG. SUPERFL.* 1331

HELENIUM. *L.*

*Recept.* nudum : *Radii* paleaceum. *Pappus* 5-*aris-*  
tatus. *Cal.* simplex , multipartitus. *Corollulae*  
radii semitrifidæ.

AUTUMNALE. *H.* foliis lato-lanceolatis , serratis :  
*L.* corollulis disci quinquefidis.

*HAB.* a Pennsylvania ad Carolinam et in occidenta-  
libus *Alléghanis*.

QUADRIDENTATUM. *H.* foliis angusto-lanceolatis ,  
întegris , latius decurrentibus : flosculis  
disci quadrifidis.

*Act. soc. hist. nat. paris.* 22. t. 4.

*HAB.* in Carolina.

*POLYGAMIA FRUSTRANEA.*

VERBESINA. L. GRON.

*Recept.* paleaceum. *Pappus* aristatus. *Cal.* duplici ordine. *Flosculi* radii circiter 5.

VIRGINICA. V. caule anguste alato : foliis alternis,  
L. lato-lanceolatis, rariter subdentatis : corymbis compositis, subfastigiatis : calycibus oblongis, dense puberulis.

*Obs.* Radii 3-4, albi.

*HAB.* in Pennsylvania, Virginia et Carolina.

SIGESBECKIA. V. caule alato : foliis oppositis, ovali-lanceolatis, utrinque acuminatis, acute serratis : panicula parva, brachiata ; ramulis summitate fasciculifloris.

SIGESBECKIA *occidentalis.* LINN.

Non autem PHAETHUSA. GERT.

*Obs.* Calyces floridi oblongi, quasi paleacei : radii 1-3, lutei : semina non marginata.

*HAB.* in Virginia et Carolina.

COREOPSIS. V. caule præalto, alato : foliis alternis, lanceolatis, serratis : panicula com-

SYNGENESIA. *POLYG. FRUSTR.* 135.

posita, laxa, foliosa : calyce laxe patulo ; disco subgloboso.

*COREOPSIS alternifolia*. LINN.

Var. *α. lutea* : radiata aut flosculosa, luteiflora.

— *β. alba* : semper flosculosa, albiflora.

*HAB. α.* in Virginia et Carolina superiore.

— *β.* in Carolina maritima.

**HELIANTHOIDES.** V. caule alato : foliis alternis, latiuscule lanceolatis, acutis, leviter dentatis, subtus subcandicanti-villosis, supra pube asperrimis : pedunculis 1-floris, parce in summitate aggregatis.

*Obs.* Flos. HELIANTHI, luteus : semina *COREOPSIS alternifoliae*.

*HAB.* in occidentalibus *Alléghanis*, territorio Tennesée, et regione Illinoensi.

**B I D E N S. L.**

*Flores* flosculosi aut etiam radiati; radiis neutris aut imperfecte masculis. *Receptaculum* paleaceum; paleis extimis quasi calycem communem constituentibus. *Semina* aristata; aristis duabus vel pluribus, retrorsum aculeatis.

**BIPINNATA.** B. glaber : foliis bipinnatis ; pinnulis

*L.* subrhomboideis : pedunculis nudis : involucello brevi, lineari : radiis quinque

136 SYNGENESIA. *POLYG. FRUSTR.*

aut paucioribus, luteis : seminibus angusto-elongatis, 3-4-aristatis.

*HAB.* a Pennsylvania ad Carolinam.

**FRONDOSA.** B. caule glabriusculo : foliis ternatim

*L.* quinatimve pinnatis; foliolis lanceolatis, serratis : involucello foliaceo, ciliato, calycibus frondosis longiore : floribus flosculosis : seminibus latis, biaristatis.

*HAB.* in regione Illinoensi; montibus *Alleghanis*, &c.

**CHRYSANTHEMOIDES.** B. glabra : foliis subconnatis, lanceolatis, rariter serratis : floribus erectis, aureis, multiradiatis : seminibus biaristatis.

*COREOPSIS perfoliata.* WALT.

*OBS.* Affinis *COREOPSISIDI Bidenti*, LINN. quæ forsane eadem, si semina tantum biaristata.

*HAB.* a Pennsylvania ad Carolinam, in aquosis.

COREOPSIS. L.

*Cal. comm. et Recept. Bidentis. Flores radiati ; radiis neutris, prolixis, luteis. Sem. compressa, aut emarginata, aut bidentata aut etiam bisetosa ; setis inermibus nec retrorsum aculeatis.*

\* *Foliis omnibus indivisis.*

LANCEOLATA. C. brevicaulis : foliis cuneato-lanceolatis linearibusve, integerrimis : pedunculis longissimis, simplicibus, nudis : seminibus orbiculatis, convexo-concavis, scabris, manifestius alatis, apice bidentatis, emarginatis.

Var. *α. glabella* : glabra, foliis margine subciliatis, angustioribus.

— *β. villosa* : caule foliisque canescenti-villosis.

HAB. in Carolina.

DICHOTOMA. C. caule glabro, superne rariter dichotomo et nudiusculo : foliis indivisis, integerrimis, in petiolum angustatis, plerisque alternis : seminibus obovalibus, breviter bisetosis, scaberulis ; ala marginali fimbriato-lacera.

*C. gladiata* ? WALT.

HAB. in sphagnosis umbrosis Carolinæ.

LATIFOLIA. C. foliis breviter petiolatis, lato-ova-



138 SYNGENESIA. *POLYG. FRUSTR.*

libus , acuminatis , crenato - dentatis ;  
crenis minute mucronatis : radiis integris : seminibus cuneato-oblongis , apertis , apice nudis.

*Obs.* Habitus *SILPHII*.

*HAB.* in excelsis montibus Carolinæ.

*\*\* Foliis partitis.*

*AURICULATA.* C. caule pubente : foliis subsessilibus , ovali-lanceolatis , integerrimis ; inferioribus auriculato-tripartitis : calyce exteriori fere ad pedunculum partito : radiis 4-dentatis : seminibus subrotundo-obovalibus , apice emarginato - bidentatis.

*HAB.* in montibus Carolinæ et Virginie.

*TRIPTERIS.* C. glabra : foliis petiolatis , subæqualiter trifoliatis vel quinato - pinnatis ; foliolis lato - lanceolatis , integerrimis : radiis integris : seminibus obovalibus , apice nudis.

*Obs.* Calycis exterioris lacinie in basin solidam distanter a pedunculo connate.

*HAB.* in altis Virginie Carolinæque montibus.

*SENIFOLIA.* C. minutissime puberula : foliis arcte sessilibus , æqualiter trifoliatis ; foliolis

SYNGENESIA. POYG. FRUSTR. 139

quasi seno-verticillatis, integerrimis : radiis integris : seminibus cuneato-oblongis.

*C. major* ? WALT.

*Obs.* Interdum fere glabra.

*HAB.* in collibus sabulosis Carolinæ.

VERTICILLATA. *C.* glabra : foliis sessilibus, ad basim tripartitis, partitionibus lineari-multipartitis vel pinnatifidis : seminibus ovalibus, muticis.

Var. *α. tenuifolia* : laciniis foliorum angustissimis et numerosioribus.

— *β. linearis* : laciniis foliorum paucioribus et latiuscule linearibus.

*Obs.* Affinitas cum *C. tripteride*.

*HAB. α.* in Virginia.

— *β.* in Carolina.

TRICHOSPERMA. *C.* minuscula, glabella, subdichotoma : foliis subquinato-pinnatis ; pinnulis lanceolatis, rariter inciso-serratis ; calyce exteriore sub8-phyllo ; foliolis spatulatis, ciliato-serrulatis : radiis integris : seminibus cuneatis.

*Obs.* Affinis *C. coronatae*. LINN. omissis synonymis VALLANTII et PLUMIERI. Folia ima septenato-pinnata : semina apice sub4-dentata ; denticulis 2 vix manifestis.

*HAB.* in humidis Carolinæ superioris.

140 SYNGENESIA. POLYG. FRUSTR.

MITIS. C. glabra : foliis petiolatis ; infimis duplicato-pinnatifidis ; supremis lineari-tripartitis : calycibus simpliciusculis , rarius subfrondosis : seminibus oblongiuscule cuneatis , apteris.

HAB. in paludosis Carolinæ.

ARISTOSA. C. tota puberula : foliis quinato-pinnatis ; foliolis serratis : radiis integris , lato-ovalibus : seminibus cuneato-obovalibus , biaristatis ; aristis longissimis , divaricatis.

HAB. in regione Illinoensi.

HELIANTHUS. L.

Recept. paleaceum , planum. *Pappus* diphyllus.  
Cal. imbricatus , subsquarrosus.

ATRORUBENS. H. totus hispidus : caule superne nudiusculo , laxè paniculato : foliis ovalibus , triplinerviis calycum squamis erectis , ovali-lanceolatis ; disco atropurpureo.

HAB. a Pensylvania ad Carolinam.

CANESCENS. H. pubescenti-canescens : caule villoso : foliis semiamplexicaulibus , subcordato-ovalibus , integrinisculis : calycibus villosissimis , foliolis lanceolatis.

HAB. in pratensibus irriguis regionis Illinoensis et Tennassée.

SYNGENESIA. *POLYG. FRUSTR.* 141

**TOMENTOSUS.** H. caule dense pubente : foliis plerisque alternis, ovali-lanceolatis, subintegris, subtus tomentosis, supra asperimis : floribus breviter pedunculatis.

*HAB.* in pratensibus Illinoensibus.

**DIVARICATUS.** H. foliis suboppositis, lanceolatis :  
**L.** caule glabello, ramosissimo : panicula quasi trichotome multiflora; ramis gracilibus, multifloris : floribus minimis.

*HAB.* in montibus Virginiae et Carolinae.

**GIGAS.** H. caule hispido, dichotomo : foliis alternis, lanceolatis, rariter serratis : calycibus hispidulis, ciliatis, disco longioribus; foliolis longo-linearibus.

*HAB.* in montibus Virginiae et Carolinae.

**ANGUSTIFOLIUS.** H. caulibus gracilibus, subunifloris :  
**L.** foliis oppositis alternisve, linearibus, integris, margine revolutis, asperis : disco convexo, subfusco.

*HAB.* a Virginia ad Floridam.

144 SYNGENESIA. *POLYG. NECESS.*

SPATHULATA. *R. parvula*, minute pubescens : caulibus gracilibus, 1-floris : foliis parvulis, alternis, spathulato-obovalibus, integris : calyce patulo : radiis tridentatis.

*HAB.* in montibus Carolinæ.

TRILOBA. *R. hirsuta* : foliis inferioribus tripartitis ; superioribus indivisis, lato-lanceolatis.

Var. *α*. foliis hirtellis, manifestius dentatis ; calyce reflexo.

— *β*. foliis subtomentosis ; calyce incumbente.

*HAB. α.* in montibus excelsis Carolinæ.

— *β*. in regione Illinoensi.

LACINIATA. *R. caule lævi*, glabro : foliis subpinnato-laciniatis, segmentis ovali-lanceolatis : disco sphæroideo ; corollulis laxiusculis.

*OBS.* Huc synonymon BAUHINI et icon MORIS. s. 6. t. 6. fig. 53.

*HAB.* in umbrosis sylvarum Virginie et regionis Illinoensis.

PINNATA. *R. caule anguloso*, pubente : foliis laciniato-pinnatis ; segmentis lanceolatis :

SYNGENESIA. POLYG. NECESS. 145

disco oblongiuscule ovoideo ; corollulis densatis.

*Obs.* Ex LINNÆO varietas *R. laciniatæ* ; sed diversa species.

MORIS. 5. 6. t. 6. f. 54.

*HAB.* in apricis Canadae et in collibus regionis Illinoensis.

SILPHIUM. L.

*Recept.* paleaceum. *Pappus* marginato-bicornis.

*Cal.* squarrosus.

LACINIATUM. S. foliis radicalibus caulinisque pinnatifidis ; laciniis lanceolatis : caule superne hispido : calycibus magnis, hirtis ; foliolis amplis, subcordatis, acuminatis.

*HAB.* a regione Illinoensi ad Ludoviciam.

COMPOSITUM. S. foliis radicalibus trifoliatis ; foliolis petiolatis, inæqualiter sinuato-multipartitis ; caulinis sinuato-pinnatifidis : caule lævi : floribus parvis, paniculatis.

*S. laciniatum.* WALT.

*HAB.* in sylvis maritimis, a Carolina ad Floridam.

TEREBINTHINACEUM. S. foliis radicalibus amplis, cordatis ; caulinis alternis, ovalibus ; supremis canaliculatis : caule lævi : panicula laxa.

*HAB.* a regione Illinoensi ad Ludoviciam.

146 SYNGENESIA. POLYG. NECESS.

TERNIFOLIUM. S. foliis terno-verticillatis, lanceolatis : caule lævi : panícula multiflora.

*S. trifoliatum*. LINN.

HAB. in montibus Virginæ et Carolinæ.

CONNATUM. S. foliis sessiliter aut quasi petiolatim

L. connatis, subovalibus : caule glabro.

OBS. *S. perfoliatum*. LINN. hujusce varietas.

HAB. in occidentalibus *Alléghanis*, Kentucky et regionis Illinoensis.

ASTERISCUS. S. caule pedunculisque hispidis : foliis

L. oppositis alternisve, sessilibus, ovali-lanceolatis, serratis vel crenatis, utrinque hirsutis : calycibus ciliatis.

HAB. a Virginia ad Carolinam.

INTEGRIFOLIUM. S. caule quadrangulato, aspero : foliis omnibus uniformibus, oppositis, sessilibus, erectis, oblongo-ovalibus, supra scaberrimis : floribus paucioribus, breviter pedunculatis.

HAB. in regione Illinoensi.

PUMILUM. S. caule pumilo, tenuiter tomentoso : foliis ovalibus, obtusis, leviter dentatis, subtomentosis.

HAB. in Florida.

**SYNGENESIA. POLYG. NECESS. 147**

**POLYMNIA. L.**

*Recept. paleaceum. Pappus nullus. Cal. exterior*  
*4-5-phyllus, interior 10-phyllus, foliolis concavis.*

**CANADENSIS.** P. viscido-villosa : foliis alternis (ra-  
**L.** rius passim oppositis), hastato-sinuatis  
angulosive : radiis minutissimis, tri-  
fidis.

*HAB. in sylvis opacis, ad flumen Ohio.*

**UYEDALIA.** P. foliis oppositis, sinuato-lobatis,  
**L.** asperiusculis : radiis elongatis.

*HAB. a Virginia ad Carolinam.*

**TETRAGONOTHECA.** P. hirsutissima : foliis opposi-  
**L.** tis, lanceolato-ovalibus, basi cuneatis,  
serratis : radiis longissimis, trifidis; exte-  
rius disci flosculis fertilibus.

*HAB. a Virginia ad Floridam.*

**PARTHENIUM. L.**

**MASC.** *Cal. comm. 5-phyllus. Cor. disci monope-*  
*talæ.*

**FOEM.** *Cor. radii 5, singulæ 2, masculæ: inter-*  
*media foeminea, supera. Sem. nudum.*

**INTEGRIFOLIUM.** P. foliis ovalibus, inæqualiter  
**L.** dentatis, asperis; superioribus amplexi-  
caulibus.

*HAB. in Virginia et in montosis Carolinæ.*



148 SYNGENESIA. *POLYG. SEGREG.*

CHRYSO G O N U M. L.

*Recept.* paleaceum. *Pappus* 1-phyllus, 5-dentatus.  
*Cal.* 5-phyllus. *Semina* calyculo 4-phylo involuta.

VIRGINIANUM. C. pumilum ; totum lanato-villo-  
L. sissimum : foliis in petiolum angustatis,  
ovalibus , dentatis.

*HAB.* a Virginia ad Carolinam , in montosis.

*POLYGAMIA SEGREGATA.*

E L E P H A N T O P U S. L.

*Calyculus* 4-florus. *Corollulae* ligulae , herma-  
phroditæ. *Recept.* nudum. *Pappus* setaceus.

SCABER. E. folijs oblongis , asperiusculis : ca-  
L. pitulis terminalibus paniculatis : pappo  
aristis rectis.

*HAB.* a Virginia ad Floridam.

*MONOGAMIA.*

*IMPATIENS. L.*

*Cal.* 2-phyllus. *Cor.* 5-petala, irregularis, nectario cucullato. *Caps.* supera, 5-valvis.

**NOLI TANGERE.** I. pedunculis multifloris, solitariis;  
*L.* foliis ovatis, geniculis caulinis tumen-  
 tibus.

Var. *α.* floribus luteis; foliis majoribus, ob-  
 tuse dentatis.

— *β.* floribus fulvis; foliis minoribus, ar-  
 gute dentatis.

*HAB. α.* in Virginia, Carolina.

— *β.* a Canada ad Carolinam.

*VIOLA. L.*

*Cal.* 5-phyllus. *Cor.* 5-petala, irregularis, pos-  
 tice cornuta. *Caps.* supera, 3-valvis, 1-locu-  
 laris.

**PENNSYLVANICA.** V. tota villosa-pubescent : cau-  
 libus summitate tantum foliosis : foliis  
 lato-cordatis : stipulis membranaceis,  
 ovalibus, subserratis : floribus luteis.

*HAB.* in umbrosis, juxta rivulos Pennsylvaniae, præ-  
 sertim ad *Skullkill*.

**HASTATA.** V. glabella : caulibus simplicissimis,

150 SYNGENESIA. *MONOGANIA*.

summitate tantum foliosis : foliis alternis , hastatis , brevissime petiolatis : stipulis minutis , denticulatis : floribus luteis.

*HAB.* in altis montibus Carolinæ.

DEBILIS. V. caulescens , debilis : stipulis membranaceis , lanceolatis , tenuiter laceris : foliis breviuscule cordatis , denticulatis : floribus albis ; longe pedunculatis.

*V. canina* ? WALT.

*HAB.* in montibus *Alléghanis*.

CANADENSIS. V. caulescens , erecta , puberula :

*L.* stipulis integris : foliis lato-cordatis , acuminatis , dentatis.

*HAB.* in Canada et altis montibus Carolinæ.

ROTUNDIFOLIA. V. acaulis : foliis orbiculato-cordatis , sinu clauso , leviter dentatis , glabriusculis ; petiolo pubescente : flore luteo ; calyce obtuso.

*HAB.* in excelsis montibus Carolinæ.

LANCEOLATA. V. acaulis , glabella : foliis lanceo-

*L.* latis , obsolete denticulatis crenulatisve : floribus albidis.

*HAB.* a Nova Anglia ad Carolinam.

SYNGENESIA. MONOGAMIA. 151

PEDATA. V. acaulis : foliis subpedato-multipar-

L. titis ; laciniis lineari-lanceolatis , eden-  
tulis : calycis laciniis linearibus , acutis.

HAB. a Nova Anglia ad Carolinam , per tractus mon-  
tium.

PALMATA. V. acaulis , pubescens : foliis hastato-

L. lobatis palmatisve ; lobis subdentatis.

OBS. Folia nonnulla interdum etiam omnia indivisa ,  
reniformi-cordata.

V. villosa et cordata. WALT. hujus varietates?

HAB. in Carolina.

LOBELIA. L.

Cal. 5-fidus. Cor. 1-petala , irregularis. Caps.  
infera , 2-3-locularis.

CARDINALIS. L. pubescens , erecta , simplex : fo-  
liis lato-lanceolatis , crasso-denticulatis :  
spica subsecunda ; floribus pedicellatis ,  
coccineis : genitalibus corolla longio-  
ribus.

HAB. a Canada ad Carolinam , in opacis sylvarum.

SIPHILITICA. L. hirtella , simplex , erecta , majus-

L. cula : foliis ovali-lanceolatis , obsolete  
subserratis : spica foliosa , longa : caly-  
cibus hirsutis , sinibus reflexis.

HAB. a Pennsylvania ad Carolinam , per montium trac-  
tus et in Kentucky.

152 SYNGENESIA. MONOGAMIA.

INFLATA. L. erecta, ramosa, hirsutissima : foliis ovali-lanceolatis, serratis : spicis foliosis; floribus pusillis : capsulis inflatis.

HAB. a Canada ad Carolinam montosam.

PUBERULA. L. erecta, simplicissima, pubescens : foliis oblongo-ovalibus, obtusis, repando-serrulatis : spica non pedunculata; floribus paucis, alternis, subsessilibus : calycibus ciliatis.

HAB. in Carolina.

CLIFFORTIANA. L. glabella, superne ramosa : foliis petiolatis, ovalibus, crenato-dentatis; infimis suborbiculatis : spicis aphyllis, laxifloris; floribus parvulis, pedicellatis.

HAB. in Virginia.

CRASSIUSCULA. L. erecta, subpubescens, lucidula : foliis lanceolatis, serrulatis, crassiusculis : spica floribus alternis, subsessilibus : calycum tubo hispido, laciniis margine reflexo dentatis.

L. glandulosa ? WALT.

Obs. Variat foliis latioribus et ovalibus : caule sæpius simpli, rarius ramoso.

HAB. in paludosis herbosis Carolinæ maritimæ usque ad Floridam.

SYNGENESIA. MONOGAMIA. 153

AMOENA. L. majuscula, erecta, glaberrima : foliis lato-lanceolatis, serratis : spica multiflora, secunda : calycis laciniis integerrimis : corollæ cœruleæ laciniis inferioribus ovalibus, acutis.

CLAYTONIANA. L. erecta, simplex, minutim pubescens : foliis oblongis, obtusis, subintegris ; imis spathulatis : spica nuda, virgata, parviflora.

RAPUNTIIUM foliis oblongis, villosis, mollibus, margine leviter crenatis, &c. CLAYT. *Fl. Virg.*

OBS. Proxima *L. urenti*. Calycum laciniae tripla tubi brevissimi longitudine.

HAB. a Pennsylvania ad Carolinam.

KALMII. L. parvula : tenuis, erecta : foliis radicalibus spathulatis ; caulinis linearibus, obtusiusculis, subintegris : spica floribus distanter alternis, pusillis, longiuscule pedicellatis.

OBS. Caulis plerumque simplex : folia armato oculo remote subdenticulata. Varietas Caroliniana foliis radicalibus suborbiculato-spathulatis ; caule graciliore ; floribus minoribus.

HAB. in Canada et Carolina.

DORTMANNA. L. foliis radicalibus linearibus, recurvis, bilocularibus, integerrimis : caule

154 SYNGENESIA. *MONOGAMIA*.

simplicissimo, erecto, subaphyllo : spica  
floribus remote alternis, pendulis.

*НЛВ.* in lacu *Senogamis* aliisque vicinis, in terri-  
torio sinus *Hudsonis*.

---

CLASSIS XX.

GYNANDRIA.

MONANDRIA.

ORCHIS. L.

*Nectarium* corniforme pone florem.

CLAVELLATA. O. bulbis tenuiter fusiformibus ;  
scapo oblonge unifolio : spica laxiuscule  
pauciflora ; bracteis brevibus : calyce  
minuto , connivente ; cornu longitudine  
ovariorum , clavato ; labello ovali , integro.

HAB. in Carolina.

HUMILIS. O. foliis ad radicem binis , latissime  
obovalibus , obtusissimis : spica rariflora ;  
bracteis ovali-lanceolatis , foliaceis : cornu  
longitudine ovariorum ; laciniis quinque-  
cullatim conniventibus ; labello depen-  
dente , subovali.

OR. Flores purpurei.

HAB. in excelsis montibus Carolinae.

QUINQUESETA. O. foliis ovalibus , acutis : spica  
floribus distanter alternis ; bracteis acu-  
minatis : cornu dupla ovariorum longitu-



156 GYNANDRIA. MONANDRIA.

dine, subbiunciali : labello in quinque lacinias setaceas partito.

*Obs.* Affinis *O. habenariae*.

*HAB.* in Carolina.

**LACERA.** *O.* foliis spicaque oblongis ; floribus distincte alternis : cornu fere ovarii longitudine : labello longiore, anguste tripartito ; laciniis subdigitatis, filiformibus.

*HAB.* in Carolina.

**CRISTATA.** *O.* bulbis indivisis : foliis lanceolatis : spica floribus confertis, parvulis, luteis : cornu dimidii ovarii longitudine : laciniis duabus interioribus rotundatis, cristato-denticulatis : labello oblongo, pennatim lacero.

*Obs.* Affinis *O. ciliari*. Ovarium itidem superne setaceo-attenuatum.

*HAB.* in sylvis Carolinæ.

**CILIARIS.** *O.* scapo folioso : spica subovata, multiflora : ovario superne in setam bractea longiorem attenuato : cornu subunciali : labello sublanceolato, pennatim lacero.

Var. *α. lutea*.

— *β. alba*.

*HAB.* a Canada ad Carolinam.

SATYRIUM. L.

*Nectarium* scrotiforme seu inflato-didymum pone  
florem.

REPENS. S. radice fibrosa, repente : foliis ra-  
dicalibus, ovalibus, venoso-maculosis :  
scapo aphylo spicaque pubentibus.

HAB. a Canada ad Floridam.

MALAXIS. SWARTZ.

*Nectarium* in medio corollæ, petalis minus, con-  
cavum; marginibus convexis, cordatum, pos-  
tice acuminatum, antice bifidum.

UNIFOLIA. M. scapo unifoliato : foliolo cordato-  
ovali : spica oblonga.

HAB. in umbrosis sylvarum, a Carolina ad Floridam.

OPHRYS. L.

*Nectarium* subtus subcarinatum.

ÆSTIVALIS. O. scapo folioso : foliis glabris, lan-  
ceolatis, acutissimis : spica pubescente,  
spirali; floribus albis : calyce quasi in  
tubum recurvum connivente; labello  
crenulato.

LIMODORUM *praecox*? WALT.

ORCHIS spiralis, alba, odorata, longo angus-  
toque folio. VAILL. *Bot. Paris.*

HAB. a Pennsylvania ad Carolinam.

158 GYNANDRIA. MONANDRIA.

CERNUA. O. foliis radicalibus, lineari-oblongis,  
L. obtusiusculis : scapo alterne vaginoso,  
puberulo : spica spiraliter subimbricata,  
pubescente : ovariis ovatis ; calycibus re-  
clinatis.

LIMODORUM? autumnale. WALT.

Obs. Affinis *O. aestivali*, spica confertiore, floribus  
subduplo majoribus.

HAB. a sinu Hudsonis ad Carolinam.

PUBERA. O. foliis ad radicem 3-4, ovalibus,  
subacutis : scapo aphylo, pubescente :  
spica laxa ; floribus pedicellatis : ovariis  
fructibusve clavatis : labello bilobo, vix  
calyce connivente longiore.

Obs. Affinis *O. ovatae*.

HAB. in Georgia.

CORDATA. O. radice fibrosa : caule gracili, me-  
L. dium versus oppositè bifolio ; foliis bre-  
vibus, cordatis : spica parviflora ; ova-  
rio subgloboso : labello oblongo, subse-  
taceo-bifido.

HAB. in Nova Anglia.

CORALLORHIZA. O. radice flexuose ramosa, den-  
L. tata scapoque aphyllis : spica laxa : la-  
bello trifido.

PLUCKN. t. 211. f. 2.

HAB. in Canada et Nova Anglia.

LIMODORUM. L.

*Nectarium* monophyllum, concavum, pedicellatum, intra petalum infimum.

TUBEROSUM. L. radice bulbosa : scapo supra basim  
L. subvaginatam unifoliato ; foliolo lineari-gramineo : spica simplici, floribus paucioribus, alternis, purpureis : labello promisse barbato.

SMITH. *ic.*

HAB. a Canada ad Floridam.

TRIFIDUM. L. foliis lanceolatis : scapo aphylo ;  
labello trifido, longitudinaliter lamellato.

Obs. Jamdudum in hortis Europæis cultum et perperam pro *L. tuberoso* traditum.

HAB. in Bahama ; cultum in Carolina, autumnomihifloruit.

ARETHUSA. L.

*Nectarium* tubulosum intra corollæ fundum : labio inferiore stylo adnato.

OPHIOGLOSSOIDES. A. radice fibrosa : scapo dissite  
L. bifoliato, 1-2-floro : foliis ovali-oblongis : labello fimbriato.

HAB. a Canada ad Carolinam, in maritimis.

160 GYNANDRIA. MONANDRIA.

BULBOSA. A. aphylla : radice bulbosa : scapo vaginato, unifloro : calyce laciniis superioribus incurvatis : labello subcrenulato.

HAB. a Canada ad Carolinam montosam.

DIVARICATA. A. radice fibrosa : scapo remote bifoliato, unifloro : foliis oblongis, obtusiusculis : calyce laciniis exterioribus patulis, longo-linearibus : labello eroso - subcrenulato.

HAB. a Carolina ad Floridam, in maritimis.

PARVIFLORA. A. parvula : scapo foliis pluribus, alternis, brevi-ovalibus ; summitate solito pluriflora ; labello integro.

HELLEBORINE mariana, flore pallide purpureo, trianthophoros. PLUCKN. *Mant.* 100. t. 348. f. 6.

HAB. in Kentucky umbrosis.

GYNANDRIA. MONANDRIA. 161

CYPRIPEDIUM. L.

*Nectarium* ventricosum, inflatum, cavum.

CANADENSE. C. totum hirsutum : caule folioso : laciniis calycis exterioribus lato-ovalibus, obtusis ; calceolo purpureo.

HAB. in Canada et occidentalibus *Alléghanis*, circa *Knoxville*.

CALCEOLUS. C. minutim pubescens : caule folioso :  
L. laciniis calycis exterioribus oblongo-ovalibus, acuminatis ; interioribus linearibus confertisque : calceolo luteo.

PLUCKN. t. 418. f. 2.

HAB. a Nova Anglia ad Carolinam superiorem.

ACAULE. C. minute pubescens : foliis ad imum  
AIT. caulem binis, oblongis, non acuminatis : scapo nudo, unifloro : laciniis calycis exterioribus lanceolatis : calceolo purpureo.

HAB. a Canada ad Carolinam montosam.

HEXANDRIA.

ARISTOLOCHIA. L.

Hexagyna. Cal. o. Cor. 1-petala, lingulata, in-  
tegra. Caps. 6-locularis, infera.

SIPHO. A. fruticosa, volubilis : foliis petiolatis, amplissime cordatis : floribus soli-

152 SYNGENESIA. MONOGAMIA.

INFLATA. L. erecta, ramosa, hirsutissima : foliis ovali-lanceolatis, serratis : spicis foliosis; floribus pusillis : capsulis inflatis.

HAB. a Canada ad Carolinam montosam.

PUBERULA. L. erecta, simplicissima, pubescens : foliis oblongo-ovalibus, obtusis, repando-serrulatis : spica non pedunculata; floribus paucis, alternis, subsessilibus : calycibus ciliatis.

HAB. in Carolina.

CLIFFORTIANA. L. glabella, superne ramosa : foliis petiolatis, ovalibus, crenato-dentatis; infimis suborbiculatis : spicis aphyllis, laxifloris; floribus parvulis, pedicellatis.

HAB. in Virginia.

CRASSIUSCULA. L. erecta, subpubescens, lucidula : foliis lanceolatis, serrulatis, crassiusculis : spica floribus alternis, subsessilibus : calycum tubo hispido, laciniis margine reflexo dentatis.

L. glandulosa ? WALT.

Obs. Variat foliis latioribus et ovalibus : caule sæpius simplici, rarius ramoso.

HAB. in paludosis herbosis Carolinæ maritimæ usque ad Floridam.

SYNGENESIA. *MONOGAMIA*. 153

**AMOENA.** L. majuscula, erecta, glaberrima : foliis lato-lanceolatis, serratis : spica multiflora, secunda : calycis laciniis integerrimis : corollæ cœruleæ laciniis inferioribus ovalibus, acutis.

**CLAYTONIANA.** L. erecta, simplex, minutim pubescens : foliis oblongis, obtusis, subintegris ; imis spathulatis : spica nuda, virgata, parviflora.

*RAFUNTII* foliis oblongis, villosis, mollibus, margine leviter crenatis, &c. *CLAYT. Fl. Virg.*

*OBS.* Proxima *L. urenti*. Calycum laciniae tripla tubi brevissimi longitudine.

*HAB.* a Pennsylvania ad Carolinam.

**KALMII.** L. parvula : tenuis, erecta : foliis radicalibus spathulatis ; caulinis linearibus, obtusiusculis, subintegris : spica floribus distanter alternis, pusillis, longiuscule pedicellatis.

*OBS.* Caulis plerumque simplex : folia armato oculo remote subdenticulata. Varietas Caroliniana foliis radicalibus suborbiculato-spathulatis ; caule graciliore ; floribus minoribus.

*HAB.* in Canada et Carolina.

**DORTMANNA.** L. foliis radicalibus linearibus, recurvis, bilocularibus, integerrimis : caule



## PODOSTEMUM.

Πους, Στήμων : *stamina pedicello suffulta.*

Plantulæ aquatiles ; cauliculis coriaceis. Flores solitarii, pedicellati, monoici ; masculo adjecto feminae.

## F E M.

CAL. nullus.

PIST. nudum, hinc basi binis squamulis subulatis stipatum : ovarium ovatum : stigmata duo, sessilia, subfiliformia, illo paulo breviora, divergentia.

CAPS. ovata, 8-striata ; 2-locularis ; 2-valvis : dissepimentum valvis parallelum, sub dehiscencia liberum.

SEM. plurima, utramque dissepimenti medio crascescentis faciem fere obtegentia, deorsum subimbricata, ovoidea.


## M A S C.

Stipes sive pedicellus filiformis, inter binas squamulas pistillum stipantes oriundus, apicē gerit.

STAMINA duo, filamentis breviusculis basi contiguis : antheris subcordatis, bilocularibus.

OBS. Genus hinc quamdam cum RUPPIA præ se ferens affinitatem, inde vero fructu ab eadem et affinibus discedens.





P. J. Radou's Del.

Plas. Sc.

PODOSTEMON *ceratophyllum*.

MONOECIA. TRIANDRIA. 165

CERATOPHYLLUM. P. foliis breviuscule setaceo-  
tab. 44. multi-partitis.

HAB. in rupibus saltus fluvii Ohio, juxta Louis-  
ville.

TRIANDRIA.

ERIOCAULON. E.

Cal. communis, capitulum imbricatum. Pe-  
tala 3, æqualia. Stam. supra ovarium.

DECANGULARE. E. maximum, glabrum : foliis gra-  
mineo-ensiformibus : scapo 10-striato :  
capitulo pomiformi, floribus numero-  
sissimis : squamis involucri vix distincti  
ovalibus, subacutis ; paleis receptaculi  
mucronatis.

E. serotinum ? WALT.

HAB. in Pennsylvania, Virginia et Carolina.

GNAPHALODES. E. glabrum : foliis brevibus, su-  
bulato-ensiformibus ; vagina iis duplo  
triplove longiore : scapo elato, 10-striato :  
capitulo subhemisphærice convexo : in-  
volucri squamis ovalibus, rotundato-ob-  
tusis, scariosis, argenteo-lucidis.

E. decangulare ? WALT.

HAB. in Carolina.

166      **MONOECIA. TRIANDRIA.**

**PELLUCIDUM.** *E. pusillum*, glabrum : foliis brevibus, subulatis, canaliculatis, pellucidis, 5-nerviis : caulibus solitariis : capitulo parvulo, subpomiformi : squamis involucri vix distincti ovalibus, obtusis.

*HAB.* in septentrionalibus Canadae.

**VILLOSUM.** *E. foliis breviusculis*, subulato-linearibus, pilosis : scapis sub4-sulcis, villosis : capitulo sphæroideo; flosculis subfuliginosis.

*E. anceps?* WALT.

*HAB.* in Carolina.

**FLAVIDULUM.** *E. minutissime puberulum* : culmis aggregatis, 5-striatis : foliis brevissimis, subulato-ensiformibus : capitulo convexo : flosculis vix pappulosis.

*HAB.* in Carolina.

## SCLERIA. BERG. GÆRT.

*Glumæ* simplices, undique imbricatæ in spiculis.  
*Spicæ* masculæ multifloræ : femineæ unifloræ :  
femineis *stigm.* 1 - 3. *Semen* globosum, sub-  
osseum.

RETICULARIS. *S. erecta*, glaberrima levissimaque :  
fasciculis lateralibus et terminali remo-  
tissimis, subsessilibus, subramosis, pau-  
cifloris : semine globoso, reticulato,  
foveolis consperso.

*HAB.* in Carolina.

CILIATA. *S. glabriuscula*, stricte erecta : foliis  
angustissimis, longissimis, partim mi-  
nutissime ciliatis : fasciculo unico ter-  
minali, fusco, paucifloro ; bracteis brac-  
teolisque ciliatis : semine globoso, sca-  
berulo.

*HAB.* in Carolina.

OLIGANTHA. *S. gracilis*, debilis, glabra : culmo  
angulis acutissimis, lævibus : foliis an-  
gustissimis, asperis : pedunculis latera-  
libus solito duobus, inter se dissitis,  
setaceis, subunifloris ; floribus termi-  
nalibus paucissimis, a se invicem dis-  
tinctis.

*HAB.* in pratensibus sylvaticis Carolinae.

168      **MONOECIA. TRIANDRIA.**

**TRIGLOMERATA.** S. glabra , angustifolia : culmo  
    acutissime triquetro , angulis asperis :  
    fasciculo terminali , subtriglomerato ;  
    glomerulis sessilibus , alternis.

*HAB.* in Carolina.

**HIRTELLA.** S. gracilis , angustifolia : foliis bracteisque  
    hirsutulis : fasciculo terminali ,  
    paucifloro.

*HAB.* in sylvis Carolinae.

**INTERRUPTA.** S. parvula , tota pubens : spica e  
    glomerulis distincte alternis , sessilibus ,  
    ebracteatis , paucifloris , hispidis.

*OBS.* Eandem in Gallo-Guyanna legit L. C. RICHARD.

*HAB.* a Carolina ad Floridam , in pratis udis.

CAREX. L. Juss.

*Flores* masculi femineis mixti, aut in spica distincta superiore segregati. Femineis *stigmata* 3, interdum 2. *Semen* tunicatum, villis destitutum.

§. I. SPICIS ANDROGYNIS.

—A. SPICULA UNICA.

**MICROSTACHYA.** C. pusilla : foliis culmisque subæqualibus, setaceis : spica unica, minuta ; floribus alternis, superioribus masculis : capsulis erectis, muticis.

**TYPHINA.** C. foliis culmum superantibus : spicula unica, crassissima, oblongo-ovata; capsulis densissimis, turbinato-inflatis.

*HAB.* in regione Illinoensi.

—B. SPICULIS PLURIBUS.

**VULPINOIDEA.** C. culmo aspero : foliis angustis : spica subinterrupta; spiculis androgynis, glomerato-sessilibus, setaceo-7-bracteatis : capsulis parvulis, introrsum planiusculis, acuminatis.

*Obs.* Valde affinis *C. vulpinae*, omnibus partibus multo minor; capsulis uti in *C. muricata* divergentibus.

*HAB.* in Canada et Nova Anglia.



170 MONOECIA. TRIANDRIA.

LEPORINA. C. spicis androgynis, in summo culmo

L. proxime alternis, distinctis, sessilibus, turgide obtuseque ovatis, ebracteatis: capsulis compacte imbricatis, convexo-planis, acuminatis.

OBS. Plerumque spiculis subviridantibus aut subflaviscantibus.

HAB. a Canada ad Carolinam.

RICHARDI. C. foliis culmisque erectis: spiculis

THUILL. androgynis, brevibus, ovatis, obtusis, pallide-viridulis, ad summitatem proxime distincteque sessilibus, subebracteatis, ovoideis.

C. Richardi. THUILLIER. Fl. Paris.

HAB. in borealibus Canadae.

TRICEPS. C. angustifolia: culmo superne aspero: spiculis tribus, approximatis, sessilibus; capsulis ovoideis, muticis.

OBS. Culmus folio infra flores iis multo longiore et sterili munitus. Spicula terminalis inferne mascula.

HAB. in Carolina.

VIRIDULA. C. pusilla, erecta: angustifolia: culmo laevi, spiculis approximatis, sessilibus; duabus inferioribus axillaribus; suprema etiam sessili, inferne mascula: capsulis triquetris, acuminatis.

OBS. Affinis C. tricipiti et flavae.

HAB. in Canada.

MONOECIA. TRIANDRIA. 171

§. II. SPICIS SEXU DISTINCTIS.

—A. MASCULA UNICA.

SCIRPOIDEA. C. planifolia, dioica : spica unica, imbricato-cylindrica : capsulis dense pubescentibus.

*HAB.* ad sinum *Hudsonis*.

FLAVA. C. diffusa, lævigata : culmis folia superantibus : spiculis fem. 2-3, turgide ovoideis, bracteatis, subsessilibus : capsulis longius acuminatis, subrecurvo-patentibus, maturitate flavis.

*HAB.* in borealibus *Canadæ*.

OEDERI. C. humilis, diffusa, lævis : foliis culmos superantibus : spiculis fem. pluribus, juxta masculam sessilem conglomerato-approximatis, subrotundo-ovoides, viridibus demumque flavis : capsulis brevius acuminatis.

*C. conglomerata.* THUILL. *Fl. Paris.*

*OBS.* Perperam cum *C. flava* ab auctoribus nonnullis conjuncta.

*HAB.* ad ripas amnis *des Goëlands* dieti, in sinum *Hudsonis* defluentis.

FOLLICULATA. C. foliis culmo multo longioribus, L. floralibus longissimis : spiculis fem. 1-3, approximatis, sessilibus, subglobosis seu

## MONOECIA. TRIANDRIA.

turgide ovoideis; mascula unica : capsulis omnium maximis, oblongo-triquetris, acuminatis.

*Obs.* Spica mascula modo sessilis, modo pedunculata.

*HAB.* in regione Illinoensi, Virginia et Carolina.

**DEBILIS.** C. foliis angustis : culmis debilibus : spiculis fem. pluribus, gracilibus, longo pedunculo recurvo - pendulis : capsulis lineari-lanceolatis, utrinque acutis; spicula mascula filiformi.

*Obs.* Affinis *C. sylvaticae* HUDSONIS.

**LENTICULARIS.** C. foliis angusto-longis, culmo gracili, triquetro subæqualibus : spiculis fem. pluribus, pedunculatis, oblongis; masculæ unica : capsulis lenticularibus, brevi-ovatis, muticis.

*Obs.* Affinis *C. paniceae*.

*HAB.* per tractus montium, a sinu *Hudsonis* ad *Canadam*, præsertim ad lacum *Cynorum* dictum.

**PAUPERCULA.** C. foliis angusto-longis, planis, erectis : culmo rectissimo, triquetro : spiculis pendulis; fem. subglobosis, squamis acutissimis; masc. unica : capsulis paucissimis, brevi-ovatis, muticis.

*HAB.* ad lacum *Mistassins*.

MONOECIA. TRIANDRIA. 173

STRIATULA. C. foliis brevibus, linearibus, planis : spiculis fem. plerumque 2, remotis, axillaribus, subpedunculatis, erectis; mascula unica, squamis rotundato-obtusis : capsulis paucioribus, subimbricatis, triquetris, utrinque acutis, striatis, glabris.

Obs. Habitus *C. paniceae*.

HAB. in Carolina.

ROSTRATA. C. stricte erecta : foliis angustissimis, planis : spicis fem. durabus, distinctis, axillaribus, subsessilibus, subglobosis, flavicantibus; mascula sessili : capsulis capitatis, erectis, oblongis, longissime rostratis.

HAB. in Canada et ad lacus *Mistassins*.

SUBULATA. C. foliis planis : culmis gracilibus, lævibus : spiculis fem. 3-4, axillaribus, valde remotis, sessilibus, suprema masculæ approximata : capsulis paucissimis, divaricato-reflexis, subulatis.

HAB. in Canadæ borealibus.

PLANTAGINEA. C. foliis radicalibus, lanceolatis, LAM. nervosis : culmo subaphyllo; vaginis remotis, monostachyis : spicis fem. gra-

ciliter linearibus, non vaginatis : mascula unica.

*Obs.* Variat 1°. vaginis spicisque castaneis et viridulis; 2°. vaginis aphyllis aut in foliolum desinentibus.

*HAB.* in Virginia et Carolina.

**MILIARIS.** C. erecta, gracilis : foliis planis, angustissimis : culmis superne asperis : spica fem. sessili, ovoidea, fusca : bractea setacea, brevi : mascula pallida, longius pedicellata : capsulis sphæroideis, lævibus.

*Obs.* Interdum spicæ fem. duæ, distantes, inferiore etiam sessili.

*HAB.* in paludosis borealibus Canadæ, præsertim ad lacus *Mistassins* dictos.

—B. MASCULIS PLURIBUS.

**OLIGOSPERMA.** C. foliis involuto-junceis culmisque erectis : spicula fem. unica, globulari, sessili; bractea setacea spicam masculam linearem, quandoque duplicem subæquante : capsulis paucissimis majusculis, turgide ovatis, acutis.

*HAB.* in Canada.

**STRIATA.** C. foliis angustis : spiculis fem. 1-2, distantibus, oblongis; squamis subalbidis, ovalibus : capsulis ovoideo-triquetris, pubescentibus; rostro bicorni : masculis 2-3, purpureo-subalbidis.

*HAB.* in Carolina.

MONOECIA. TRIANDRIA. 175

LANUGINOSA. C. culmo erecto, superne asperiusculo foliisque glabris : spiculis fem. 2, distantibus, axillaribus, arcte sessilibus, ovaleo-oblongis; masculis duabus, teretibus, pallidis : capsulis densissimis, brevibus, acuminatis, minuta lanugine tomentosis.

HAB. ad lacus Mistassins.

TRAGIA, L.

MASC. Cal. 3-partitus. Cor. o.

FOEM. Cal. 5-partitus. Cor. o. Stylus 3-fidus.

Caps. 3-cocca, 3-locularis. Sem. solitaria.

OBS. A Plumeriana specie meæ differunt spicis oppositifoliis et numero staminum terno, qui binus tantum in priore.

URENS. T. erecta : caule minutissime puberulo : foliis sessilibus, ovali-lanceolatis linearibusve, rariter dentatis : capsulis hirsutulis.

L.

Var.  $\alpha$ . *subovalis* : foliis oblongo-ovalibus, nonnullis subcuneatis.

—  $\beta$ . *lanceolata* : foliis lanceolatis, subdentatis integrisque.

T. *innocua*. WALT.

—  $\gamma$ . *linearis* : foliis linearibus, fere omnibus integris.

HAB. in Carolina et Georgia.

176      **MONOECIA. TRIANDRIA.**

**URTICIFOLIA.**    *T. erecta* : caule hirsutissimo : foliis cordato-ovalibus.

*Obs.* Huc, ni fallor, synonyma *GRONOVII* et *CLAYTONI* perperam ad *T. Mercurialem* refert *LINNEUS*.

*HAB.* in Georgia.

**CORDATA.**    *T. scandens*, hispidula : foliis latioribus, eximie cordatis, acute dentatis : capsula majuscula.

*Obs.* Capsula in genere grandis, coccis majoris *Pini* crassitie.

*HAB.* in Kentucky.

**TETRANDRIA.**

**PACHYSANDRA.**

Πάχυς, Ἀνρ : *stamina crassa.*

Folia alterna. Spicæ ex ima caulis herbacei parte nuda, oblongæ; floribus majusculis, sessilibus; paucissimis inferioribus foemineis, cæteris masculis.

**M A S C.**

BRACTEA squamiformis, calyci incumbens, paulo minor et ipsi consimilis.

CAL. sub4-phyllus; foliolis ovalibus; duobus paulo interioribus.

STAM. 4 : filamenta tripla calycis longitudine, erecta, crassissima, compresso-clavata : antheræ dorso affixæ, oblongæ, 2-loculares; demum arcuatae.

**F O E M.**

BRACTEÆ 3, squamiformes; una exteriore.

CAL. fere maris, minor; ovario appressus.

PIST. ovarium subrotundum, 3-sulcatum : styli 3, recurvi, sulco exarati; desinentes in totidem stigmata ligulata.



178 MONOECIA. TETRANDRIA.

CAPSULA subglobosa , 3-cocca , stylis persistentibus 3-cornis , 3-ocularis.

SEMINA in singulis loculis bina , oblonga , summo loculo appensa , lævigata.

OBS. Character fere Buxi : habitus diversissimus.

PROCUMBENS. P. caule procumbente : foliis brevibus , ovalibus , superne crenato-dentatis : calycibus minutim ciliatis : capsula puberula.

HAB. in occidentalibus montium *Alleghanis*.

U R T I C A. L.

MASC. Cal. 4-phyllus. Cor. o. Nectarium centrale , cyathiforme.

FOEM. Cal. 2-valvis. Cor. o. Sem. 1 , nitidum.

CANADENSIS. U. foliis alternis , lato-ovalibus : racemis subpaniculatis ; inferioribus sessilibus , masculis , plerumque pentandris ; superioribus pedunculatis , foemineis aut submonoicis.

OBS. Synonymum PLUCKNETI ad *U. divaricatam* relatum huc pertinet.

HAB. in Canada et ad ripas fluminis *Ohio*.

PUMILA. U. glabella , lucida : caule carnosissimo : foliis ovalibus , sub3-nerviis : racemis



P. J. Redoute' Del.

Plée. Sc.

**PACHYSANDRA** *procumbens*.



**MONOECIA. TETRANDRIA. 179**

monoicis, 3-andris, brevissimis, sessilibus.

*Obs.* Altitudine variat; 4-uncialis ad 2-pedalem.

*HAB.* copiose ad occidentem montium *Alléghanis* et in Carolina.

**CYLINDRICA.** U. herbacea: foliis oblongo-ovalibus,

*L.* subtrinerviis: spicis simplicissimis, dioicis; masculis glomerato-interruptis; foemineis cylindraceis.

*HAB.* in dumetosis, a Canada ad Floridam.

**DIOICA.** U. foliis oppositis, oblongiuscule cor-

*L.* datis: racemis geminis, compositis, elongatis, patulis; plerumque dioicis.

*Obs.* Sæpius vidi monoicam.

*HAB.* in Canada.

**MORUS. L.**

*MASC. Cal.* 4-partitus. *Cor.* o.

*FOEM. Cal.* 4-phyllus. *Cor.* o. *Styli* 2. *Cal.* baccatus. *Sem.* 1.

**RUBRA.** M. foliis cordatis, longius acuminatis,

*L.* æqualiter serratis, subtus pubentibus: spicis foemineis cylindraceis.

*Obs.* Interdum mere dioica.

*HAB.* a Canada ad Floridam.

BETULA. L.

MASC. *Cal.* 1-phyllus, 3-fidus, 3-florus. *Cor.* 4-partita.

FOEM. *Cal.* 1-phyllus, sub3-fidus, 2-florus. *Sem.* utrinque membrana alata.

NANA. B. humillima, glaberrima : foliis perpusillis, subcuneato-orbiculatis, inciso-crenulatis, reticulato-venosis : amenti squamis profunde 3-partitis, laciniis oblongis : capsulis orbiculatis, subapteris.

HAB. in sphagnosis, a sinu *Hudsonis* ad lacus *Mistassins*.

GLANDULOSA. B. pumila : ramis glanduloso-punctatis, glabris : foliis subrotundo-obovalibus : amenti squamis semi3-fidis : capsulis orbiculatis, anguste marginatis.

Obs. Affinis *B. pumilae*.

HAB. circa lacus, a sinu *Hudsonis* ad *Mistassins*.

PAPYRIFERA. B. foliis ovalibus, acuminatis, subæqualiter serratis : petiolo glabro : amento fem. cylindrico ; squamis intus minutissime puberulis ; lobis lateralibus multo brevioribus, suborbiculatis, non divergentibus.

*B. papyracea*. AIT. *Hort. Kew.*

MONOECIA. TETRANDRIA. 181

**LANULOSA.** B. foliis deltoideo - ovalibus , non acuminatis , duplicato-serratis ; petiolo brevi : amento fem. subovoideo ; squamis extus dense lanulosis ; lobis subæqualibus , oblongis , divergentibus.

*Obs.* Videtur *B. nigra* Auctorum ; cujus nomen , ideam falsam movens , immutavi.

**CARPINFOLIA.** B. petiolis brevissimis , puberulis ; foliis sinu profundo cordatis , oblongiuscule ovalibus , acuminatis ; nervis oblique parallelis : amento fem. subsessili , cylindrico ; squamis glabris ; lobis brevibus , subæqualibus , rotundatis , divergentibus.

*HAB.* in montosis Canadæ , Pensylvaniæ et alibi.

**SERRULATA.** B. Alnus : stipulis ovalibus , obtusis ,  
*AIT.* foliis lato-ovalibus , acutis : amentis foem. oblongiusculis.

*Obs.* Folia sæpe obovalia , interdum subglanduloso-repanda , basi semper acuta.

*HAB.* a Pensylvania ad Carolinam.

**CRISPA.** B. Alnus : foliis ovalibus , subacutis ,  
*AIT.* duplicato-serratis , subundulatis ; nervorum axillis subnudis : stipulis lato-ovalibus , obtusis.

*Obs.* Folia basi obtusiuscula , interdum etiam subcordata.

*HAB.* in Canada.

*PENTANDRIA*.*XANTHIUM*. *L*.

**MASC.** *Cal.* communis imbricatus. *Cor.* 1-petalæ,  
5-fidæ, infundibulif. *Recept.* paleaceum.

**FOEM.** *Cal.* involucrum 2-phyllum, 2-florum.  
*Cor.* o. *Drupa* sicca, muricata, 2-fida. *Nu-*  
*cleus* 2-locularis.

**STRUMARIUM.** *X.* inerme : foliis subcordatis, an-  
*L.* gulosis, asperis.

*Obs.* Fructus densius spinosus.

*HAB.* a Canada ad Carolinam.

*AMBROSIA*. *L*.

**MASC.** *Cal.* communis 1-phyll. *Cor.* 1-petalæ,  
5-fidæ, infundib. *Recept.* nudum.

**FOEM.** *Cal.* 1-phyllus, integer, ventre 5-dentato,  
1-florus. *Cor.* o. *Nux* e calyce indurato, 1-  
sperma.

**BIDENTATA.** *A.* hirsutissima : foliis arcte sessili-  
bus, simplicibus, oblongis, supra basim  
latescentem utrinque unidentatis : fructu  
4-gono, infra apicem 4-spinoso.

*Obs.* Involucra mascula superne ultra flosculos in mu-  
cronem desinentia.

*HAB.* in regione Illinoensi.

MONOECIA. *PENTANDRIA.* 183

TRIFIDA. *A. hirsuta, aspera* : foliis 3-partitis,  
L. serratis ; lobis ovali-lanceolatis , acuminatis : fructu infra apicem 6-spinoso.

*OBS.* Interdum foliis simplicibus variat

*HAB.* in Pennsylvania et Carolina.

ABSYNTHIFOLIA. *A. foliis subtus incanis, bipinnatifidis* ; sinubus laciniisque obtusis.

*OBS.* Fructus spinæ submediæ, acutæ , apice ejus conspicue breviores.

*HAB.* in montosis Carolina.

PANICULATA. *A. caule ramosissimo, superne paniculato petiolisque villosis* : foliis utrinque viridibus, bipinnatifidis ; laciniis lanceolatis : fructibus aggregatis , pusillis , globoso-obovatis , subinermibus.

*IVA monophylla.* WALT.

*OBS.* Affinis *A. elatiori*, quæ glabritie spicarum, fide LINNÆI, differt.

Variat foliis subsimpliciter pinnatifidis, uti in icone PLUCKNETI t. 10. f. 5 ; et sæpe, luxuriante planta, subtripinnatifidis.

*HAB.* a Canada ad Floridam.



I V A. L.

MASC. *Cal.* communis 3-phyllus. *Cor.* disci 1-petalæ, 5-fidæ. *Recept.* pilis distinctum.

FOEM. *Cal.* flosculi radii 5. *Cor.* 6. *Styli* 2. longi. *Sem.* obtusum.

ANNUA. I. annua, hirsuta : foliis ovali-lanceolatis, rariter serratis : spica confertiuscula; bracteis acuminatis involucrisque hirsutis.

*Obs.* Folia modo opposita, modo alterna, fere *BRUNELLE vulgaris*. Spica quasi *AMEROSILÆ*. Flores feminei intra singula involucria 3-5.

*HAB.* in regione Illinoensi.

IMBRICATA. I. suffruticosa, glabra : foliis crassis, *WALT.* cuneatim lineari-lanceolatis, integris : glomerulis florum squamis imbricatis.

*HAB.* in maritimis Carolinae et Georgiae.

FRUTESCENS. I. fruticosa : foliis lanceolatis, serratis : glomerulis florum depresso-globosis.

*Obs.* Occurrunt individua floribus omnibus masculis.

*HAB.* in maritimis, a Nova Angliâ ad Floridam.

CROTONOPSIS.

Κρότων, Οψίς : *facies* CROTONI.

Caulis herbaceus, dichotomus : folia alterna : spicilla terminalis ; floribus minutis, solitarie alternis ; inferioribus foemineis, superioribus masculis. Pubes miculacea, fimbriata, in omnibus partibus.

M A S C.

CAL. profundissime 5-partitus ; laciniis ovalibus, obtusis, concavulis.

COR. petala 5, calyce minora, alterna, lineari-oblonga.

STAM. 5 : filamenta longitudine calycis : antheræ 2-loculares ; loculis, interposita filamenti summitate incrassata, disjunctis.

F O E M.

CAL. maris ; laciniis subulatis.

COR. o.

PIST. ovarium ovoideum : stigmata 3, sessilia, brevissima, duplicato bifida.

CAPS. brevis, ovata, teres, 1-ocularis, non dehiscens.

186      **MONOECIA. POLYANDRIA.**

**SEMEN** unicum, subglobosum; summo loculo adnexum.

Embryo uti semen inversus, albumine crassissimo, oleaginoso-carnoso inclusus.

**OBS.** Genus **CROTONI** affine.

**LINEARIS.** C. caule erecto, dichotome ramosissim<sup>o</sup> : foliis linearibus, obtusis.  
*tab 46.*

**HAB.** in maritimis Carolinæ, juxta *Long-bay*, et in regione Illinoensi.

**P O L Y A N D R I A.**

**P O T H O S. L.**

*Spatha. Spadix simplex, tectus. Cal. o. Petala 4. Baccæ dispermæ.*

**FOETIDA.** P. acaulis : foliis ovalibus, concavis : spadice subgloboso.

**DRACONTIUM** *fœtidum.* LINN.

**OBS.** Forsan potius congener *CALLÆ palustri*; habitu et habitatione certe affinis.

**HAB.** a Canada ad Virginiam; aquatica.



P. J. Redouté Del.

Plée Sc.

**CRÓTONOPSIS** *linearis*.

1. The first part of the document is a list of names and addresses of the members of the committee. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

2. The second part of the document is a list of the names of the members of the committee who have been elected to the office of the chairman. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

3. The third part of the document is a list of the names of the members of the committee who have been elected to the office of the secretary. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

4. The fourth part of the document is a list of the names of the members of the committee who have been elected to the office of the treasurer. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

5. The fifth part of the document is a list of the names of the members of the committee who have been elected to the office of the clerk. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

6. The sixth part of the document is a list of the names of the members of the committee who have been elected to the office of the auditor. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

7. The seventh part of the document is a list of the names of the members of the committee who have been elected to the office of the assessor. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

8. The eighth part of the document is a list of the names of the members of the committee who have been elected to the office of the collector. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

9. The ninth part of the document is a list of the names of the members of the committee who have been elected to the office of the recorder. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

10. The tenth part of the document is a list of the names of the members of the committee who have been elected to the office of the clerk of the court. The names are listed in alphabetical order, and the addresses are given below each name. The list includes names such as Mr. J. H. Smith, Mr. W. B. Jones, and Mr. C. D. Brown, among others.

CALLA. L.

*Spatha* plana. *Spadix* tectus flosculis. *Cal.* o.  
*Petala* o. *Baccae* polyspermæ.

PALUSTRIS. C. foliis subrotundo-cordatis : spatha  
L. brevi-ovali.

*OBS.* Sexus uniformiter per totum spadicem oblongo-  
ovatum conjuncti.

*HAB.* in aquosis Canadæ.

SAGITTIFOLIA. C. foliis subhastato-sagittatis ; au-  
ribus oblongis : spatha cucullata, su-  
perne ovali.

*OBS.* Spadix longo-cylindraceus, a parte circiter media  
ad apicem masculus.

*HAB.* in paludosis Georgiæ et Floridæ.

VIRGINICA. C. foliis sagittato-hastatis, auribus  
obtusis : spatha elongata, incurva.

*OBS.* Spadix superne longius masculiflorus.

ARUM *virginicum*. LINN.

*OBS.* Variat foliis angusto-oblongis.

*HAB.* in aquosis, a Nova Anglia ad Virginiam.

A R U M. *L.*

*Spatha* monophylla , cucullata. *Spadix* supra nudus , inferne foemineus , medio stamineus.

**DRACONTIUM.** A. foliis pedatis : foliolis sublan-  
L. ceolato-oblongis , integerrimis.

*Obs.* *Spatha* parvula , *spadice* brevior.

*HAB.* in umbrosis Carolinæ et Floridæ.

**TRIPHYLLUM.** A. acaule : foliis 3 - foliolatis ; fo-  
L. liolis ovalibus , integerrimis.

*Obs.* *Spadix* *spatha* dimidio brevior.

Var. *α. virens* : *spatha* virescente.

— *β. atropurpurea* : *spatha* atropurpurea.

*HAB.* a Canada ad Carolinam.

SPARGANIUM. L.

MASC. *Amentum* subrotundum. Cal. 3-phyllus.  
Cor. o.

FOEM. *Amentum* subrotundum. Cal. 3-phyllus.  
Cor. o. *Stigma* 2 - fidum. *Drupe* exsucca,  
1-sperma.

ANGUSTIFOLIUM. S. foliis planis, angustissimis,  
caulem longissime superantibus, debi-  
liter fluitantibus.

OBS. Folia vix sesquilineam lata.

HAB. in omnibus Canadae.

SAGITTARIA. L.

MASC. Cal. 3-phyllus. Cor. 3-petala. *Filamenta*  
fere 24.

FOEM. Cal. 3-phyllus. Cor. 3-petala. *Pist.* multa.  
*Sem.* multa, nuda.

SAGITTIFOLIA. S. foliis sagittatis, auribus depen-  
dentibus : seminibus plano-compressis.

OBS. Folia interdum *S. obtusifoliae*; auribus vero de-  
pendentibus discrepantia. Occurrunt etiam scapi  
floribus omnibus masculis.

HAB. a Canada ad Carolinam.

EAUCIFOLIA. S. foliis lato-lanceolatis, inferne  
L. angustatis : seminibus planis, falcatis,  
rostratis.

HAB. in Carolina.



190 MONOECIA. POLYANDRIA.

GRAMINEA. S. foliis lanceolato-linearibus : capitulis pistillorum pusillis.

HAB. in Canada.

NATANS. S. foliis natantibus, oblongo-ovalibus, obtusis; infimis subcordatis.

Obs. Habitus ALISMÆ natantis.

HAB. in Carolina.

MYRIOPHYLLUM. L.

MASC. Cal. 4-phyllus. Cor. o. Stam. 8.

FOEM. Cal. 4-phyllus. Cor. o. Pist. 4. Stylus o. Sem. 4, nuda.

SPICATUM. M. spica nuda, interrupta : floribus L. masculis polyandris.

HAB. a sinu Hudson ad Mont-réal.

VERTICILLATUM. M. floribus omnibus verticillatis, L. bracteatis; bracteis pinnatifidis; inferioribus foemineis; superioribus masculis aut hermaphroditis imperfectis, 8-andris.

HAB. in Canada, juxta Quebec.

SCABRATUM. M. foliis pinnatifidis : floribus omnibus verticillato-axillaribus; superioribus masculis, 4-andris; inferioribus foemineis, fructu jugatim 8-gulato.

POTAMOGETON pinnatum? WALT.

HAB. in Carolina et Georgia.

MONOECIA. POLYANDRIA. 191

HETEROPHYLLUM. M. foliis inferioribus capillaceo-pinnatis ; superioribus floralibus ovalibus , argute serratis ; floribus masculis 6-andris.

POTAMOGETON *verticillatum*? WALT.

HAB. in Carolina et Georgia.

JUGLANS. L.

MASC. Cal. 1-phyllus , squamiformis. Cor. 6-partita. Filamenta 18.

FOEM. Cal. 4-fidus, superus. Cor. 4-partita. Styli 2. Drupa nucleo sulcato.

\* Amentis masculis simplicibus.

REGIA. J. foliolis subseptenis, ovalibus, abrupte acuminatis, subintegris , glabellis : fructu globoso , lævigato ; nuce corrugata.

L.

HAB. in *Mediae* sylvis, juxta mare *Caspium* vulgo *Ghilan*.

NIGRA. J. foliolis subcordatis, superne angustatis , serratis : fructu globoso , punctato , scabriusculo : nuce corrugata.

L.

HAB. a Pennsylvania ad Floridam.

CINEREA. J. foliolis basi rotundato-obtusis, subtus tomentosis, leviter serratis : fructu oblongo-ovato , apice mammoso ; nuce

192 MONOECIA. POLYANDRIA,

oblonga, acuminata, insigniter insculpto-  
scabrosa.

*HAB.* præsertim in montibus *Alléghanis* et in mari-  
timis Novæ Cesaræ, Pensylvaniæ, &c.

PTEROCARPA. *J.* foliolis plurimis, lanceolatis :  
spica numerose foeminiflora : fructu pu-  
sillo, supra basim transverse diptero.

*HAB.* in umbrosis humidis sylvarum, juxta mare  
*Caspium*.

\*\* *Amentis masculis compositis.*

OLIVÆFORMIS. *J.* foliolis plurimis, subpetiolatis,  
falcatis, serratis; impari longius petio-  
lato : fructu oblongo, prominulo-qua-  
drangulo; nuce olivæformi, lævi.

*OBS.* Hæc vera incolarum *Pacane* est.

*HAB.* in regione Illinoensium.

MUCRONATA. *J.* foliolis 7-9-nis, argute serratis,  
glabriusculis; impari breviter petiolato :  
fructu subrotundo-ovoideo, superne su-  
turis 4 prominulis; nuce subglobosa,  
lævi, longissime mucronata.

*HAB.* in humidis sylvarum Virginæ, Carolinæ.

TOMENTOSA. *J.* foliolis septenis, leviter serratis,  
impari sessili; junioribus amentisque  
eximie tomentosis : fructu globoso, lævi;  
pericarpio crassissimo; nuce sexangu-  
lata.

*HAB.* a Pensylvania ad Carolinam.

MONOECIA. POLYANDRIA. 193

ALBA. J. foliolis quinis, conspicue serratis ;  
impari subsessili : squamis amenti atratis,  
lacinula terminali elongata : fructu sub-  
turbinato-globoso ; nuce ex utraque facie  
triangulata.

HAB. in Pennsylvania, Virginia et Carolina.

CASTANEA. TOURN. GÆRT.

MASC. Cal. subhexaphyllus. Cor. nulla. Stam. 5-20.

HERM. Cal. 5-6-phyllus. Cor. nulla. Stam. sub-  
duodena, in tomento latitantia. Styli 6, cartila-  
ginei. Ovarium inferum, 6-loculare : ovulis rostel-  
latis, pendulis, in singulo loculamento geminis.  
Nux matura 1-locularis, evalvis. Sem. 1-3.

VESCA : americana. C. foliis lanceolatis, acuminatis,  
serratis.

OBS. Ab Europæa vix distinguenda ; foliis paulo latio-  
ribus, subtus inter nervos non puberulis.

HAB. a New-Yorck ad Carolinam montosam.

PUMILA. C. foliis oblongo-obovalibus, serratis,  
subtus incano-tomentosis.

OBS. Variat serratura plus minus expressa ; foliis ro-  
tundato-obtusis seu acutis, interdum sublanceo-  
latis.

HAB. a Marilandia ad Floridam.

F A G U S. TOURN. GERT.

MASC. *Cal.* campanulatus, subsexfidus. *Cor.* nulla.  
*Stam.* 8.

FOEM. *Cal.* superus, hexaphyllus, tomentosus. *Cor.*  
nulla. *Stylus* unicus, 3-fidus. *Ovarium* imma-  
turum 3-loculare : ovulis subulato-caudatis, in  
singulo loculamento geminis. *Nux* matura 1-lo-  
cularis. *Sem.* 1-2.

SYLVESTRIS. F. foliis ovalibus, obsolete serratis.  
GERT.

Q U E R C U S. L.

MASC. *Cal.* 4-5-fidus. *Cor.* o. *Stam.* 4-10.

FOEM. *Cal.* plerumque 6-dentatus. *Cor.* o. *Ovarium*  
inferum, 3-loculare; *Stylus* 1. *Stigm.* 3. *Glans*  
ovata aut globosa, 1-loc., 1-sperma; rarissime  
2-3-sperma.

OBTUSILOBA. Q. foliis subtomentosis, profunde  
sinuato-lobatis, lobis retusis; basi acute  
cuneata : fructu mediocri; cupula crate-  
rata; glande brevi-ovata.

HAB. a Canada et Nova Anglia ad Floridam.

MACROCARPA. Q. foliis subtomentosis, profunde  
lyratimque sinuato-lobatis: lobis obtusis,  
subcrenato-repandis : fructu maximo :

MONOECIA. POLYANDRIA. 195

cupula profundius craterata, superne  
crinita; glande turgide ovata.

*HAB.* in montibus *Alleghanis*, Kentucky, Tennesseé,  
regione Illinoensi, &c.

LYRATA. Q. foliis subsessilibus, glabris, lyrato-  
WALT. sinuosis; summitate dilatata, divaricato-  
triloba; lobis acutangulis, terminali tri-  
cuspidate: cupula depresso-globosa, mu-  
ricato-scabrata; glande subtecta.

*HAB.* in Carolina meridionali, Georgia, &c.

ALBA. Q. foliis subæqualiter pinnatifidis; la-  
L. ciniis oblongis, obtusis, plerumque in-  
tegerrimis: fructu majusculo; cupula  
craterata, tuberculoso-scabrata; glande  
ovata.

Var. *α. pinnatifida*: foliis pinnatifidis, lævi-  
bus; lobis sinus subæquantibus; supra sa-  
turate viridibus, subtus glaucis; glandibus  
magnis, ovatis. WALT. *Car.* p. 235, n° 10.

— *β. repanda*: foliis repandis.

*HAB.* *α.* a Canada ad Floridam.

— *β.* in sylvis Carolinæ.

PRINUS. Q. foliis oblongo-ovalibus, acumina-  
L. tis acutisve, subuniformiter dentatis,

deciduis : cupula craterata, subsquamosa ; glande ovata.

Var. *α. palustris* : foliis longiuscule petiolatis, obovalibus ; fructu magno ; cupula modice concava ; conspicue squamosa.

— *β. monticola* : foliis brevi-petiolatis, subrhombico-ovalibus ; fructu majusculo, cupula turbinata, scabrosa ; glande oblonga.

— *γ. acuminata* : foliis longe petiolatis, basi obtusis, acutissime serratis ; fructu mediocri ; cupula subhæmisphærica.

— *δ. pumila* : foliis modice petiolatis, sublanceolatis, subtus glaucis.

— *ε. tomentosa* : foliis subsessilibus, obovalibus, dentibus obtusissimis ; subtus tomentosis.

*HAB.* *α.* in utraque Carolina, Georgia, Florida, &c.

— *β.* a Massachusetts ad Virginiam et in altis montibus utriusque Carolinæ.

— *γ.* in montibus *Alléghanis*.

— *δ.* in occidentalibus Virginis et Carolinæ.

— *ε.* in regione Illinoensi.

**VIRENS.**     *Q.* foliis perennantibus, coriaceis ; ovali-oblongis ; junioribus dentatis, vetustioribus integris : cupula turbinata, squamulis abbreviatis ; glande oblonga.

*AIT.*

*HAB.* a Virginia ad Floridam.

MONOECIA. POLYANDRIA. 197

PHELLOS. Q. foliis lineari-lanceolatis, integerrimis, glabris, apice setaceo-acuminatis; junioribus dentatis lobatisve : cupula scutellata ; glande subrotunda.

Var. *α. sylvatica* : foliis angusto-lanceolatis, utrinque acutis, deciduis.

— *β. maritima* : foliis latiuscule lanceolatis, perennantibus.

— *γ. pumila* : fruticulosa : foliis oblongis, basi obtusis.

HAB. *α.* a New-Jersey ad Floridam.

— *β.* in Carolina.

CINEREA. Q. foliis petiolatis, lanceolato-oblongis, acutis, integerrimis; subtus cinereo-pubescentibus : cupula scutellata, squamis marginalibus introrsum manifestis ; glande sphaerica.

HAB. in utraque Carolina et Georgia.

IMBRICARIA. Q. foliis subsessilibus, ovali-oblongis, acutis, integerrimis, subtus pubescentibus.

OBS. Fructus praecedentis : squamæ cupulae paulo majores.

HAB. in montibus Alleghensis.

LAURIFOLIA. Q. foliis subsessilibus, ovali-lanceolatis, inferne angustatis, acutis, inte-



198 MONOECIA. POLYANDRIA.

gerrimis, glabris; cupula subturbinata;  
glande subglobosa.

*HAB.* in Carolina meridionali et Georgia.

AQUATICA. Q. foliis obovali-cuneatis, basi acutis;

*CATESB.* summitate subintegris varieve trilobis,  
glabris; cupula modice craterata; glande  
subglobosa.

*HAB.* a Marilandia ad Floridam.

NIGRA. Q. foliis coriaceis, cuneatis, summi-

*CATESB.* tate dilatata retuso-subtrilobis, basi retu-  
sis; subtus rubiginoso - pulverulentis:  
cupula turbinata, squamis obtusis, sca-  
riosis; glande brevi-ovata.

*HAB.* a Marilandia ad Floridam.

TINCTORIA. Q. foliis petiolatis, subtus pubescen-

*BARTR.* tibus, lato-obovalibus, leviter et sub-  
rotunde lobatis, basi obtusis: cupula  
subscutellata aut turbinata; glande de-  
presso-globosa aut ovata.

Var. *α. angulosa*: foliis leviter lobatis, lobis  
angulosis: cupula subscutellata; glande de-  
presso-globosa.

— *β. sinuosa*: foliis profundius sinuosis:  
cupula turbinata; glande ovata.

*HAB. α.* ad lacum *Champlain*, in Pennsylvania et altis  
montibus utriusque Carolinæ et Georgiæ.

— *β.* in Carolina meridionali et Georgia.

MONOECIA. POLYANDRIA. 199

TRILOBA. Q. foliis petiolatis, oblonge cuneatis, summitate lobato-tricuspidibus; subtus eximie tomentosis; cupula scutellata; glande globosa.

HAB. a Nova Anglia ad Georgiam.

BANISTERI. Q. foliis longe petiolatis, acutangulo-quinquelobis, margine integris; subtus cinereo - tomentosis : cupula subturbinata; glande subglobosa.

HAB. in New-Yorck, New-Jersey, &c.

FALCATA. Q. foliis longe petiolatis, basi obtusis, divaricatim subpalmato - lobatis, lobis subfalcatis : cupula crateriformi; glande globosa.

HAB. a Virginia ad Floridam.

CATESBÆI. Q. foliis brevissime petiolatis, basi angustatis, acutis, subpalmato-lobatis, lobis interdum subfalcatis : cupula majuscula, squamis marginalibus introflexis; glande subglobosa.

HAB. in Marilandia, Virginia et Carolinis.

COCCINEA. Q. foliis longissime petiolatis, 5-7-lobis; lobis dentibusque acutissime angustatis : cupula turbinata, insigniter squamosa; glande brevi-ovata.

HAB. in Virginia et in utraque Carolina.

200 MONOECIA. POLYANDRIA.

PALUSTRIS. Q. foliis longe petiolatis, profundius  
DU ROI. septemlobis; sinubus latis, lobis oblon-  
gis, acute subdivisis: fructu parvo; cu-  
pula scutellata, lævi; glande subglo-  
bosa.

HAB. a Nova Anglia ad Virginiam et in occidenta-  
libus montium *Alléghanis*.

RUBRA. Q. foliis longe petiolatis, glabris; 7-9-  
L. lobis; lobis brevibus, dentibus angulisve  
acutissimis, sinubus subacutis: fructu  
majusculo; cupula scutellata, sublævi;  
glande turgide ovata.

HAB. a Canada ad Georgiam et in occidentalibus mon-  
tium *Alléghanis*.

CORYLUS. L.

MASC. *Cal.* 1-phyllus, 3-fidus, squamiformis,  
1-florus. *Cor.* o. *Stam.* 8.

FOEM. *Cal.* 2-phyllus, lacerus. *Cor.* o. *Styli* 2.  
*Nux* ovata.

AMERICANA. C. foliis lato-cordatis : involucri fructifero hispido, pilis apice glandulosis ; limbo ampliato, inæqualiter multifido : nuce subglobosa, basi latius derasa.

*HAB.* a Canada ad Floridam, etiam in maritimis.

ROSTRATA. C. pumila : foliis cordato-ovalibus : amenti masculi squamis ciliatis : involucri fructifero densissime hirsuto, in longum angustioremq̃ue tubum desinente.

*HAB.* in Canada et excelsis montibus *Alleghanis*.

CARPINUS. L.

MASC. *Cal.* 1-phyllus : squama ciliata. *Cor.* o.  
*Stam.* 20.

FOEM. *Cal.* 1-phyllus : squama ciliata. *Cor.* o.  
*Ovaria* 2. *Styli* singulis 2. *Nux* ovata.

AMERICANA. C. foliis oblongo-ovalibus : involucrorum laciniis acute dentatis.

*Obs.* Habitus *C. Betuli* : involucris fructiferis multo minoribus et conspicue dentatis.

*HAB.* a Canada ad Floridam.

202 MONOECIA. POLYANDRIA.

OSTRYA : americana. C. foliis cordato - ovalibus :  
amentis foemineis oblongioribus : invo-  
lucris fructiferis compresso-vesicariis.

*Obs.* Vix differt ab *Ostrya Micheliana* ; nec nisi  
amentis foemineis paulo longioribus.

*HAB.* a Canada ad Carolinam montosam.

LIQUIDAMBAR. L.

MASC. *Cal.* comm. 4-phyllus. *Cor.* o. *Filam.* nu-  
merosa.

FOEM. *Cal.* in globum , 4-phylli. *Cor.* o. *Styli* 2.  
*Caps.* multæ , in globum , bivalves , poly-  
spermæ.

STYRACIFLUA. L. foliis palmatis ; lobis acuminatis :  
L. axillis nervorum barbatis.

*HAB.* a Pensylvania ad Floridam.

COMPTONIA. GÆRT.

MASC. Amentum cylindricum, undique laxe imbricatum squamis 1-floris. *Cal.* diphyllus. *Cor.* nulla. *Filam.* tria, bifurca.

FOEM. Amentum ovatum. *Cal.* hexaphyllum. *Cor.* o. *Ovarium* subrotundum *Styli* duo, capillares. Nux unilocularis, evalvis.

ASPLENIFOLIA. C. foliis longe linearibus, alternatim crenato-pinnatifidis.

HAB. 4 Nova Anglia ad Virginiam et in excelsis montibus Carolinæ.

MONADELPHIA.

LARIX. TOURN.

Folia plura, fasciculata. Strobili per caulem sparsi seu axillares. Cætera fere ut in ABETE.

AMERICANA. L. foliis brevioribus: strobilis parvis, ovoideo-subglobosis; squamis paucioribus.

Obs. *Epinette rouge* Canadensium.

HAB. 2 sinu Hudson ad Pensylvaniam.

P I N U S. TOURN. JUSS.

*Squamae* foemineae interiores oblongae, clavatae, apice incrassato, inaequali, anguloso et extus umbilicato. Amenta in racemum terminalem compacta. Folia 2-5 ex eadem vagina simul prodeuntia. Caetera ut in *ABETE*.

INOPS. P. foliis binis, crassiusculis : amentis  
*AIT.* masculis ovato-oblongis, rufidulis : strobilis lateralibus, solitariis, conoideo-ovatis ; mucronibus tessularum compresso-conoideis, rigidis, deorsum subinclinatis.

*OBS.* Folia subsesquiuncialia. Ramuli obscure et squallide fusi.

*HAB.* in Nova Caesarea et Pennsylvania.

MITIS. P. foliis binis, tenuiusculis : strobilis  
 solitarie lateralibus, oblongiuscule conoideo-ovatis ; tessulis muticis.

*OBS.* Folia subsesquitriuncialia, interdum in nonnullis ramulis terna. Amenta mascula parva, ovata, divergentia, capitata.

*HAB.* in Virginia, Kentucky et Carolina.

PALUSTRIS. P. foliis ternis, longissimis : amentis  
*MILL.* masculis longo-cylindricis, fusco-glaucis, divergentibus : strobilis longissime conoi-

**MONOECIA. MONADELPHIA. 205**

deis ; tessularum tuberculo tumido , mucrone minutissimo terminato.

**OBS.** Squamæ gemmaceæ confertissimæ , flavidulo-canescentes.

**HAB.** a Carolina septentrionali ad Floridam , præsertim in maritimis.

**TÆDA.** P. foliis ternis , prælongis : amentis  
**L.** masculis oblongo-cylindricis , patulis ,  
divergentibus : strobilis floridis flavescen-  
tibus , sursum imbricatis , fructiferis  
oblongo-conoideis , tessulis mucrone sur-  
sum rugide-uncinato.

**HAB.** in Virginia , Carolina et Georgia.

**SEROTINA.** P. foliis ternis , prælongis ; amentis  
masculis in ovoideum congestis , erecto-  
incumbentibus , oblongo-ovatis , luteis :  
strobilis floridis squamis patulis , sub-  
squarrosis , glauco-subviolaceis ; fructi-  
feris subgloboso-ovatis ; tessularum mu-  
crone minutissimo.

**OBS.** Strobili anno tantum sequenti dehiscunt.

**HAB.** in humidis Carolinæ et Pensylvaniæ cupres-  
setis.

**STROBUS.** P. foliis quinis , gracilibus , glaucis :  
**L.** amentis masculis minimis , rufis : stro-



CUPRESSUS. L.

MASC. *Cal.* Amenti squama. *Cor.* o. *Antherae* 4,  
sessiles, absque filamentis.

FOEM. *Cal.* strobilus squamis multifloris. *Cor.* o.  
*Styli*? puncta concava. *Nux* angulata.

THUYOIDES. C. foliis squamulatim imbricatis;  
L. frondescencia compressa : strobilis mi-  
nutis, globulosis.

HAB. a Nova Cæsarea ad Carolinam septentrionalem,  
in paludosis.

DISTICHA. C. foliis planis, quasi pinnatim disti-  
L. chis : floribus masculis aphylo - race-  
mosis : strobilis subgloboso-ovoideis.

HAB. a Marylandia ad Floridam et a Tennassée fluvio  
ad Ludoviciam.

THUJA. L.

MASC. *Cal.* Amenti squama. *Cor.* o. *Stam.* 4.

FOEM. *Cal.* strobili ; squama 2-flora. *Cor.* o.

*Pist.* 1. *Nux* 1, cincta ala emarginata.

OCCIDENTALIS. T. foliationis articulis suborbiculatis : strobilis ovatis , teretibus ; squamis oblongo - ovalibus , obtusis : seminibus alatis.

*HAB.* in Canada et in montibus Virginiae.

PHYLLANTHUS. L.

MASC. *Cal.* 6-partitus , campanulatus. *Cor.* o.

FOEM. *Cal.* 6-partitus. *Cor.* o. *Styli* 3 , bifidi.

*Caps.* 3-locularis. *Sem.* solitaria.

CAROLINENSIS. P. herbaceus, erectus : foliis oblongis : limbo extrorsum appendicibus petaloidis plerumque quinque ; introrsum lacinulis totidem alternis et denticulatis (quasi bractearum loco.)

*HAB.* in Carolina et Georgia.

EUPHORBIA. L.

*Involuc.* commune calyciforme, subcampanulatum : limbo extrorsum appendicibus petaloidis plerumque quinque ; introrsum lacinulis totidem alternis et denticulatis (quasi bractearum loco.)

FLORES MASCULI tot quot lacinulae , iis oppositi.

210 MONOECIA. MONADBLPHIA.

CAL. subpolyphyllus : foliolis quasi pennatim laciniatis.

STAM. 4-5, aut plura.

FLOS FOEM. centralis, unicus : quandoque deest.

Cætera sat nota.

\* *Folius alternis.*

COROLLATA. E. perennis : caulibus erectis, simpliciusculis : foliis sparsis, obtusis, margine integro revolutis, ovalibus oblongis linearibusve ; umbella terminali, dichotoma, subaphylla : calycinis appendicibus petaloideis, tubo duplo longioribus, obovalibus, albis,

HAB. ubique in America septentrionali.

GRAMINIFOLIA. E. annua, pusilla, erecta, ab imo ramosa, minutissime puberula : foliis sparsis, linearibus, integerrimis ; supremis basi discoloribus : floribus fasciculato-terminalibus.

Obs. Affinis E. *hissopifoliae*.

HAB. in maritimis Floridae et Georgiae.

HETEROPHYLLA. E. frutescens, inermis : foliis petiolatis, subovalibus, subdentatis, diver-

**MONOECIA. MONADELPHIA.** 211

siformibus ; summis basi discoloribus :  
floribus fasciculato-terminalibus.

*HAB.* in Florida.

**\*\* Foliis oppositis.**

**DENTATA.** *E.* annua ; pumila , hirsuta : foliis oppositis , ovalibus , dentatis : floribus ad summitates congestis.

*Obs.* Folia superiora maculosa.

*HAB.* in Tennessee , juxta Nashville.

**POLYGONIFOLIA.** *E.* perennis , glaberrima , *sup.*  
*L.* perne dichotoma : foliis oppositis , subsessilibus , linearibus , integris : pedunculis axillaribus , solitariis , 1-floris.

*Obs.* Interdum procumbens , sæpe erecta. Folia sparsa ,  
*rara.*

*HAB.* in Carolina.

**HYPERICIFOLIA.** *E.* glabra , dichotome ramosissima ,  
*L.* erectiusculo - patula , ramis divaricatis : foliis oppositis , subfalcato-oblongis , argute serratis : ramusculis in summitate fasciculatim multifloris.

*HAB.* in arenis ripariis amnium fluviorumque occidentalis Americae septentrionalis.

**MACULATA.** *E.* dichotoma , erecto-patula : foliis  
*L.* oppositis ; oblongo-ovalibus , serratis ,

212 MONOECIA. *MONADELPHIA*.

pilosus : floribus subsolitariis : appendicibus calycinis albis.

*HAB.* in Florida.

**IPECACUANHÆ.** *E. perennis*, procumbens, *pub.*  
*L.* mila, glabra : foliis oppositis, sessilibus, obovalibus oblongisve, integris : pedunculis solitariis, 1-floris, elongatis.

*Obs.* Folia interdum linearia. Quandoque plantæ arena obrutæ. Cauliculi fasciculati quasi aphylli.

*HAB.* in sabulosis, a Pennsylvania ad Carolinam.

**MERCURIALINA.** *E. perennis* : caulibus gracilibilibus, simpliciter trifidis : foliis oppositis ternisve, subsessilibus, ovalibus, integris : pedunculis terminalibus, solitariis, 1-floris.

*Obs.* Habitus quodammodo *MERCURIALIS annuæ*, sed non ita ramosa.

*HAB.* in collibus, circa *Knoxville*.

**PUBENTISSIMA.** *E. perennis*, erecta; caulibus subdichotomis calycibusque pubentibus : foliis oppositis, sessilibus, subcordato-ovalibus, obtusis; pedunculis solitariis, capillaribus, unifloris : appendicibus calycinis albis, petaloideis.

*HAB.* in Carolina.

**THYMIFOLIA.** *E. humifusa*, dichotoma, gracilis, tota dense puberula : foliis parvulis, op-

**MONOECIA MONADELPHIA. 213**

positis, ovali-oblongis, obtusis, superne  
subserratis : fasciculis axillaribus, fo-  
liolis multifloris.

*HAB.* ad ripas fluminum *Mississipi et Ohio.*

**STILLINGIA. L.**

*MASC. Cal.* hemisphæricus, multiflorus. *Cor.* tu-  
bulosa, erosa.

*FOEM. Cal.* 1-florus, inferus. *Cor.* supera. *Stylus*  
3-fidus. *Caps.* 3-cocca.

**SYLVATICA.** *S.* radice crassissima : caulibus herba-  
ceis : foliis sessilibus, ovalibus, serru-  
latis : flosculis masculis squamam flora-  
lem vix superantibus.

*OBS.* Variat foliis sublanceolato-oblongis.

*HAB.* a Carolina ad Floridam.

**BIGUSTRINA.** *S.* fruticosa : foliis petiolatis, ovali-  
lanceolatis, utrinque acutis, integris :  
flosculis masculis brevissime pedicellatis.

*OBS.* Basis elapsæ capsulæ persistens, quasi triptera,  
truncata.

*HAB.* in sylvarum umbrosis, ad ripas ænium Caro-  
linæ et Georgiæ.

**SEBIFERA.** *S.* arborescens : foliis longe petiolatis,  
subrhombéo-ovalibus, acuminatis; infra

214 MONOECIA. MONADELPHIA.

basim glandula petiolari : flosculis masculis pedicellatis.

*CROTON sebiferum*. LINN.

*HAB.* in urbibus *Charlstown* et *Savannah* culta, et nunc fere spontanea in maritimis Carolinae.

CROTON. L.

MASC. *Cal.* cylindricus, 5-dentatus. *Cor.* 5-petala. *Stam.* 10-15.

FOEM. *Cal.* polyphyllus. *Cor.* α. *Styli* 3, bifidi. *Caps.* 3-locul. *Sem.* 1.

GLANDULOSUM. C. annuum, erectum, hispidissimum : foliis ovalibus, basi biglandulosis : fructibus quasi glomeratim sessilibus.

*C. multiglandulosum*. JACQ. *ic. rar.*

*HAB.* in Carolina.

CAPITATUM. C. herbaceum, erectum : foliis oblongo-ovalibus, obtusis, basi rotundatis, integris, utrinque tomentosis : floribus foemineis ad basim spicillæ marium capitata congestis.

Affine *C. argenteo*.

*OBS.* Petioli, ramuli calyces capsulaeque tomentoso-lanata.

*HAB.* in regione Illinoensi.

DISJUNCTIFLORUM. C. fruticosum : foliis breviter

**MONOECIA. MONADELPHIA. 213**

subcordato-ovalibus, obtusis, integris :  
floribus foemineis subsolitariis.

*C. maritimum?* WALT.

*Obs.* Flores foeminei a spicillis marium sessilium in  
sede diversa sejuncti.

*HAB.* in maritimis Carolinæ.

**MONANTHOGYNUM.** *C. herbaceum*, humile, erectum : foliis subcordato-ovalibus, obtusis, integris, tomentosis : spicilla dichotomali paucissimiflora; flore foemineo unico, pedicellato, cernuo.

*HAB.* in Tennassée, juxta Nashville.

**ARGYRANTHEMUM.** *C. caule fruticuloso* : foliis integerrimis, obtusis, obovalibus : racemo terminali, brevi, congestim multifloro; calycibus pedicellatis, nitidule argenteis.

*HAB.* in sylvis aridis Georgiae et Floridae.

**A C A L Y P H A. L.**

MASC. *Cal.* 3-4-phyllus. *Cor.* o. *Stam.* 8-16.

FOEM. *Cal.* 3 - phyllus. *Cor.* o. *Styli* 3. *Caps.* 3-cocca, 3-locularis. *Sem.* 1.

**VIRGINICA.** *A. annua*, puberula : foliis breviter  
L. petiolatis, sublanceolato-oblongis, serratis : involucris subsessiliter axillaribus,



216 MONOECIA. *MONADELPHIA.*

nervoso-plicatis, acutissime quasi cristato-incisis.

*HAB.* a Canada ad Virginiam.

CAROLINIANA. *A.* annua, puberula : foliis longe

*WALT.* petiolatis, subrhomboideo-ovalibus, serratis : involucris axillaribus, ampliatis, subcampanulatis, incisis.

*OBS.* Involucris sæpe ramulis aphyllis incidentes.

*HAB.* a Virginia ad Floridam.

J A T R O P H A. *L.*

MASC. *Cal.* o. *Cor.* 1-petala, infundibuliformis.

*Stam.* 10 : alterna breviora.

FOEM. *Cal.* o. *Cor.* 5-petala, patens. *Styli* 3 ;

bifidi. *Caps.* 3-locularis. *Sem.* 1.

STIMULOSA. *J.* herbacea, stimulis patulis horridissima : foliis subpalmato-lobatis ; lobis obtusiusculis, subsinuato-incisis : cymis brevi-pedunculatis : floribus candidis.

*PLUK.* t. 220. f. 3.

*MARCG. bras.* 79. fig. 2.

*J. urens.* *WALT.*

*OBS.* Radix filipendulo-tuberosa ; tuberibus subcylindraceis, edulibus.

*HAB.* a Virginia ad Floridam.

SYNGENESIA.

SICYOS. L.

MASC. *Cal.* 5-dentatus. *Cor.* 5-partita. *Filamenta* 3.

FOEM. *Cal.* 5-dentatus. *Cor.* 5-partita. *Stylus* 3-fidus. *Drupa* 1-sperma.

ANGULATA. S. ramis, petiolis pedunculisque lanosis : foliis sublobato-angulosis : fructibus capitatis, hispidis.

HAB. a Canada ad Tennassée, et in Carolina.

LOBATA. S. glabra : foliis semi5-fido-lobatis ; lobis acutissimis : fructibus solitariis, hispidis.

OBS. Fructus dupla magnitudine præcedentis.

HAB. in occidentalibus Pensylvaniæ, juxta fluvium Ohio.

MELOTHRIA. L.

*Cal.* 5-fidus. *Cor.* campanulata, 1-petala. *Bacca* 3-locularis, polysperma.

PENDULA. M. foliis subreniformibus, lobato-angulosis ; lobo impari cæteris æquali : bacca cylindræco-ovata.

OBS. Plures, ni fallor, hujus generis species permixtæ sunt sub eodem nomine !

HAB. in Carolina et Georgia.

## SCHISANDRA.

*Σχισίς, Anp* : *fissurae* antheris interjectæ.

Frutices scandentes : folia alterna : flores axillares.

## M A S C.

CAL. subtriplici serie 9-phyllus, deciduus : foliolis subrotundis, concavis ; interioribus gradatim minoribus, intensius coloratis.

COR. o.

STAM. antheræ 5, subsessiles ; basi contiguous, apicibus supra loculos connatæ, lateribus totidem interjectis rimis discretæ.

## F O E M.

CAL. maris.

PIST. ovaria plurima, in receptaculo demum protractili subcapitatim congesta, ovoidea, stigmate brevi acuminata.

FRUCT. bacca receptaculo elongato quasi spicatim affixæ, inæqualiter ovoideæ, 1-spermæ.

SEM. ovato-oblongum, teretiusculum, læviusculum.





P. J. Redouté Del.

Plac. Sc.

**SCHISANDRA** *coccinea* L.

MONOECIA. SYNGENESIA. 219

Embryo erectus; albumine carnoso et vi-  
rente inclusus : radícula oblonga, cylin-  
drica; cotyledonibus ovalibus, plano-  
appressis.

*Obs.* Genus MENISPERMO-affine.

COCCINEA. C. glabra : foliis lanceolato-ovalibus,  
utrinque acutis, rariter subdentatis : flo-  
ribus coccineis.

*HAB.* in umbrosis Carolinæ et Georgiæ.

## CLASSIS XXII.

## DIOECIA.

## MONANDRIA.

## NAJAS. L.

MASC. *Cal.* cylindricus, 2-fidus. *Cor.* 4-fida. *Filamentum* nullum.

FOEM. *Cal.* o. *Cor.* o. *Pist.* 1. *Caps.* ovata, 1-locularis.

CANADENSIS. *N.* pusilla, filiformis, lævis : foliis angustissime linearibus.

HAB. in Canadæ lacu *S. Joannis*.

## DIANDRIA.

## VALLISNERIA. L.

MASC. *Spatha* 2-partita. *Spadix* tectus flosculis. *Cor.* 3-partitæ.

FOEM. *Spatha* 2-fida, 1-flora. *Cal.* 3-partitus, superus. *Cor.* 3-petala. *Gly.* 3. *Caps.* 1-locul. polysperma.

AMERICANA. *V.* foliis erectis : pedunculis non spiralibus.

OBS. Folia minus quam in *spiral*i elongata, stantia, inferne non angustata.

HAB. in flumine *Mississipi* et in fluvio *S. Joannis* Floridæ.

## CERATIOLA.

Frutices habitu et florescentia Ericoidei. Gemmulæ minutæ, sessiliter axillares, unifloræ, ovoideæ, imbricatæ : squamis 8, oppositis, appressis; extimis gradatim minoribus; intimis sexum obvolventibus : calyce corollaque nullis.

## M A S C.

STAM. duo, summitate gemmulæ hiantē exerta : filamenta erecta, plana : antheræ tantisper supra medium dorsum appensæ, majusculæ, inferne semibifidæ, superne emarginatæ, scaberulæ; loculis oblongo-ovatis, apice obtusis, sulco longitudinali dehiscentibus.

## F O E M.

PIST. ovarium gemmula apice angustato clausa involutum, obovato-subglobosum : stylus subsemiexertus : crassiusculus, erectus : stigma inæqualiter multipartitum; laciniis subfiliformi-ligulatis, sæpe geminatis et inde quasi bicornibus, radiatim recurvo-patentissimis.

BACCA minima, squamulis gemmulaceis



BIOECIA. *DIANDRIA*.

(omissa summitate) obvoluta, spheroidæ : caro paucissima : nuculæ duæ, subhemisphæricæ ; testa crassiuscula, lignosa ; uniloculares, 1-spermæ.

SEMEN compresso - obovatum, hinc convexius.

*Obs.* Genus *EMPETRO* valde affine.

**ERICOIDES.** C. ramis minutissime lanuginosis : foliis quaterno-verticillatis, angusto-linearibus seu acerosis : gemmularum flandarum squamis lanoso-fimbriatis.

*HAB.* in aridis sabulosis Georgiæ et Floridæ.

O L E A. *L.*

*Cor.* 4-fida, laciniis subovatis. *Drupa* monosperma.

**AMERICANA.** foliis late oblanceolatis, coriaceis, lucidis, integerrimis : drupa globosa.

*Obs.* Arbor : flores albidii, composite racemosi : rudimenta staminum in foemina.

*HAB.* in maritimis, a Carolina ad Floridam.

## ADELIA. BROWN.

Caulis fruticosus. Folia opposita. Flores minuti, axillares, glomerato-fasciculati; fasciculis opposite squamoso-bracteatis, plerumque simplicibus, rarius ramulosis; squamis unifloris.

## M A S C.

CAL. minutissimus, 4-phyllus: foliolis erectis, sublanceolatis.

COR. nulla.

STAM. 2-3: filamenta intra calycem summo pedicello contigue affixa, longiuscule filiformia: antheræ subcordato-ovatae, obtusae.

## F O E M.

CAL. 4-phyllus: foliolis 2-oppositis majoribus, oblongis, erecto-patentiusculis; 2 minutissimis, quandoque deficientibus.

COR. nulla.

PIST. ovarium biloculare; loculis biovulatis; summo pedicello immediate insidens, subrotundo-ovoideum: stylus crassiusculus, cylindraceus: stigma depresso-capitatum, levissimo sulco subbilobum.

**BACCA** ovoidea; pariete interna (nucis tenuis instar) cartilaginea et quasi plicato-sulcata; abortatione fere constanter monosperma.

**SEM.** oblongum, utrinque attenuatum, longitudinaliter sulcatum nervosumque; ad summitatem dissepimenti ad alterum cavitatis latus propulsi adnexum et ideo inversum.

**INTEG. PROPRIUM** membranaceum, rufescens.

**EMBR.** inclusus in materie subcorneo-carnosa, crassa; uti semen inversus, rectus. Radicula cylindrica, longiuscula. Cotyledones sublanceolato-oblongæ, planoligulatæ.

**OBS.** Genus affine **CHIONANTHO**.

**PORULOSA.** A. foliis coriaceis, sessilibus, ovalilanceolatis, obtusis, margine revolutointegris, subtus porulis pertusis.

**OBS.** Folia subtus quasi rubiginosa.


**HAB.** in maritimis Floridæ.

**LIGUSTRINA.** A. foliis submembranaceis, subpetiolatis, lanceolato-oblongis, integerrimis: drupa breviuscule ovata.

**OBS.** Habitus et folia **LIGUSTRI**.

**HAB.** in fruticetis Illinoensibus, Tennassée, &c.





P. J. Redoute Del.

Plac. Sc.

**ADELIA** *acuminata.*

DIOECIA. DIANDRIA. 225

ACUMINATA. A. foliis membranaceis, quasi rhomboideo - lanceolatis, superne longius  
*tab. 48.* angustatis, levissime serratis : drupa oblonga, ante maturitatem superne rostratim attenuata.

*HAB.* ad ripas fluviorum Carolinæ et Georgiæ.

S A L I X. L.

MASC. *Amenti* squamæ. Cor. o. *Glandula* baseos nectarifera.

FOEM. *Amenti* squamæ. Cor. o. *Stylus* 2-fidus. *Caps.* 1-locularis ; 2-valvis. *Sem.* papposa.

ERIOCEPHALA. S. diandra : ramulis minutim tomentosis : foliis oblongo-ovalibus, basi subretusis, serrulatis : amentis ovalibus, confertim villosissimis.

*Obs.* Affinis *S. capreae*.

*HAB.* in regione Illinoensi.

CORDATA. S. ramulis foliisque villosis : foliis cordato-ovalibus, acuminatis, argute serrulatis ; stipulis foliaceis, maximis.

*HAB.* in Canada, ad lacum *S. Joannis*.

INCANA. S. foliis lanceolato-oblongis, integerrimis, utrinque tomentosis, subtus candidioribus ; petiolis brevibus, basi auriculatis.

*HAB.* in Canada, juxta lacum *S. Joannis*.

**LONGIROSTRIS.** *S. pumila*, diandra : foliis subsessilibus, longiuscule lanceolatis, subcuneatis, integriusculis, utrinque tenuiter tomentosis : amentis oblongo-ovatis, cinereis : capsulis divergentibus, longissime rostratis.

*S. alpina* ? WALT.

*OBS.* Affinis *S. arenariae*.

*HAB.* a Nova Anglia ad Carolinam; copiose in dumetosis apricis Tennassée.

**CAROLINIANA.** *S. foliis lanceolatis*, subtiliter argutisque serrulatis, subsessilibus : staminibus 4-6 : amenti foeminei squamis oblongis, minutissime partimque lanuginosis; ovariis oblongis, glabris.

*S. pentandra* ? WALT.

*OBS.* Affinis *S. triandrae*.

*HAB.* in Carolina et Georgia.

T E T R A N D R I A.

H I P P Ô P H A E. L.

MASC. *Cal.* 2-partitus. *Cor.* o.

FOEM. *Cal.* 2-fidus. *Cor.* o. *Styl.* 1. *Bacca* monosperma.

CANADENSIS. H. foliis petiolatis, ovalibus : spicis  
L. masculis distinctifloris.

HAB. in Canada et Noveboraco, juxta lacum Champlain.

M Y R I C A. L.

MASC. *Amenti* squama lunata. *Cor.* o.

FOEM. *Amenti* squama lunata. *Cor.* o. *Styli* 2.  
*Bacca* 1-sperma.

GALE. M. foliis cuneato-lanceolatis, superne  
L. rariter leviterque serratis : amentis masculis imbricatis, squamis ciliolatis : fructibus squamoso-capitatis.

HAB. in Canada, a sinu Hudsonis ad Pensylvaniam.

CERIFERA. M. foliis cuneato-lanceolatis, superne  
L. rariter serratis : amentis masculis



DIOECIA. *TETRANDRIA*.

laxis : fructibus sphaericis , nudatis , distinctis.

Var. *α. arborescens* : foliis lanceolatis , utrinque angustatis , acutissimis : fructibus minoribus.

— *β. media* : foliis latioribus : fructibus majusculis.

— *γ. pumila* : fruticulosa : foliis minoribus , magis cuneatis.

*HAB.* *α.* a Carolina ad Floridam , in umbrosis.

— *β.* a Nova Anglia ad Floridam , in udibus et juxta rivulos.

— *γ.* in aridis , a Carolina ad Floridam.

I L E X. *L.*

*Cal.* 4-dentatus. *Cor.* rotata. *Stylus* o. *Bacca* 4-sperma.

*OPACA.* I. foliis ovalibus , rigide patuleque dentato-spinosis : fasciculis florum laxis ; pedunculis compositis : calycibus acutiusculis , glabellis : fructibus ovoideis.

*AIT.*

*HAB.* in umbrosis humidis , a Pennsylvania ad Floridam.

*DAHOON.* I. ramulis calycibusque pubescentibus : *WALT.* foliis laurinis , sub lanceolato-oblongis , rariter serratis : fasciculis florum pedunculatis.

*I. Cassine.* LINN. Ait.

*HAB.* in paludosis , a Carolina ad Floridam.





P. J. Redouté Del.

Plée Sc

**ILEX** *canadensis*.

DIOECIA. *PETRANDRIA*. 229

MYRTIFOLIA. I. foliis pusillis, lineari-lanceolatis,  
*WALT.* subintegris, mucronatis, utrinque gla-  
berrimis : floribus fertilibus solitariis ;  
calycibus acutis, glabris.

*HAB.* in stagnosis Carolinæ et Georgiæ.

CASSENSA. I. foliis parvulis, oblongiuscule ova-  
libus, utrinque obtusis, crenato - ser-  
ratis.

*I. Cassine. WALT.*

*CASSENSA vera, CATESB.*

*HAB.* in maritimis, a Carolina ad Floridam. 5.

PRINOIDES. I. foliis deciduis, ovali-lanceolatis,  
*AIT.* utrinque angustatis, leviter serratis; pe-  
dunculis simplicibus, longiusculis, 1-flo-  
ris; fertilibus solitariis.

*I. decidua. WALT.*

*HAB.* in umbrosis, ad ripas rivulorum Carolinæ et  
Georgiæ.

CANADENSIS. I? foliis ovalibus, integris seu rariter  
serratis: pedunculis solitariis, longis, uni-  
*tab. 49.* floris: fructu obtuse subtetragono.

*HAB.* in præruptis montibus, a sinu *Hudsonis* ad  
lacum *Champlain*.

*P E N T A N D R I A.*

HUMULUS. L.

MASC. *Cal.* 5-phyllus. *Cor.* o.

POEM. *Cal.* 1-phyllus, oblique patens, integer.

*Cor.* o. *Styli* 2. *Sem.* 1. intra calycem foliatum.

LUPULUS. HUMULUS.

L.

VITIS. L.

*Petala* apice cohærentia, emarcida. *Bacca* 5-sperma, supera.

Obs. Species mæx omnes dioicæ? *V. vulpina* LINNÆI adeo est obscura, ut nullam hoc nomine appellarim.

LABRUSCA. V. foliis amplissimis, lato-cordatis,  
L. sublobato-angulatis, subtus incano-tomentosis : racemis fertilibus parvulis :  
baecis majoribus.

*V. taurina.* WALT.

Obs. Omnibus incolis hæc *vulpina* dicta. (*Fox grape*) uti recte docet phrasis CLAYTONI.

Antillana videtur speciei diversa.

HAB. a Pennsylvania ad Floridam.

ESTIVALIS. V. foliis lato-cordatis, 3-5-lobatis,  
subtus rariuscule tomentosis, pube rufa,

**DIOECIA. PENTANDRIA. 231**

sinubus rotundato-obtusis : racemis fertilibus oblongis : baccis parvulis.

*V. Labrusca.* WALT.

*Obs.* Vulgo *summer grape* seu æstivalis nominatur.

*HAB.* in sylvis, a Virginia ad Carolinam.

**CORDIFOLIA.** V. foliis cordatis, acuminatis, subæqualiter dentatis, utrinque glabris : racemis laxè multifloris, polycarpiis : baccis parvulis, serotinis.

*Obs.* *Winter grape* sive uva hiemalis incolis dicta.

*HAB.* a Pennsylvania ad Floridam.

**RIPARIA.** V. foliis inæqualiter inciso-dentatis, breviuscule trifidis : petiolo, nervis marginæque pubescentibus.

*Obs.* Incolis gallis *Vigne des battures* dicitur.

*HAB.* ad ripas et in insulis fluviorum *Ohio*, *Mississipi*, &c.

**ROTUNDIFOLIA.** V. foliis utrinque lucidis, reniformi-cordatis, subæqualiter dentatis : racemorum floribus pluries capitulatis : baccis magnis.

*Obs.* *Muscadin grape* vulgo audit.

*HAB.* a Virginia ad Floridam.

**PYRULARIA.**

Caulis fruticosus : folia alterna, exstipulata : flores perpusilli, spicati ; sexibus dioicis.

M A S C.

**CAL.** breviter campanulatus, subsemi-5-fidus:

tubus subhemisphærice concavus : limbus laciniis revolutis.

COR. o.

STAM. 5, summo calycis tubo inserta, laciniis opposita brevioraque, erecta : filamenta brevia, crassiuscula : antheræ subdidymæ, loculis ovatis, emisso polline patentissimis.

DISCUS. Materies carnosa, tubum calycis intus incrassans, ad ambitum inter singulorum staminum bases desinit in totidem processus rotundato-obtusos.

#### F O E M.

CAL. turbinato-campanulatus, 5-fidus; laciniis revolutis.

STAM. 5, antheris incassis.

DISCUS epigynicus, processibus manifestioribus.

PIST. ovarium inferum : stylus limbo calycis brevior, erectus, crassus : stigma depresso-capitatum.

DRUPA pyriformis, limbo calycino latius coronata : nux subglobosa, tenuiuscula, 1-ocularis, 1-sperma.

SEMEN nuci conforme : fundo loculi affixum; tegumine membranaceo, quasi fibrilloso; nucleo oleaginoso.

OBS. Genus *Nyssoidium*.

DIOECIA. *PENTANDRIA.* 233

**PUBERA.** P. ramulis, foliis calycibusque pubentibus : foliis ovali-oblongis, acuminatis, integerrimis.

**OBS.** Frutex 3-6-pedalis; etiam ultra; radice ingrate olente.

**HAB.** in montibus Carolinæ occidentalis et in Virginia.

**A C N I D A. L.**

Habitus et florescentia **AMARANTHI.**

**M A S C.**

**CAL.** quasi campanulatus, 5-partitus, pelucido-membranaceus.

**COR.** o.

**STAM.** imo calyci inserta, hujus laciniis opposita, iisque paulo longiora : antheræ utrinque bipartitæ et inde quasi acute bicornes.

**F O E M.**

**CAL.** minor, 3-partitus; laciniis inæqualibus, lineari-subulatis, acutis, erecto-appressis.

**COR.** o.

**PIST.** ovarium calyce multoties longius, oblongum, inæqualiter 3-gonum : stylus nullus, nisi brevissimum ovarii acumen;



**DIOECIA. PENTANDRIA.**

stigmata 3, capillaria, patula, glanduloso-puberula.

**CAPS.** calyce minuto basi stipata, inæqualiter obovoideo-trigona; non dehiscens, 1-locularis, 1-sperma.

**SEM.** suborbiculato-obovatum, compressum, lævigatum; funiculo basilari annexum fundo pericarpium.

**OBS.** Interdum stigmata 4 seu 5, tuncque fructus 4-5-gonus.

Genus **AMARANTHO** conterminum.

**CANNABINA.** A. foliis lanceolatis : capsulis læviusculis, acutangulis.

**HAB.** ad ripas fluviorum inundatas Virginie.

**RUSOCARPA.** A. foliis ovali-lanceolatis : capsulis *tab. 5o.* obtusangulis, rugosis.

**OBS.** Herba maxima, erecta, altitudine humana seu ultra; caule crasso, fistuloso, anguloso.



P. J. Redouté Del.

Plée Sc.

*ACNIDA rusocarpa.*



## ZANTHOXYLUM. L.

MASC. *Cal.* 5-partitus. *Cor.* o.

FOEM. *Cal.* 5-partitus. *Cor.* o. *Pist.* 5. *Caps.* 5,  
1-spermæ.

RAMIFLORUM. *Z.* foliis pinnatis; foliolis ovali-  
lanceolatis, integerrimis, subsessilibus,  
utrinque pubentibus: floribus rameis:  
capsulis stipitatis.

*Z. caribaeum.* GÆRTN.

*Z. Clava Herculis*, β. LINN.

*Z. fraxinifolium.* MARSH.

*HAB.* a Canada ad Virginiam et in Kentucky.

TRICARPUM. *Z.* foliis glaberrimis, pinnatis; fo-  
liolis petiolatis, oblongo-ovalibus, sur-  
sum angustatis, falcato-obliquis: cap-  
sulis ternis, sessilibus.

*Z. fraxinifolium?* WALT.

*Obs.* Variat capsulis binis nec plus quam ternis.

*HAB.* a Carolina ad Floridam.

*H E X A N D R I A*.

## P R I N O S. L.

*Cal.* 6-fidus. *Cor.* 1-petala, rotata. *Bacca* 6-sperma.

GLABER. P. foliis cuneato-lanceolatis, superne rariter subdentatis.

*Obs.* An vere congener *P. verticillati*. LINNÆI?

*HAB.* a Nova Anglia ad Floridam.

GRONOVII. P. foliis ovalibus, serrulatis, acuminatis : fasciculis florum masc. axillariibus, umbelluliformibus; floribus foemineis aggregatis; utrisque 6-partitis.

*P. verticillatus*. LINN.

*HAB.* a Canada ad Virginiam.

AMBIGUUS. P. foliis ovalibus, utrinque acuminatis; pedicellis masc. 1-floris, ad imos ramunculos numerose congestis; foem. solitariis : florum partitione quaternaria.

*Obs.* Interdum florum partitio quinaria; tunc videtur

*CASSINE caroliniana*. WALTERI. Certo tamen *P. verticillati* LINN. congener.

*HAB.* in Carolina.

## SMILAX. L.

MASC. *Cal.* 6-phyllus. *Cor.* o.

FOEM. *Cal.* 6-phyllus. *Cor.* o. *Styli* 3. *Bacca*  
3-locularis. *Sem.* 2.

\* *Caule fruticoso.*

ROTUNDIFOLIA. *S.* ramis angulosis, rariter aculeatis : foliis subrotundatim cordato-ovalibus, acutis, glaberrimis, 5-nerviis : baccis sphæricis.

*OBS.* Affinis *S. deciduae*. LINN.

*HAB.* a Nova Anglia ad Virginiam.

GLAUCA. *S.* ramis subteretibus, passim aculeatis : foliis quasi cordato-ovalibus, acuminatis, glabris, subtus glaucis, 5-nerviis : umbellulis longiuscule pedunculatis, floribus parvulis.

*HAB.* in sylvis Carolinæ.

BONA NOX. *S.* ramis angulosis, aculeatis : foliis  
*L.* hastato-lanceolatis, auribus rotundato-obtusis, 5-nerviis : baccis globosis.

*HAB.* in maritimis Carolinæ et Floridæ.

LAURIFOLIA. *S.* ramis teretiusculis, subinermibus :  
*L.* foliis coriaceis, lanceolato-oblongis, trinerviis, lævigatis, persistentibus : um-

*OCTANDRIA*.

## POLYGONELLA.

Caulis fruticosus. Folia alterna, stipulis vaginantibus. Ramuli spiciflori; bracteolis 1-floris, minutis, vaginantibus, subimbricatis: floribus parvulis, breviter pedicellatis, dioicis.

CAL. petaloideus, patens, 5-partitus: laciniis subæqualibus, ovalibus.

## M A S C.

STAM. 8; (interdum 7) imo calyci inserta, ejusque circiter longitudine: filamenta subulata, patula: antheræ subrotundæ.

PISTILLI incrementum oblongo-triquetrum, flaccidum; stigmatulis tribus imperfectis terminatum.

## F O E M.

PIST. ovarium ovato-triquetrum, apice angustato terminatum stigmatulis tribus brevibus, subclavato-obtusis.

Nequidem vestigia staminum.

FRUCT. capsula oblongo-triquetra, non dehiscens, 1-sperma; vestita calyce laciniis tribus ampliatis.

Obs. Genus affine *ATRAPAXI*.







DIOECIA. DECANDRIA. 241

PARVIFOLIA. P. caulibus stricte erectis : foliis pusillis, cuneato-linearibus : floribus spicatis.

HAB. in aridissimis Carolinæ.

DECANDRIA.

GYMNOCLADUS. LAM.

Flores racemosi. *Calyx* tubulosus, 5-fidus. *Petala* 5, æqualia, tubo inserta. *Stamina* non exerta. *Legumen* magnum, crassum, intus pulposum.

CANADENSIS. G. foliis bipinnatis; foliolis ovalibus; *tab. 51.* acuminatis, puberulis.

POLYANDRIA.

MENISPERMUM. L.

MASC. *Petala* 4 exteriora; 8 interiora. *Stam.* 16.  
FOEM. *Cor.* maris. *Stam.* 8, sterilia. *Baccae* binæ monospermæ.

CANADENSE. M. foliis peltatis, subreniformibus, angulosis : racemis longe pedunculatis.

Obs. Flores masculi polyandri. Fructus maturi nigri.

HAB. a Canada ad Carolinam.

## DIOECIA. POLYANDRIA.

CAROLINIANUM. M. foliis vix peltatis, ovalibus seu

L. quasi hastato-cordatis : racemis subsessilibus.

Obs. Flores masculi hexandri. Fructus maturi rubri.

HAB. in Carolina, Georgia, Florida.

## ZAMIA. L.

MASC. *Amentum* strobiliforme, squamis subtus tectis polline.

FOEM. *Amentum* strobiliforme in utroque margine. *Drupa* solitaria.

PUMILA. Z. foliis glabriusculis ; foliolis subcuneatim lanceolatis, rotundato-obtusis.

HAB. in Florida.

POPULUS. L.

**MASC.** Amenti *Cal.* lamina lacera. *Cor.* turbinata, obliqua, integra.

**FOEM.** Amenti *Cal.* et *Cor.* maris. *Stigma* 4-fidum. *Caps.* 2-locularis. *Sem.* multa, papposa.

**OBS.** Glandulæ basis foliorum nihil valent ad distinguendas species ! quippe quæ, in eadem specie, aut desunt aut adsunt.

**ANGULOSA.** *P.* ramis acutangulis : foliis deltoideis, glaberrimis, serratis : floribus masculis polyandris ; foemineis remotiuscule distinctis, glabris : ovario subgloboso.

*P. angulata.* AIT.

**HAB.** a Pennsylvania ad Carolinam.

**GRANDIDENTATA.** *P.* petiolis superne compressis : foliis subrotundo-ovalibus, acuminatis, utrinque glabris, inæqualiter sinuato-grandidentatis.

**OBS.** Affinis *P. albae* ; foliis itidem quandoque basi biglandulosis.

**HAB.** in Canada.

**TREMULOIDES.** *P.* foliis parvulis, suborbiculatis, abrupte acuteque acuminatis, serrulatis, margine pubescentibus.

**HAB.** in Canada et Noveboraco.

244 DIOECIA. *MONADELPHIA*.

HETEROPHYLLA. P. ramulis teretibus; foliis am-

L. plis, sinu parvo cordatis, obtusis, leviter dentatis; junioribus tomentosis: floribus masculis polyandris; amenti foem. glaberrimi floribus remote distinctis, longius pedicellatis.

HAB. a Nova Anglia ad Carolinam et copiose in regione Illinoensium.

NIGRA. P. foliis deltoidibus, acuminatis, serratis.

L.

BALSAMIFERA. P. foliis ovatis, serratis, subtus

L. albidis, stipulis resinosis.

HAB. in America septentrionali.

*MONADELPHIA*.

IRESINE. L.

MASC. Cal. 2-phyllus. Cor. 5-petala. Nectaria 7.

FOEM. Cal. 2-phyllus. Cor. 5-petala. Stigmata 2, sessilia. Caps. seminibus tomentosis.

CELOSIODES. I. foliis primariis lato-ovalibus; superioribus rameisque ovali-lanceolatis.

L.

Obs. Paniculae mitis laxae quam in Antillana.

HAB. in sabulosis fluvii Ohio ripariis hieme inundatis.

TAXUS. L.

MASC. Cal. 3-phyllus gemmæ, Cor. o. Stam. multa. Antheræ peltatæ, 8-fidæ.

FOEM. Cal. 3-phyllus gemmæ, Cor. o. Stylus o. Semen 1, calyculo baccato, integerrimo.

BACCATA :  $\beta$ . minor T. foliis approximatis.

L.

Obs. Humilior, patulus : floribus fructibusque minoribus.

HAB. in umbrosis Canada.

JUNIPERUS. L.

MASC. Amenti calyx squamæ. Cor. o. Stam. 3.

FOEM. Cal. 3-partitus. Petala 3. Styli 3. Bacca 3-sperma, tribus tuberculis calycis inæqualis.

COMMUNIS. J. foliis ternis, non decurrentibus,

L. patentibus, mucronatis : fructu sphærico, nigro longioribus.

HAB. a sinu Hudsoni ad Noveboracum.

VIRGINIANA. J. foliis vetulis ternis, adnato-decurrentibus, subulato-spinescentibus; junioribus oppositis, modo acerosis, modo imbricatis, lanceolato-ovalibus, acutis.

Obs. Sunt individua foliis plerorumque ramorum et ramulorum acerosis laxisque; sunt etiam alia ramulis plerisque aut omnibus minutim imbricatis : unde facilis erroris ansa nonnullis subministrata est cultoribus; qui priora *J. virginianam*, posteriora *J. carolinianam* dixere.

HAB. a lacu Champlain ad Floridam.

246 DIOECIA. *MONADELPHIA*.

SABINA. J. ramulis confertis : foliis vetulis ad-  
L. nato-squamiformibus, mucrone libero  
breviter acutis; junioribus imbricatis,  
acutiusculis.

*Obs.* Caules patuli aut procumbentes.

*HAB.* in præruptis rupibus, a sinu *Hudson* ad ostium  
fluvii *Sagney*.

ARBADENSIS. J. foliis quadrifariam imbricatis,  
L. ovalibus, obtusiusculis : fructibus pu-  
sillis.

*HAB.* in Floridae maritimis et in *Bahama*.

---

POLYGAMIA. MONOECIA. 247

CLASSIS XXIII.

POLYGAMIA.

MONOECIA.

PLANERA. GMEL. *Syst.*

Frondescencia et floratio ULMI; fasciculorum squamis 1 - floris, inferioribus masculis, superioribus hermaphroditis aut rarius foemineis.

MASC.

CAL. membranaceus, subcampanulatus, 4-5-fidus.

STAM. 4-6, plerumque 4, exerta.

PISTILLI nequidem vestigium.

HERMAPHROD.

CAL. marium.

STAM. marium; antheris tardius dehiscen-  
tibus.

PIST. ovarium ovoideum, scabrum seu  
laeve: stigmata 2, oblonga, glandulosa,  
recurvo-divergentia.

CAPS. subglobosa, membranacea, 1-locu-  
laris, non dehiscens; aut laevis, aut un-



248      **POLYGAMIA. MONOECIA.**

dique cutaneis processibus quasi squamulosa.

**SEM.** unicum , ovoideum , acutum.

*Obs.* Genus **ULMO** valde affine.

**GMELINI.** P. foliis manifestè petiolatis , oblongo-ovalibus , sensim angustatis , acutis , basi obtusis , æqualiter serratis : capsula scabra.

*HAB.* ad ripas fluviorum *Ohio* , *Mississipi* et *Carolinæ*.  
Primo vere floret , eodem abeunte fructif. 5 .

**RICHARDI.** P. foliis subsessilibus , oblongo-ovalibus , subcrenatim grandi-dentatis , basi emarginatis : capsula lævi.

*Obs.* Jamdudum in hortis gallicis , ignota appellationis *Orme de Sibérie* origine , colitur. Prima vice in horto Trianonensi floruit Aprili 1779. In dissertatione de **ULMO** , ineunte anno 1781 , *Academiæ Scient. Parisinæ* proposita a cl. **L. C. RICHARD** , hæc species fuit descripta , et **ULMUS polygama** dicta.

*HAB.* circa mare *Caspium*.

CELTIS. L.

HERMAPHROD. *Cal.* 5-partitus. *Cor.* o. *Stam.* 5.

*Styli* 2. *Drupa* 1-sperma.

MASC. *Cal.* 6-partitus. *Cor.* o. *Stam.* 6.

OCCIDENTALIS. *C.* foliis ovalibus, acuminatis; petiolo partimque nervis pubentibus : floribus inferioribus ternis ; superioribus solitariis.

*Oss.* Pedunculi fructiferi longitudine petiolorum.

*HAB.* a Nova Anglia ad Carolinam : copiose ad *Mississipi*.

VERATRUM. L.

HERMAPHROD. *Cal.* o. *Cor.* 6-petala. *Stam.* 6.

*Pist.* 3. *Caps.* 3, polyspermæ.

MASC. *Cal.* o. *Cor.* 6-petala. *Stam.* 6. *Pist.* rudimentum.

Genus forsân in posterum cum MELANTHIO in unicum cogendum. Distinguitur foliis latioribus , quasi plantagineis : calycis laciniis non ita abrupte unguiculatis ; præsertimque glandularum defectu. Aliunde vix accersitur recta differentiæ ratio.

ALBUM. *V.* foliis lato-ovalibus, sulcatis : floribus subsessilibus , virescentibus : calycis laciniis oblongo-ovalibus , acutis.

*Oss.* Omnibus partibus pubens.

*HAB.* in rivulis montanis, a Carolina ad Cånadam.

250      **POLYGAMIA. MONOECIA.**

**PARVIFLORUM.** V. foliis ovalibus lanceolatisve, planis, glabris : paniculæ ramulis filiformibus ; floribus pedicellatis, viridibus : calycibus stellatis ; laciniis ovali-lanceolatis, utrinque acutis, ungue staminiferis.

*Obs.* Foliatio **VERATRI**, non ita sulcata : floratio **MELANTHII** ; calyce eglanduloso : planta utrumque genus connectens.

*Aff.* **PLUCK.** t. 434. f. 8.

*HAB.* in excelsis montibus Carolinæ. Julio floret.

**MELANTHIUM. L.**

Folia elongata, debilia : scapus foliosus : panicula subpyramidata ; ramis spicifloris, inferioribus rarius subdivisis : floribus pedicellatis, bracteolis totidem brevibus.

Ex floribus ejusdem paniculæ alii hermaphroditi fertiles ; alii, modo pistilli vitio, modo hujus defectu, steriles : interdum etiam omnes steriles ; nunquam vero omnibus conceditur fertilitas. Pubescentia quasi pulverulenta.

**CAL.** petaloideus, rotatus, profundissime 6-partitus : laciniis æqualibus, inferne unguiculatis ; unguiculis angustis, distanter a pistillo staminiferis ; laminis introrsum ad basim glandula gemina, colorata insignitis.

**STAM.** 6, calyce breviora, assurgentia : filamenta setacea, incurva : antheræ rotundato-reniformes.

**POLYGAMIA. MONOECIA. 251**

**PIST.** ovarium subovato-trigonum ; lateribus depressis ; angulis, in totidem stylos apice stigmatico obtusos, desinentibus.

In nonnullis floribus pistilli tantum rudimentum.

**FRUCT.** capsula calyce persistente multoties major, membranacea, subovata, apice subtrifida et persistentibus stylis quasi tricornis, compresso-tricocca; immediata coccorum ad axim connexione inseptatim trilocularis ; loculis oligospermis.

**SEMINA** in singulo loculo circiter 6 seu pauciora; planiuscule compressa; subovalia, ambitu alato-membranacea.

**VIRGINICUM.** M. elatius : panicula pyramidata :  
L. laciniis calycinis ovalibus.

*Obs.* Planta 2-3-pedalis : paniculæ fami omnes simplices, laxiflori : flores obsolete pallido-albidi, interdum intus obscure lividi.

Icon *PLUCKNETI* non quadrat.

*HAB.* in humidis herbosis, a Pennsylvania ad Carolinam.

**RACEMOSUM.** M. panicula racemoso-oblonga, confertiuscula : laciniis calycinis orbiculatis.

*Obs.* Priore minus altum, scapo firmiore et rectiore. Flores obsolete pallidi, immutabiles. Rami paniculæ inferiores interdum compositi.

*HAB.* in excelsis montibus Carolinæ septentrionalis.

## A C E R. L.

HERMAPHROD. *Cal.* 5-fidus. *Cor.* 5-petala. *Stam.* 8.

*Pist.* 1. *Caps.* 2 seu 5, monospermæ, ala terminatæ.

MASC. *Cal.* 5-fidus. *Cor.* 5-petala. *Stam.* 8.

SACCHARINUM. A. foliis subpalmato-5-lobis, margine integris : floribus corymbosis : capsulæ lobis turgide ovatis, alis assurgentibus.

*Oss.* Habitus *A. platanoidis*. Folia modo, uti supra dicti, subtus glabella, modo pubentia.

*HAB.* a sinu *Hudson* ad Carolinam et Tennassée.

BARBATUM. A. foliis breviter trilobis, serratis : pedunculis masc. ramosis, foem. simplicissimis : capsulæ alis erectis.

*A. carolinianum* ? WALT.

*Oss.* Flores pallido-viriduli, modo omnes masculi, modo hermaphroditi iis immixti : pedunculi solito pilosi. Calyces masc. intus densissima barba obsiti.

*HAB.* in Carolina.

PENSYLVANICUM. A. foliis inferne rotundatis, superne acuminato-tricuspidibus, argute serrulatis, glabris : racemo simplici, rarifloro.

*Oss.* Gallis *Erable jaspé*.

*HAB.* in Canada ; rarius in montibus *Alléghanis*, Pennsylvaniae et Carolinae.

POLYGAMIA. MONOECIA. 253

MONTANUM. A. foliis 3-sub5-lobis, rugosis, dentatis, subtus pubentibus : racemo composito.

Obs. Flores parvi.

HAB. in Canada, raro in montibus excelsis Alleghaniis, usque ad Floridam.

ERIOCARPUM. A. foliis palmato-5-lobis, inæqualiter dentatis, subtus glabriusculis glaucisque; sinubus obtusis : floribus fertilibus subsessiliter conglomeratis.

A. tomentosum. Hort. paris.

Obs. Fructus junior lanosus; maturus pubens, alis amplissimis, decussato-convergentibus.

RUBRUM. A. foliis palmato-5-lobis, inæqualiter dentatis, subtus pubentibus demumque glaucis; incisuris acutis : floribus fertilibus aggregatis, longiuscule pedicellatis.

Obs. Fructus junior maturusque glaberrimus; alis decussato-convergentibus.

HAB. a Pennsylvania ad Carolinam.

NEGUNDO. A. glaberrimum : foliis pinnatis, ad summum 5-foliolatis : floribus dioicis.

HAB. a Pennsylvania ad Carolinam.

MIMOSA. L.

HERMAPHROD. *Cal.* 5 - dentatus. *Cor.* 5 - fida.  
*Stam.* 5 seu plura. *Pist.* 1. *Legumen.*  
 MASC. *Cal.* 5 - dentatus. *Cor.* 5 - fida. *Stam.* 5 - 10 ,  
 plura.

ILLINOENSIS. *M. inermis*, pumila : caule herbaceo,  
 gracili, divaricato : foliis bipinnatis ; pin-  
 nis 5 - 8 - jugis : capitulis solitariis, breviter  
 pedunculatis : leguminibus paucioribus.

Affinis *M. glandulosae*.

*OBS.* Glandula unica ad infimum par pinnarum. Le-  
 gumina planissima, non arcuata.

*HAB.* in pratensibus regionis Illinoensis.

GLANDULOSA. *M. inermis* : caule herbaceo : foliis  
 multijugo-bipinnatis : pedunculis solita-  
 riis, monocephalis : leguminibus conges-  
 tis ; planis, arcuato-falcatis.

Affinis *M. punctatae*. LINN.

*OBS.* Glandulæ tot quot pinnarum paria.

*HAB.* ad ripas fluvii *Tennassée* ; etiam juxta *Missis-  
 sippi* legit BARTRAM'S.

HORRIDULA. *M.* caule herbaceo, diffuso seu pro-  
 cumbente petiolisque uncinatim aculeo-  
 latis : foliis multijugo-bipinnatis : pe-  
 dunculis geminis, monocephalis : legu-

POLYGAMIA. DIOECIA. 255

minibus densissime aculeato - echinatis,  
quadri- valvibus.

*M. Intsia.* WALT.

*Obs.* Affinis *M. quadri- valvi*. Legumina quasi longo-  
linearia, teretiuscula, recta.

*HAB.* a Virginia ad Floridam.

D I O E C I A.

EMPETRUM. L.

MASC. *Cal.* 3-partitus. *Cor.* 3-petala. *Stam.* longa.

FOEM. *Cal.* 3-partitus. *Cor.* 3-petala. *Styli* 9.  
*Bacca* 9-sperma.

NIGRUM. E. caule procumbente : foliis oblon-  
L. gis, obtusis, margine revolutis : fructu  
nigro, 9-spermo.

*HAB.* in rupibus maritimis Canadæ.

F R A X I N U S. L.

HERMAPHROD. *Cal.* o. seu 4-partitus. *Cor.* o. seu  
4-petala. *Stam.* 2. *Pist.* 1. *Sem.* 1, lanceola-  
tum.

FOEM. *Pist.* 1, lanceolatum.

QUADRANGULATA. F. ramulis quadrangulatis : fo-  
liolis ad summum 4-jugis, subsessilibus,



**POLYGAMIA. DIOECIA.**

ovali-lanceolatis, argute serratis, subtus  
pubentibus : capsulis utrinque obtusis.

*HAB.* in Kentucky et Tennesse.

**PLATICARPA.** F. foliolis ad summum 3-jugis, sub-  
sessilibus, ovalibus, serratis : capsulis  
lato lanceolatis.

*HAB.* in Carolina.

**EPIPTERA.** F. foliolis ad summum 4-jugis, oblongo-  
ovalibus, acuminatis, subintegris : cap-  
sulis obverse lanceolatis, apice obtuso  
emarginatis, ima parte teretibus, ap-  
teris.

*HAB.* in Virginia et Carolina.

**P A N A X. L.**

**HERMAPHROD.** Umbella. *Cal.* 5-dentatus, supe-  
rus. *Cor.* 5-petala. *Stam.* 5. *Styli* 2. *Bacca* 2-  
sperma.

**MASC.** Umbella. *Cal.* integer. *Cor.* 5 - petala.  
*Stam.* 5.

**QUINQUEFOLIUM.** P. radice fusiformi : foliis ternis;  
**L.** foliolis quinis, ovalibus, acuminatis,  
petiolatis.

*HAB.* in excelsis montibus Americæ septentrionalis  
sive a Canada ad Tennesse.

POLYGAMIA. *DIOECRA*. 257

TRIFOLIUM. *P. pusillum* : radice subrotundo-bulbosa : foliis ternis ; foliolis subsessilibus.

*OBS.* Foliola terna , rarius quina.

*HAB.* in Noveboraco et Pennsylvania,

GLEDITSIA. L.

HERMAPHROD. *Cal.* 4-fidus. *Cor.* 4-petala. *Stam.* 6.  
*Pist.* 1. *Legumen.*

MASC. *Cal.* 3-phyllus. *Cor.* 3-petala. *Stam.* 6.

FOEM. *Cal.* 5-phyllus. *Cor.* 5-petala. *Pist.* 1.  
*Legumen.*

TRIACANTHOS. *G. spinosa* : leguminibus oblongis ,  
*L.* polyspermis.

*Var. α. macrocarpos* : legumine longissimo.

— *β. brachycarpus* : legumine brevior ;  
subtriunciali.

*HAB.* in occidentalibus *Alléghanis*.

MONOSPERMA. *G. spinosa* : legumine subrotundo ,  
*WALT.* 1-spermo.

*HAB.* in Carolina , Georgia , Florida et regione Illinoisensi.

DIOSPYROS. *L.*

HERMAPHROD. *Cal.* 4-fidus. *Cor.* urceolata, 4-fida. *Stam.* 8. *Stylus* 4-fidus. *Bacca* 8-sperma.  
 MASC. *Cal.* *Cor.* *Stam.* prioris.

VIRGINIANA. *D.* foliis longe petiolatis, oblongo-  
*L.* ovalibus, acuminatis, subtus pubentibus.

*HAB.* a New-Yorck ad Ludoviciam et in occidentilibus ultra montes *Alléghanis*.

NYSSA. *L.*

HERMAPHROD. *Cal.* 5-partitus. *Cor.* o. *Stam.* 5.  
*Pist.* 1. *Drupa* infera.  
 MASC. *Cal.* 5-partitus. *Corolla* o. *Stam.* 10.

VILLOSA. *N.* foliis ovalibus, integerrimis; petiolo, nervo medio margineque villosis: pedunculis foem. longis, plerumque 3-floris: nuce brevi-obovata, obtuse striata.

*N. multiflora*? *WALT.*

*N. montana.* Hortulanorum.

*Obs.* Interdum cultura villositatem amittit. Affinis  
*N. biflorae.*

*HAB.* a Nova Anglia ad Carolinam.

POLYGAMIA. *prostrata*. 259

BIFLORA. N. foliis oblongo-ovalibus, integerri-  
mis : pedunculis foem. bifloris : drupa  
WALT. brevi-obovata ; nuce obtuse striata.

OBS. Folia adulta glabella et coriaceo-laurina. Pedun-  
culi foem. rarius 1-flori.

*N. aquatica?* LINN.

HAB. in stagnis Virginiae rarius ; frequentius Caro-  
linae et Georgiae.

CANDICANS. N. foliis brevissime petiolatis, subcu-  
neato-oblongis, subtus subcandicanti-  
bus ; pedunculis foem. 1-floris : bracteolis  
calycisque tomentosí laciniis brevibus :  
drupa oblonga.

*N. capitata?* WALT.

OBS. Variat foliis obovalibus ; integris et rariter sub-  
dentatis. Flores masculi in capitulum conglobati.

HAB. ad amnem Ogechee.

TOMENTOSA. N. foliis longe petiolatis, subtus to-  
mentosus, ovalibus, acuminatis : pe-  
dunculis foem. 1-floris : drupa oblonga.

OBS. Folia passim acute grandi-dentata. Bracteolæ  
ovario sublongiores. Calycis lacinae cuneatæ.

HAB. ad amnem S. Mary et in Florida.

ANGULISANS. N. foliis longe petiolatis ; inferiori-  
bus subcordatis ; plerisque ovalibus ,

**POLYGAMIA. DIOECIA.**

**acuminatis; rariter angulato-dentatis :  
pedunculis fœm. 1-floris : drupa oblonga.**

***N. uniflora.* WALT.**

***HAB.* in aquosis Carolina.**

---

CLASSIS XXIV.

CRYPTOGAMIA.

FILICES.

§. I. ANNULATE PERIGENESÆ.

VITTARIA. SM.

FRUCTIF. Linea continua sub replicato frondis margine, ab istius sinu distincta.

ANGUSTIFRONS. V. frondibus simplicibus, integerrimis, angustissime longissimeque linearibus, debilibus.

HAB. in Florida, juxta amnem *Aisa-hatcha*.

PTERIS. L.

FRUCTIF. Linea continua sub membrana quasi ab ipso margine replicato facta; *sporangii* ex ipsomet membranæ sinu oriundis, directione introrsa.

ATROPURPUREA. P. parvula : stipite castaneo, villosiusculo : fronde remote pinnata; pinnis suboppositis, stipitatis, obtuse-ovalibus etiamve lineari-lanceolatis, inte-

262 CRYPTOGAMIA. *PSILICES.*

gerrimis ; inferioribus modo auritis ,  
modo 3-5-foliolatis.

PLUCK. t. 401. f. 3.

*HAB.* in montibus *Alleghensis* saxosis.

GRACILIS. *P. pusilla* , glabra : stipite debili , graciliter setaceo : fronde subbipinnata ; pinnis pinnatifidis ; laciniis ovali-lanceolatis integrisque , sterilibus rotundioribus , crenulatis.

*HAB.* in rupibus *Canadæ* , juxta *Malbaye*.

AQUILINA. *P. magna* : stipite glabro : fronde tripinnata ; pinnulis contigue adnatis , ovalibus , integris ; pinnis supremis linearibus.

*HAB.* a sinu *Hudson* ad *Floridam*.

**ADIANTUM. L. Sm.**

**FRUCTIF.** Lineolæ sub totidem membranulis marginalibus introrsum dehiscentibus et quasi creniformibus.

**PEDATUM.** *A. glaberrimum* : fronde pedata ; partialibus pinnatis ; pinnis subtriangulis , superno latere incisis , lacinulis subquadratis , apice truncato fructificantibus.

*HAB.* in Pennsylvania, &c.

**§. II. ANNULATE HYPOGENESÆ.**

**WOODWARDIA. Sm.**

**FRUCTIF.** Linea utrinque nervo medio adjecta , interrupta ; partialibus oblongis , membrana introrsum dehiscente singulatim contextis.

*OBS.* Genus vix a sequenti distinguendum.

**BANISTERIANA.** *W. fronde pinnata ; pinnis pinnatifidis* : fructificatione juxta nervum pinnarum pinnularumque interrupta.

*PLUCK.* t. 179. f. 2.

*HAB.* in montibus Carolina.



BLECHNUM. L.

FRUCTIF. Linea utrinque nervo medio adjecta ,  
continua , et membrana introrsum dehiscente  
primum oblecta.

SERRULATUM. B. majusculum , stricte erectum :  
frondibus pinnatis ; pinnis longo-lanceo-  
latis , acutis , basi minime dilatatis , ar-  
gute rigideque serrulatis.

HAB. in Florida , juxta amnem *Aisa-hatcha*.

ASPLENIUM. L. SM.

FRUCTIF. Lineæ in disco frondis rectæ , obliquatæ ,  
distinctæ ; membranula ex altero margine de-  
hiscente primum oblectæ.

RHIZOPHYLLUM. A. pusillum , glabrum : frondibus  
L. simplicibus , subhastato-lanceolatis , in-  
tegris , summitate cirrhosa radicanibus :  
fructificatione lineis brevibus , inordi-  
nate sparsis.

HAB. in montium rupibus , a Canada ad Tennassée.

TRICHOMANES. A. parvulum , glaberrimum : sti-  
L. pitibus atro-fuscis : frondibus pinnatis ;  
pinnulis suborbiculatis , crenulatis.

HAB. in Canada , Pensylvania , et in excelsis mon-  
tibus Carolinæ.

CRYPTOGAMIA. *FILICES*. 265

TRICHOMANOIDES. A. fronde pinnata, glabra; pin-  
L. nis sessilibus, subhastato-oblongis, obtu-  
siusculis, leviter serratis : fructificatione  
lineis brevissimis.

PLUCK. t. 89. f. 8. et t. 287. f. 2. var. minor.

MORIS. III. s. 14. t. 2. f. 12.

*HAB.* in Carolina.

ANGUSTIFOLIUM. A. glabrum : fronde pinnata ;  
pinnis remote alternis, longo-linearibus,  
integris : fructificatione lineis densis, to-  
tum fere discum obtegentibus.

*Lonchitis virginiana*, &c. MORIS. III. s. 14.  
t. 2. f. 25.

*HAB.* ad ripas fluminis *Ohio*.

THELYPTERIOIDES. A. majusculum , glabellum :  
fronde pinnata ; pinnis longis , pinnati-  
fidis ; laciniis ovalibus , rotundato-obtu-  
sis , levissime denticulatis : fructifica-  
tione lineis breviusculis , æqualibus ,  
oblique approximato-parallelis.

*OBS.* Habitus *POLYPODII Thelypteridis*, LINN.

*HAB.* in montibus Virginie et Carolinæ septentrio-  
nalis.

ADIANTUM NIGRUM. A. parvulum : fronde glabra ,  
L. bipinnata ; pinnulis ovalibus seu lanceo-

CRYPTOGAMIA. *FILICES*.

latis, subpinnatifidis; lobulis ad apicem denticulatis.

*Obs.* Europæo multoties minus.

*HAB.* in altis montibus Carolinæ.

**RUTA MURARIA.** A. parvulum, glabrum: frondibus alternatim decompositis; pinnulis cuneatis, crenulatis.

*Obs.* Europæo multo minus.

*HAB.* in fissuris rupium montium excelsorum Carolinæ septentrionalis.

**NEPHRODIUM.** *Hort. med. paris. Cat.*

*Nεφρος, Ειδος:* fructificatio punctis subreniformibus.

**POLYPODIUM.** *L.*

**FRUCTIF.** Puncta in disco frondis sparsa seu seriata; primum oblecta membranula reniformi aut lunata, hinc (servato nexu laterali) dehiscente.

*Obs.* Præterea, bina ex **POLYPODIIS LINNÆI** excerpanda sunt genera:

1°. **CYATHEA.** *Sm.*

Puncta sphaeroidea, inferne membranula cupulatum circum cincta.

2°. **HYPOPELTIS.**

Puncta subrotunda; *sporangis* circum stipitem membranulæ peltatæ affixis.

CRYPTOGAMIA., *FILICES.* 267

**ACROSTICHOIDES.** N. stipite paleaceo : fronde longa, pinnata ; pinnis alternis , subsessilibus , basi hinc auriculatis , levissime serratis , ciliatis ; superioribus tantum fructificantibus.

**OBS.** Affine *POLYP. auriculato.* LINN. Fructificatio primum punctis crebris , demum totum discum dense obtegens.

**HAB.** in Pennsylvania , Carolina et Tennassée.

**THELYPTERIOIDES.** N. stipitibus erectis , glabris : fronde pinnata ; pinnis sublinearibus , pinnatifidis ; pinnulis oblongiusculis , subintegris : punctis minusculis , juxta marginem pinnularum seriatis.

**OBS.** Valde affine *POLYP. Thelypteridi* ; punctis etiam maturitate non ita coeuntibus. Frons pubens , praesertim in scapo nervoque pinnarum medio ; longa ; remote pinnata.

**HAB.** in Canada et in montibus *Alleghanis.*

**MARGINALE.** N. stipite insigniter paleaceo : fronde glabra , bipinnata ; pinnis distantibus ; pinnulis angustissime decurrentibus , rotundato-obtusis , crenatis : punctis ad singularum crenarum incisuras marginalibus.

*POLYPOD. marginale.* LINN.

**HAB.** a Canada ad Kentucky et Virginiam.

268 CRYPTOGRAMIA. *FILICES.*

**PUNCTILOBULUM.** N. majusculum : stipite nudo, ramis pinnulisque pubescentibus : fronde longa , bipinnata ; pinnulis decurrentibus , subovali-oblongis , semi et ultra pinnatifidis ; lobulis oblongiusculis , apice 2-4-dentatis , singulis unipunctiferis.

*Obs.* Habitus **POLYP.** *Filicis feminae*, LINN.

*HAB.* in Canada.

**BULBIFERUM.** N. glabrum : fronde bipinnata , passim bulbifera ; pinnis distanter suboppositis ; pinnulis oblongiuscule ovalibus , obtuse inciso-dentatis , dentibus subbidenticulatis.

*Obs.* Habitus fere **POLYP.** *Filicis masc.*

*HAB.* in Canada.

**FILIX FÆMINA.** N. majusculum , glabrum : fronde oblonga , bipinnata ; pinnis elongatis ; pinnulis lanceolatis , subbipinnatifidis seu inciso-serratis : punctis utrinque juxta nervum sublunatis.

*HAB.* in Canada.

**ASPLENIOIDES.** N. majusculum , glabrum : stipite nudo : fronde bipinnata ; pinnulis subovali-oblongis , inciso-dentatis ; dentibus inferioribus obtusis , supremis mutice

CRYPTOGAMIA. *FILICES.* 269

acutis : punctis utrinque juxta nervum lunatis.

*OBS.* Affine *POLYP. Filici* fœm. et forsan varietas. Puncta ante maturitatem quasi lineolæ plerumque arcuatæ; ita ut proximitatem ad *ASPLENIUM* indigent.

*HAB.* a Nova Anglia ad Carolinam.

**CRISTATUM.** N. majusculum : stipite paleaceo : fronde glabra, subtripinnata; pinnulis subovali-oblongis, acutis, pinnatifidis; lobulis argute dentatis, dentibus mucronatis : punctis minusculis, distinctis, subreniformibus.

*POLYP. cristatum?* LINN.

*HAB.* in Canada.

**TENU.** N. pusillum, glaberrimum, debile : fronde bipinnata; pinnulis anguste decurrentibus, ovalibus, retusis, leviuscule dentatis : punctis majusculis, ad singulos dentes singulis.

*OBS.* Valde affine *POLYP. fragili.* LINN.

*HAB.* in Canada.

**RUFIDULUM.** N. pusillum, totum rufo-villosissimum : fronde subbipinnata; pinnis pin-

CRYPTOGAMIA. *FILICES.*

natifidis ; pinnulis lobulisve ovalibus,  
retusis , subcrenatis.

Affine *N. lanoso*.

*OBS.* Fructificatio demum confluens et villositate totum pinnularum discum quasi refo-tomentosum efficiens.

*HAB.* in rupibus Canadæ et Novæ Caesariæ.

**LANOSUM.** *N.* parvulum ; elegans ; totum lanosissimum : fronde oblonga , bipinnatifida ; pinnis distantibus ; pinnulis pinnatifidis ; lobulis subrotundo-ovalibus , integris : punctis demum contiguis.

*OBS.* Habitus quodammodo *POLYP. fontani* ; paulo majus.

*HAB.* in montibus saxosis Tennassée et Carolinæ septentrionalis.

**DRYOPTERIS.** *N.* parvulum : fronde bipinnata ; pinnis oppositis ; pinnulis pinnatifidis ; laciniis oblongis , obtusis , leviter crenatis : punctis submarginalibus , demum contiguis.

*POLYP. Dryopteris. LINN.*

*OBS.* Frons circumscriptione quasi subæquilateraliter triangula : pinnarum par primum (interdum et alterum) longiuscule stipitatum.

*HAB.* in Canada.

POLYPODIUM. L. SM.

FRUCTIF. Puncta subrotunda, nudissima, in disco frondis sparsa seriatave.

VULGARE. P. stipite glabro : fronde pinnatifida ;  
L. pinnis oblongis, levissime subdentatis : punctis grossis, utrinque a nervo medio uniseriatis.

*P. virginianum?* LINN.

*HAB.* a Canada ad Carolinam.

CETERACCINUM. P. fronde pinnata ; pinnis oblongis, integerrimis, obtusis, subtus stipiteque squamulosis : punctis utrinque uniseriatis.

*ACROSTICHUM polypodioides*. LINN.

*HAB.* parasiticum in Kentucky, Tennesse, Florida.

HEXAGONOPTERUM. P. fronde pubente ciliataque, bipinnatifida ; membranis pinnae oppositas connectentibus oblongo-hexagonis ; pinnulis inciso-crenatis : punctis sparsis.

PLUCK. t. 284. f. 2.

*OBS.* Frons quasi subæquilateraliter triangula. Puncta vix perceptibilia, paucissime granulosa.

*HAB.* in Carolina et Virginia.

CONNECTILE. P. stipite paleaceo : fronde pubente ciliataque, bipinnatifida ; pinnis opposi-



272 CRYPTOGRAMIA. FILICES.

tis, adnatis, contiguus; pinnulis sub-  
ellipticis: punctis minutis.

HAB. in Canada.

ACROSTICHUM. L. SM.

FRUCTIF. *Sporangia* indefinite nudeque discum  
frondis obtegentia.

AUREUM. A. maximum, glabrum: fronde pin-  
L. nata; pinnis stipitatis, oblongis, in-  
tegerrimis.

Obs. Variat pinnis subacutis.

HAB. in Florida, ad amnem *Aisa-hatcha*.

ONOCLEA. L.

FRUCTIF. Frondis fertilis pinnulæ subtus convo-  
lutæ et quasi conversæ in totidem urceolos bise-  
riatos occultantesque numerosa *sporangia*.

SENSIBILIS. O. pinnis connatis, crenato-incisis:  
L. fronde fertili urceolis distinctis.

HAB. a Pennsylvania ad Georgiam.

NODULOSA. O. pinnis sessilibus, linearibus, pin-  
natifidis: fronde fertili pinnis quasi no-  
duloso-articulatis; urceolis contiguus.

ACROSTICHUM *areolatum*. LINN.

OSMUNDA *caroliniana*. WALT.

HAB. in Carolina limosis umbrosis,

§. III. EXANNULATÆ.

OSMUNDA. L. Sm.

FRUCTIF. *Sporangia* numerosa, nuda, indefinite  
seu quasi acervatim receptacula sua contegentia.

REGALIS. O. magna, glabra : fronde bipinnata ;  
L. pinnulis distinctissimis, oblongis, subserrulatis : stipite summo composite racemifero.

HAB. a Canada ad Floridam.

CINNAMOMEA. O. frondibus sterilibus pinnatis ;  
L. pinnis elongatis, pinnatifidis ; laciniis subovalibus, integriusculis : frondibus fertilibus oppositè racemiferis.

MORIS. III. s. 14. t. 4. f. 3.

OBS. Magna, lanugine gossypina et rufidula vestita.

HAB. a Nova Anglia ad Carolinam.

INTERRUPTA. O. majuscula : stipite glabro : frondibus pinnatis ; pinnis oppositis, pinnatifidis ; laciniis subovalibus, integriusculis : pinnis aliquot intermediis fructificantibus.

HAB. in Canada, Pennsylvania et Kentucky.

BOTRYPUS. *Hort. med. paris. cat.*

*Stipes* supra radicem 1-2-frondifer; ultra frondem continuatus in pedunculum simplicem duplicemve, nudum, summitate racemiferum.

FRUCTIF. Racemus plus minus compositus; *sporangii* duplici serie ordinatis, discretis, omisso nexu postico liberis, nudis, globosis, transverse æqualiter bivalvibus.

*Obs.* Est quasi OPHIOGLOSSUM fronde spicaque compositis.

VIRGINICUS. B. stipite villosiusculo : fronde sub3-pinnata; pinnulis lanceolato-ovalibus, inciso-dentatis : sporangiis minutis.

OSMUNDA *virginica*. Linn. excluso PLUMIERI synonymo. MORIS. III. s. 14. t. 4. f. 5.

*HAB.* a Canada ad Virginiam et in montibus Carolinæ.

LUNAROIDES. B. pumilus, glaber : fronde subtrichotome 3-pinnata; pinnulis reniformi-rotundatis, leviter crenulatis : racemo brevi, subtrichotomo.

*Obs.* Pinnæ secundariæ frondem LUNARIÆ æmulantur, multo minores.

*HAB.* in pascuis, circa Charlestown.

## CTEISIUM.

*Krus*, pecten: *frondes propter spicillas fructificantes quasi pectinatim marginatae.*

**FRUCTIF.** Frons scandens, composita: spiculæ lineares, subtus a reflexis utriusque marginis lobulis bifariam imbricatæ. *Sporangia* singulatim sub singulis lobulis retenta, transversim ovoidea, ab apice interno dissimili ad externum bivalvihiliscentia.

**PANICULATUM.** C. frondibus partialibus suborbiculatis, basi sinu profundo emarginatis, 5-6-lobo-palmatis, lobis subæqualibus, ovali-oblongis, obtusis: spicillis compositis paniculatis.

*HAB.* in occidentalibus Virginiae, ad fines Kentucky et in Tennassée.

## OPHIOGLOSSUM. L.

**FRUCTIF.** Spica simplex, linearis, utroque margine quasi articulato unifariam multilocularis; loculis sutura transversa medianaque extrorsum dehiscentibus.

**AULGATUM.** O. fronde ovali; scapo elongato: spica longo-lineari.

*HAB.* in Nova Cæsarea.

276 CRYPTOGRAMIA. HEPATICÆ.

BULBOSUM. *O.* pusillum : radice subgloboso-bulbosa : fronde subcordato-ovali , obtusa : spica brevi, mucronata.

*O. crotalophoroides* ? WALT.

Obs. *O. lusitanico* paulo majus.

HAB. in sabulosis Carolinæ inferioris.

H E P A T I C Æ.

M A R C H A N T I A. L.

*Sporangia* sub radiis pilei solitarii et peltato-stipitati propendentia , irregulariter ruptilia.

TENELLA. *M.* pusilla , brevifrons : stipite capillari : pileo conoideo, inferne membranaceo - multifido ; laciniis ligulato - linearibus.

HAB. in Canada.

CRINITA. *M.* pusilla : fronde margine colorato erosa , e disco lobulorum rotundatorum fructificante : stipite brevissimo : pileo subgloboso , 5-lobo , ex incisuris prolixè filamentoso.

Obs. *Frons* vetuli TARGIONIÆ ; subtus , marginibus , sæpeque etiam supra atro-purpureo imbuta. *Stipes* 3-4 lin. altus : *pilei* lobi rotundati.

Affinis *M. hemisphæricæ*.

HAB. in versuris fossarum sylvaticarum Carolinæ.

**CRYPTOGAMIA HEPATICE. 277**

**POLYMORPHA.** M. stipitibus marginalibus : pileis.

**L.** stérilibus plano-convexis, crenatis; fertilibus stellato-multipartitis : scyphulis campanulatis.

**HAB.** varietates  $\alpha$  et  $\beta$  promiscue crescent in humidis Canadae.

**JUNGERMANNIA. L.**

*Sporangium* e vagina perigonica pedunculato-exertum, 4-valve.

\* *Pinnulis distichis, inauriculatis.*

**VITICULOSA. JUNGERMANNIA.**

**L.**

N<sup>a</sup>. Sterilem tantum offendi; qualem delineat DILL.  
t. 69. f. 7. c.

**HAB.** in Canada.

**ORBICULARIS.** J. perpusilla, repens : foliis orbiculatis, integerrimis; duplici serie connivente verticaliter assurgentibus : pedunculo caulino, brevi : sporangio ovato.

**OBS.** J. *bienspidata* paulo minor. *Folia*, primo obtutu, quasi lenticularia; neque auriculata; neque basi replicata; glabra; binis ordinibus supra aurculos serpentes sese convergendo erigentia, singulorum ordinum sibi lateraliter incumbentia.

**HAB.** in provincia New-York.

**\*\*** *Pinnulis distichis, auriculatis.*

**TRIDENTICULATA.** J. procumbenti-repens : foliis complanato-distichis, auriculatis, majusculis, subovalibus, apice 3-denticulatis : pedunculis caulinis, longis.

Affinis DILL. t. 70. f. 15 ?

*Oss.* Majuscula; ramis in eodem plano ac caules patulis. *Folia* præter apicem integerrima. Vagina perigonalis angusto-oblonga. *Sporangium* ovatum.

**VARIA.** J. surculis erectiusculis : foliis bifidis, subdistiche conduplicato-æqualibus, integerrimis : pedunculo terminali, longiusculo : sporangio ovato.

*Oss.* Mea varietas dimidio minor quam Europæa. *Folia* conduplicato-bifida; altero lobo auriculam constituentem laciniæ superiori subæquali; unde foliatio depressa et quasi quadrifariam imbricata.

*HAB.* in Canada et in monte *Grand-Father* Carolinæ.

**SETULAROIDES.** J. repens, capillaris : foliis setaceo-bipartitis; setis quasi articulatis : sporangiis caulinis : perigonii dentibus apice setuliferis.

*HAB.* in montibus Carolinæ, in diversarum congenerum cæspitibus prorepens.

**LACINIOSA.** J. repens, composite ramosissima : foliis bifidis, profunde fimbriato-laci-

**CRYPTOGAMIA. HEPATICÆ. 279**

niosis : perigoniis lateralibus , oblongis,  
glabellis : sporangiis brevi-pedunculatis,  
globosis.

*Obs.* Affinis *J. ciliari*. *Caulibus* protractis copiosisque  
ramulis intricatim repens. *Folia* minuta , acute  
bifida , profunde et inæqualiter laciniosa; laciniis  
subpiliformibus , recurvatis. *Pedunculi* vix dupla  
perigoniorum conspicuorum longitudine.

*HAB.* in Canada , cortici arborum adnata.

**TOMENTELLA.** *J. assurgens* ; ramis distiche 2-3-

*ERH.* pinnatis : foliis lanuginiformibus.

*Obs.* Fructificantem non vidi.

*HAB.* in paludosis sylvaticis Carolinæ.

\*\*\* *Foliis imbricatis.*

**TAMARISCI.** *J. distiche* compositeque ramosa ; lu-

*L.* cidule fusca : foliis rotundatis , convexis ,  
bifariam imbricatis , anguste auriculatis ;  
foliolis emarginatis , subtus uniseriatis :  
perigonio terminali.

*HAB.* in arboribus montosæ Carolinæ.

**PEATIPHYLLA.** *J. distiche* compositeque ramosa :

*L.* imbricata ; supra , foliis subovalibus , bi-  
fariis ; subtus , auriculis ovalibus , inte-  
gris , consimilibusque foliolis trifariis :  
perigoniis lateralibus , subgloboso - ur-  
ceolatis.

*Obs.* *Sporangium* sæpe 5-6-valve , pariter ac in Eu-  
ropæa.

*HAB.* in Canada et in montibus Carolinæ.



280 CRYPTOGRAMIA. HEPATICÆ.

\*\*\* *Fronde laminata.*

FURCATA. J. fronde lineari, ciliata; extremita-  
L. tibus obtusis, furcatis.

*HAB.* in arboribus Carolinæ.

ANTHOCEROS. L.

Frons laminata. Vagina perigonica emittens *sporangium* corniforme, maturitate bivalve.

CAROLINIANUS. A. frondibus breviter rotundato-  
lobatis, margine subintegris, supra vix  
puncticulatis : corniculis subsemiuncia-  
libus.

*HAB.* in humidis Carolinæ inferioris.

*EQUISETACEÆ.*

EQUISETUM. L.

*Stipes* caulescens; articulationibus (frondis loco) vaginiferis.

*Sporangia* sub peltis in conum collectis.

*Sporae* spiralliter involutæ filis (vascularibus) subita materiæ connectentis ruptura extensilibus.

SCIRPOIDES. E. cauliculis simplicibus, setaceis : vaginis trisetis : capitulo brevissimo, nigricante.

*Obs.* Pusillum, ab radice confertim ramosum et inde quasi multicaule.

*HAB.* in vetustis sylvis Canadæ.

*LYCOPODACEÆ.*

PSILOTUM. Sw.

*Sporangia* globosa, subtrilocca (rarius 4-cocca) sparsa, axillari-lateralia, sessilia, 5-locularia, valvulis tribus apice dehiscentibus.

*Sporae* numerosissimæ.

FLORIDANUM. P. frondis furcationibus ultimis breviusculis : sporangiis stipitulatis.

*Obs.* Species hujus generis difficillime definiendæ ; usquedum a pluribus sedulo collatis, eruantur differentiæ.

*HAB.* in Florida.

LYCOPODIUM. Sw.

*Sporangia* reniformia, axillaria, sessilia, bivalvia,  
elastice dehiscentia.

*Sporae* numerosæ, minutissimæ.

CLAVATUM. L. caule repente : foliis confertis, su-

L. bulatis bracteisque sensim apice piliferis :  
pedunculis ramulorum terminalibus, ple-  
rumque distachyis.

*Obs.* Pedunculi sæpe monostachyi.

*HAB.* in Canada.

DENDROIDEUM. L. surculis assurgentibus, inferne  
simplicibus nudiusculisque, superne den-  
droideo-ramosis ; ramulis sessiliter 1-sta-  
chyis : foliis lineari-lanceolatis, integer-  
rimis : spica crassa, compacte imbricata ;  
bracteis abbreviato - dilatatis, mucro-  
natis.

DILL. t. 64.

*HAB.* a Canada et Nova Anglia ad Carolinam mon-  
tosam.

ALPINUM. L. repens ; surculis dichotome ramo-

L. sissimis, fertilibus altioribus : foliis 4-fa-  
riam appressis imbricatisve, lanceolatis,  
convexis : spicis dichotomias solitarie  
terminantibus ; bracteis subcordatis, acu-  
minatis, integris.

*Obs.* DILL. t. 58. f. 2. quadrat.

*HAB.* in Canada.

CRYPTOGAMIA. LYCOPODACEÆ. 283

COMPLANATUM. L. surculis passim dichotome ramulosis ; ramulis alato-complanatis : pedunculis longiusculis , subnudis ; spicis teretibus.

Affinis *L. alpino*.

*Obs.* Ramuli sæpe cæteris complanatis intermixti occurrunt foliis 4-fariam imbricatis. Spicæ interdum 5.

*HAB.* in Canada et Carolina.

ANNOTINUM. L. caulibus procumbentibus : foliis lanceolato-linearibus , serrulatis , patulis : spicis terminalibus , solitariis ; bracteis brevi-cordatis , eroso-denticulatis.

*HAB.* in Canada.

INUNDATUM. L. caulibus radicante-repentibus : scapis radicalibus , brevibus , foliosis , monostachyis : foliis confertis , subulatis , integris : spica bracteis foliiformibus.

*HAB.* a sinu *Hudson* ad lacum *Mistassins*.

ALOPECUROIDES. L. repens : foliis confertis , patulis , subulatis , ciliosis : scapis radicalibus , altioribus , dense foliosis : spica bracteis longissimis dense foliosa.

*Obs.* Rami recurvati sæpe summitate radicales.

*HAB.* in humidis herbosis Carolinæ.

CAROLINIANUM. L. caule radicato-repente : foliis subfalcato-lanceolatis , integerrimis : scapis nudiusculis , longissimis , mono-

284 CRYPTOGAMIA. LYCOPODACEÆ.

stachyis : spica longa ; bracteis sublan-  
ceolatis , integris.

*HAB.* in pratensibus udis Carolinæ.

SELAGINOIDES. *L. parvulum* , procumbens ; sur-  
*L.* culis fertilibus assurgentibus : foliis pa-  
tentibus , ovali-lanceolatis , ciliosis : spi-  
cis solitarie terminalibus , subalternatim  
foliosis.

*HAB.* in borealibus Canadæ.

APODUM. *L. pusillum* : foliis remotiusculis , sub-  
*L.* rotundo-ovalibus , membranaceis , bifa-  
riis : summitatibus subspicatum fructifi-  
cantibus.

*HAB.* in umbrosis sylvarum herbosis , a Virginia ad  
Floridam.

RUPESTRE. *L. pusillum* , Hypniforme , repens :  
*L.* foliis confertim incumbentibus , subula-  
tis , ciliatis : spicis sessiliter terminalibus ,  
tetragono-prismaticis.

*HAB.* in rupibus montanis , a sinu *Hudson* ad Virgi-  
niam et Carolinam montosam.

LUCIDULUM. *L. caulibus* assurgentibus rariter lon-  
giusculisque ramosis : foliis patentibus ,  
lucidis , lineari-lanceolatis , serratis : fruc-  
tificationibus caulino-axillaribus.

*Obs.* Affine *L. Selagini* ; pariter passim bulbiferum.

*HAB.* a Canada ad Carolinam montosam.

## MUSCI.

## SPHAGNUM. HEDW.

*Peristomium* nullum.

*Flos masculus* clavatus terminalis ramorum superiorum : *fæmineus* inter ramulos fastigiatos in summitate trunci ; hinc nonnunquam fructus maturi in ramorum superiorum alis : uterque distincti individui.

VULGARE. S. ramulis teretibus, deflexis : foliis brevi-ovalibus, obtusiusculis, concavioribus ; perigonalibus obtusis.

*S. latifolium.* HEDW.

HAB. in Canada, Carolina inferiore, &c.

GRACILE. S. ramulis gracilibus, deflexis, subsquarrosis : foliis minutis, crebrioribus, lanceolatis, acutis ; perigonalibus acuminatis.

*S. capillifolium.* HEDW.

HAB. in Canada.

## GYMNOSTOMUM. HEDW.

*Peristomium* nullum.

*Flos* masculus *femineusque* terminalis.

**TURBINATUM.** G. brevi caule, simplex : foliis lanceolatis, apiculatis, serrulatis : sporangio oblongiuscule turbinato ; operculo convexo, submamillato.

*OBS.* Habitus et magnitudo *G. pyriformis* ; pedunculo longiore ; *sporangio*, excusso operculo, magis truncato minusque crasso.

*HAB.* in Carolina inferiore.

**PROREPENS.** G. stipite repente ; cauliculis erectis, *HEDW.* brevibus, obtusis : foliis confertissime imbricatis, ovali-lanceolatis, acuminatis ; sporangio globoso ; operculo subulato.

*OBS.* Dense cauliculosum, obscure olivaceum. *Folia* integerrima, nervo valido. *Operculum* longitudine sporangii, abrupte longeque acuminatum, erectum.

*HAB.* in Carolina.

## ANICTANGIUM. HEDW.

*Peristomium* nullum.

*Flos masculus* alaris.

FILIFORME. *A.* caulibus gracili-filiformibus : foliis  
arcte imbricatis, ovalibus, in acumine  
concolore denticulatis; perigonalibus  
apice ciliosis : operculo subconoideo : ca-  
lyptra subvillosa.

*Obs.* Affine *A. ciliato*; minus et multo gracilius,  
apice foliorum non scarioso.

*HAB.* in rupibus Carolinæ et Canadæ.

## TETRAPHIS. HEDW.

*Peristomium* simplex : dentibus quatuor pyrami-  
dalibus.

PELLUCIDA. *T.* caule simplici; foliis sporangiferi  
*HEDW.* lanceolatis, cæterorum ovalibus.

*HAB.* in Canada et in monte *Grand-Father* Carolinæ.

## SPLACHNUM. HEDW.

*Peristomium* simplex octodentatum seu octopa-  
rium.

*Sporangium* apophysii conoideæ aut umbraculi-  
formi insidens.

*Flores hermaphroditi* dimorphi : disciformium  
genitalia foeminea infœcunda, elongatorum fer-  
tilia.

SETACEUM. *S.* caulibus tomento radicali confertim



cohærentibus : foliis lanceolatis , sensim  
setaceo-acuminatis , serratis : pedunculo  
non exerto : apophysi clavato-turbinata.

*Obs.* Valde affine *S. angustato* , forsânque ejusdem  
varietas. Habitus quasi *ANICTANGII lapponici*. *Caules*  
graciles 1 - 2 - unciales , pertinaciter fere adusque  
summitates cohærent. *Pedunculus* brevissimus , cir-  
citer longitudine foliorum perigonialium. *Calyptra*  
longo-conica , pallens ; acumine capillari , fusces-  
cente. *Peristomii* flaviduli dentes ecto , biparti-  
biles.

*HAB.* in paludibus Canadæ.

#### WEISSIA. HEDW.

*Peristomium* simplex , 16-dentatum : denticulis  
reflexilibus.

*Flos masculus* terminalis.

MICROODONTA. *W.* subacaulis , simplex : foliis li-

*HEDW.* neari-subulatis , insigniter cirrhato-in-  
volutilibus : pedunculo pallido : sporan-  
gio oblongo-ovato ; operculo abrupte  
longeque acuminato.

*Obs.* Foliatio dilute virens , constanter arefactione cir-  
rhata. Magnitudo varians , 3-6 lin. *Operculum* e la-  
tiore basi abrupte setaceum ; modo sporangio sub-  
brevius , modo eodem paulo longius ; solito basi  
rubellum cæteroque pallens ; interdum fere totum  
rubens. *Peristomii* rubentis denticuli minuti et bre-  
vissimi.

*HAB.* in Carolina.

## TREMATODON.

*Fructificatio* utraque terminalis.

*Sporangium* stipitatum.

*Peristomii* simplicis dentes 16, distincti, subulati, recti, foraminulis pertusi.

LONGICOLLIS. T. brevicaulis, simplex : foliis subulato-setaceis : pedunculo longissimo, tortuoso : sporangii oblongi stipite elongato ; operculo e basi conica graciliter rostrato.

*Obs.* *Caules* simplices, 2-3 lin. alti. *Folia* dilute viridula, fere capillaria. *Pedunculus* 1-2-uncialis, tenuiter capillaceus, pallidus, flexuoso-tortilis. *Sporangium* subcroceo-flavum, cylindraceo-oblongum, obliquatum ; inferne angustatum in collum seu stipitem ipso duplo triplove longiorem arcuatumque : *operculum* e basi eximie conica in longum gracileque acumen desinens : *Peristomii* subobliqui dentes purpurei, foraminulis seriatim pertusi.

Affinis DICRANO *ambiguo*, Hedw. sed distinctissimus.

*HAB.* in arenosis Carolinae.

## GRIMMIA. HEDW.

*Peristomium simplex* : dentibus sedecim , latiusculis , reflexilibus.

*Flores masculi* axillares gemmacei , terminales capituliformes in eadem planta.

**GRACILIS.** *G. graciliter caulescens* : foliis lanceolatis, in acumen subpiliforme, exalbentipellucidum scabriusculumque desinentibus : pedicello terminali ; sporangio brevi-ovato, lævi.

*Obs.* *Caules* cespitosi , erecti , subramosi , graciles ; 3-9 lin. alti. *Folia* obscure viridia , minutula , sensim quasi in pilum decolorem , obsolete asperiusculum desinentia. *Pedunculus* 2 lin. longus , capillaris , pallidus. *Sporangium* subrotundo-ovatum , læviusculum , flavum : *operculum* rubellum , e lata et subconvexa basi mediocriter rostellatum. *Peristomii* simplicis denticuli minuti , brevissimi , obsolete purpurascens , erecti.

*HAB.* in Carolina ; prior detexit D. BEAUVOIS.

## PTERIGYNANDRUM. HEDW.

*Peristomium* simplex, 16-dentatum.

*Flos* masculus *fœmineusque* alaris.

TRICHOMITRION. P. ramificatione compressa : foliis ovalibus, acuminatis, longitudinaliter striatis : pedunculo perigonii longitudine : sporangio ovato : calyptra prolixè pilosa.

OBS. *Cauliculi* subbiunciales, breviuscule et composite ramulosi, ramis erectis. *Folia* confertissime imbricata, superne erecto-patentiuscula, tenuiter acuminata, longitudinaliter 3-5 plicato-striata. *Perigonia* longa, pedunculos sæpe breviores involucrantia. *Sporangia* erecta, rufidulo-flava. *Peristomii* denticuli minuti, pallidi. *Calyptra* sporangium adhuc dum perigonio immersum deserens.

HAB. in montibus Carolinæ.

JULACEUM. P. cauliculis erectis, teretibus : foliis crebro-imbricatis, ovalibus, acuminatis, lævibus : pedunculis perigonio longissimo vix emersis : sporangiis breviovatis; operculo conico : calyptra glabra.

OBS. Habitus FISSIDENTIS *sciuroidis*; subdimidio minor. *Pedunculus* erectus, brevis, perigonio angustolongo vix æqualis. *Peristomii* denticuli pallidi, subulares.

HAB. in truncis arborum Carolinæ.

**HIRTELLUM.** P. ramulis gracili-filiformibus, simplicibus, brevibus : foliis cylindraceo-imbricatis, subcordatis, hirtellis, fimbriolatis : sporangio oblongo; operculo conoideo-elongato.

**Obs.** Minusculum gracileque. *Folia* minutissima, piloso-acuminata; extus convexa et minutissime pulverulento-hirtella, margine ciliolato-fimbriolata. *Pedunculus*  $\frac{1}{4}$  lin. longus, capillaris, rubello croceus. *Perigonii* folia intima conspicue longiora. *Sporangium* cylindraceo-oblongum, sursum paulisper angustatum, sub microscopio subtili sime scabriusculum. *Calyptra* angusto-longa, subobliquata, hinc fissilis, glabra, pallens, apiculo fuscescens. *Operculum* dimidii sporangii longitudine, subobliquatum, pallidum, a basi ad apicem obtusiusculum sensim attenuatum. *Peristomii* simplicis denticuli 16, albi, setaceo-subulati, denticulato-scabri; materie olim illos connectente passim sæpe persistente.

**HAB.** ad imos arborum truncos Carolinæ.

**MARGINATUM.** P. fasciculato-ramosissimum; ramulis erectis, attenuato-filiformibus : foliis ovali-acuminatis; utraque ora revoluta quasi marginatis; nervo prominulo : sporangio oblongo; operculo subulato.

**Obs.** Affine P. *subcapillato*. Parvulum, densissime cæspitosum; cauliculis ramulosissimis, confertim erectis. *Folia* cylindraceo-imbricata, subovalia, sensim et longiuscule acuminata; nervo medio et revolutis marginibus quasi prominulo-trinervia.

CRYPTOGAMIA. *MUSC.* 293

*Pedunculus* capillaris, 4-5 lin. longus, rubens : *sporangium* oblongum, sursum paulisper angustatum : *operculum* e latiore basi longe cuspidatum, sub-obliquatum : *peristomii* denticuli pallidi.

*HAB.* in Carolina.

POLYTRICHUM. HEDW.

*Calyptra* quasi villosa.

*Peristomii* duplicis dentes exteriores numerosi, brevissimi ; membrana interior indivisa, ambitu foraminulosa.

*Flos* masculus foemineusque terminalis.

\* *Apophysi* sub *Sporangio*.

PERIGONIALE. P. simplex, majusculum ; foliis serrulatis ; perigonalibus dissimilibus, elongatis, membranaceis, capillatim desinentibus : sporangio tetragono ; operculo orbiculato, centro breviter apiculato.

*OBS.* Affine *P. communi* majori.

*HAB.* in Carolina : etiam in Europa.

\*\* *Apophysi* nulla.

AMBIGUUM. P. subsimplex : foliis longo-subulatis, aculeato-serratis : sporangio subcernua, cylindraceo : operculo promisse acuteque cuspidato.

*OBS.* Affine *P. alpino* DILLENII : foliis basi non ita dilatatis.

*HAB.* in Canada.

294 CRYPTOGRAMIA. *MUSCI*.

**FASCICULATUM.** P. caulibus breviusculis, superne fasciculato-subramosis : foliis linearibus, acutis, rariuscule aculeato-serratis : pedunculo breviusculo : sporangio cylindraceo, recto; operculo cum brevi mucrone convexiusculo.

*Obs.* Affine *P. urnigero*. *Caules* vix unciales et breviores; aut simplices, aut ramulis brevissimis superne compositi. *Folia* siocitate recta, conferta; linearia, sensim acutissima; aculeato-serrata. *Pedunculus* subuncialis, terminalis : *sporangium* erectum, cylindraceum : *operculum* depresso-convexum, mucrone centrali breviusculo.

*HAB.* in Canada.

**CAPILLARE.** P. brevicaule, simplex : foliis paucioribus, laxiusculis, linearibus, aculeato-serratis : pedunculo longiusculo, capillari : sporangio erecto, ovoideo; operculo convexo, mucrone abrupto, gracili, longiusculo.

*Obs.* Statura *P. nani*. *Caules* sub3-lineares, basi nudi. *Folia* acuta quidem, sed non angustato-subulata. *Calyptra* longiuscule acuminata.

**PENNSYLVANICUM.** P. acaule : foliis paucissimis; in-  
*HAB.* ferne in oblongum dilatato-appressis; superne subulato-mucronatis, obscure denticulatis : sporangio erecto, oblongo-cylindraceo; operculo convexo, mucrone longiusculo, inclinato.

*HAB.* in monte *Grand-Father* Carolinæ.

CRYPTOGAMIA. *musci*. 295

BRACHYPHYLLUM. *P. acaule* : foliis ad radicem conglobatis, brevissimis, ovali-oblongis, apice crassiusculo obtusatis : sporangio subeorneo, oblongiuscule obovoideo; operculo breviter apiculato.

*Obs.* Statura *P. nani*. Folia vix 1 lin. longa; madore incurvulo-patentia. Pedunculus subuncialis et solito brevior. Sporangium maturum exsiccatione subclavato-oblongum.

*HAB.* in Carolina inferiore.

UNDULATUM : minus. *P. simplex* : foliis lanceolatis, *HEDW.* serratis, undulosis, laxe patentibus : sporangio oblonge cylindrico, subarcuatim obliquo : calyptra glabriuscula.

*HAB.* in Carolina.

TORTULA. *HEDW.*

*Peristomium simplex* : dentibus capillaribus, spiraliter convolutis.

*Flos masculus* gemmaceus in eodem individuo.

TORTUOSA. *T. altiuscule caulescens*, ramosa : foliis *HEDW. sp.* lineari-lanceolatis, acutis, subdenticulatis, crispatis : sporangio recto, gracili-cylindrico; operculo subulato.

*Obs.* Varietas paulo minor; pedunculis vix semiunciam superantibus.

*HAB.* in Carolina et Canada.



## TRICHOSTOMUM. HEDW.

*Peristomium simplex.* Dentes 16, e basi exigua  
in crura duo, filiformia, recta divisi.

*Flos masculus* alaris.

CANADENSE. T. prostratum, ramis assurgentibus:  
foliis lineari-lanceolatis, non margina-  
tis; acumine decolore, vix perceptibi-  
liter denticulato: pedunculo breviusculo:  
sporangio oblongo-ovato, operculo an-  
guste conoideo.

*Obs.* Affine *T. lanuginoso*; statura rectius quadrans  
cum DILL. t. 47. f. 28.

*Folia* integriuscula, acumine albido, mediocri.

*HAB.* in pinetis Canadæ.

PALLIDUM. T? subacaule, simplex: foliis setaceo-  
*HEDW.* subulatis: pedunculo capillari, pallido:  
sporangio oblonge cylindraceo-ovato;  
operculo conico, obtuso.

*Obs.* Variat pedunculis vix semiuncialibus.

*HAB.* in Carolina et Virginia.

DICRANUM. HEDW.

*Peristomium* simplex : dentibus 16 brevioribus, inflexis, bifidis.

*Flos masculus* capituliformis vel gemmiformis terminalis in distincto individuo.

SCOPARIUM. *D. maximum*, assurgenti-ramosum :

*HEDW.* foliis homomallis, lanceolato-subulatis, summitate serrulatis, lævibus : sporangio subarcuato, cylindraceo-oblongo ; operculo longissime subulato.

*HAB.* in Carolina.

POLYSETUM. *D. maximum*, assurgens, pauci-ra-

*SW.* mosum : foliis lanceolato-subulatis, superne subaculeato-serratis, disco unduloso-crispis : pedunculis ex singulo perigonio pluribus.

*OBS.* Affine priori : *Folia* sæpius undique versa, rarius tamen homomalla.

*HAB.* in Canada.

GLAUCUM : pumilum. *D. caulibus* ramosis, densis-

sima foliatione crassis : foliis glaucis, convexo-canaliculatis : sporangio subinclinato, oblongo-ovato ; operculo graciliter subulato, subinflexo.

*OBS.* Mea varietas humillima ; foliis acuminatis.

*HAB.* in Carolina.

ORTHOCARPUM. *D. pusillum*, brevi-caule, simplex:

*HEDW.* foliis setaceis, basi dilatatis, subheteromallis : pedunculo laterali, longiusculo : sporangio suberecto, oblongo-ovato ; operculo rostrato.

*Obs.* *Caules* 2-4-lin. alti. *Folia* dilute flavidulo-viridula, sæpe undique versa. *Pedunculus* 6-9 lin. longus, tenuiter capillaris, rectus, plerumque tortilis, pallens. *Sporangium* croceum, non ita angustum ac in icone *HEDWIGII*; *peristomio* saturate purpureo.

*HAB.* in arenosis Carolinæ.

PURPUREUM. *D. erecto-caulescens*, ramosum :

*HEDW.* foliis lineari-lanceolatis, carinato-canaliculatis : pedunculo laterali : sporangio ovato, arefactione sulcato ; operculo oblongiuscule conico, mutico.

*Obs.* Meum humile et statura *D. purpurascens*.

*HAB.* in Carolina.

## FISSIDENS. HEDW.

*Peristomium simplex* : dentibus sedecim, latiusculis, bifidis.

*Flos masculus* gemmaceus axillaris, in eadem planta.

SUBBASILARIS. F. cauliculis distichophyllis : pedunculo subbasilari : sporangio erecto, oblongo-ovato.

*OBS.* Pusillus, densis et indefinitis crescens cæspitibus.  
*Folia* lineari-oblonga, obtusiuscula. *Pedunculus* brevis.

*HAB.* in monte *Grand-Father* Carolinae.

SCIUROIDES : vaginatus. F. cauliculis simpliciuscule elongatis, siccitate curvis : foliis confertim imbricatis, lanceolatis, striatis : pedunculo laterali breviusculo, intra perigonium longum prorsus vaginato.

*OBS.* Varietatis istius foliatio uno versu subsquarrosa.

*HAB.* in Canada et in monte *Grand-Father* Carolinae.

## ORTHOTRICHUM. HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim, latiusculis : *interius* dentibus totidem, latiusculis, erosis; vel nullum. *Calyptra* conica, pilis erectis.

*Flos masculus* gemmaceus, axillaris et terminalis.

Obs. Vix recte distinctum a GRIMMIA.

STRIATUM. O. foliis lanceolatis, subacutis, muticis : sporangio ovoideo, inferne turbinate; arefactione clavato, conspicue sulcato-striato : peristomii utriusque denticulis pallidis.

HAB. in Canada et Carolina.

## BARTRAMIA. HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim, cuneiformibus : *interius* membranaceum, plicato-carinatum, apice lacinulatum.

*Flores masculi* foemineis approximati.

*Character emendatus* ex SWARTZ.

*Sporangium* subrotundum.

*Peristomium* duplex : *exterius* dentibus sedecim, apice inflexilibus : *interius* e membrana carinata, 16-partita, lacinulis bifidis.

VULGARIS. B. decumbenti - assurgens, pauciramosa : foliis subulato-setaceis, serrulatis;

CRYPTOGAMIA. *mscr.* 301

varie curvatilibus : pedunculo subterminali, longiusculo.

*B. pomiformis.* HEDW.

*Obs.* Omnium hujus generis specierum, tum Europæarum quum Americanarum *sporangium* sphæroidæum : ideoque mutandum mihi videtur nomen triviale, in methodo Linnæana bonum et in Hedwigiana malum.

*HAB.* in Carolina.

LONGISETA. *B. brevicaulis* ; cauliculis simpliciusculis : foliis non ita densis, subulatis, margine carinaque vix perceptibiliter denticulatis; siccitate erectis : pedunculo subbasilari, longissimo.

*Obs.* *Caules* sub3-unciales; inter folia non ita ac in præcedente conferta partim visibiles. *Pedunculi* ex imis cauliculis oriundi, erecti, non tortiles, subrubenti-crocei, 1  $\frac{1}{2}$ -2 unc. longi : *sporangium* uti in congeneribus crassum itidemque obliquatum : *peristomii externi* dentes angustiusculi et subcrocei; *interni* membrana subalbido-pallida.

*HAB.* in Carolina, ex herbario D. Bosc, et in Virginia ex D. BEAUVOIS.

BUXBAUMIA. *L.* HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim, truncatis : *interius* membranaceum, plicatum.

*Flos masculus* discoideus, terminalis.

FOLIOSA. *B. subacaulis* : foliis linearibus ; radi-  
*L. HEDW.* calibus muticis ; perigonalibus ærista-

tis : sporangio sessili, ventricoso-ovato,  
vertice obliquato.

*HAB.* in Carolina inferiore.

W E B E R A. HEDW.

*Peristomium duplex* : *exterius* dentibus sedecim,  
latiusculis, acutis : *interius* membranaceum :  
dentibus acuminatis.

*Flos hermaphroditus* terminalis.

**PYRIFORMIS.** W. cauliculis simplicibus ; peduncu-  
*HEDW.* liferis depressis ; rosuliferis elongatis :  
foliis alternis, capillaceis : sporangio cer-  
nuo, ob stipitem insignem oblongo-py-  
riformi ; operculo convexo, mutico.

*HAB.* in humidis Canadæ.

## FUNARIA. HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim, obliquis, apice coherentibus : *interius* ciliis totidem membranaceis, planis.

*Flores masculi foemineique terminales* : illi discoidei in diversa planta.

FLAVICANS. F. foliis longius setaceo-acuminatis : pedunculo capillari, pallido.

*Obs.* Statura *F. hygrometricae* : folia pallido-viridula, angustiora : pedunculo graciliore, constanter pallido-flavescente : sporangio minore, flavo.

*HAB.* in Carolina.

## MEESIA. HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim, brevibus, obtusis : *interius* ciliis totidem, acutis, distinctis vel opere reticulato connexis.

*Flores masculi foemineis approximati* vel discoidei in diversa planta.

LONGISETA. M. caulescens : foliis sensim angustato-acutis : superioribus e basi lata subulato-acuminatis : pedunculo longissimo : operculo convexo : peristomii interni, exteriore subquadruplo longioris, dentibus setaceis.

*Obs.* Folia lanceolata, sensim in acutum angustata ; superiora et perigonica inferne dilatato-amplexantia,



acumine longius subulato; arefactione involutitia. *Pedunculi* quasi laterales, ramulorum seu cauliculorum, cauliculis sterilibus multo demissorum, terminales, 3-4  $\frac{1}{2}$  unc. longi, rubentes. *Sporangium* crassum, arcuato-cernuum, elongato-subpyriforme; siccitate ovoideum et angustius stipitatum. *Operculum* subconoideo convexum, muticum. *Peristomii* utriusque dentes pallidi; *exterioris* brevissimi; *interioris* gracili-setacei, scabriusculi.

*HAB.* in paludosis Canadæ.

### T I M M I A. Hedw.

*Peristomium* duplex: *exterioris* dentibus sedecim, latiusculis: *interius* obsolete sulcatum, laciniatum; laciniis subæqualibus, pertusis, unifornibus.

*Flores masculi* monoici, gemmacei, pedunculati, axillares: *feminei* terminales.

**CUCULLATA.** T. foliis linearibus, angustato-acutis, serrulatis: pedunculo breviusculo: sporangio cernuo, arcuato, obconice turbinato, e latere persistentis calyptræ erumpente: peristomii externi dentibus articulatis.

*Obs.* *Caules* erecti, subbiunciales, pauciter longiusculeque ramosi. *Folia* non ita crebra, Polytrichoides; basibus paulum dilatatis amplexanti-incumbentia; cætero patula; nervo totali, validissimo. *Pedunculus* vix uncialis, sæpe cauliculos non superans; setaceus, rectus, non tortilis, rufus. *Sporangium* subito

## CRYPTOGAMIA, *musci*. 305

propria recurvatione cernuum. *Calyptra* angusta, erecta persistit, summitem pedunculi basimque sporangii fissura laterali erupti inferne involvens. *Operculum* jam excussum non vidi. *Peristomii exterioris* dentes validi, fere subulati, sensim in acutissimum angustati; supra basim conspicue articulati; infra articulationem obscure flavi, supra pallidi; superiore ista parte vetustate sæpe separabili a *interioris*, ejusdem longitudinis pallentisque lacinulæ capillares, elegantissime varieque in partialia reticula connexæ.

*HAB.* in humosis udis Canada.

### M N I U M. HEDW.

*Peristomium* duplex: *exterius* dentibus sedecim, latiusculis, acutis; *interius* membranaceum, plicato-carinatum, laciniatum, laciniis latiusculis capillaribusque alternis.

*Flos masculus* terminalis, discoideus.

PALUSTRE. *M. maximum*: caulibus elongatis, erectis, tomento radiculario insigniter squalentibus: foliis confertis, lanceolatis, brevi-acuminatis, integriusculis: sporangio erectiusculo, oblongo-ovoideo; operculo conico.

*Obs.* Pariter ac in Europæo, folia interdum versus summitem minutissime denticulata.

*HAB.* in paludibus Carolina.

366 CRYPTOGRAMIA. MUSCI.

CUSPIDATUM. M. caulibus sporangiferis erectis ,  
*HEDW.* simpliciusculis : foliis subalternis , lato-  
ovalibus , abrupte acuminatis , ciliato-  
denticulatis : sporangio reflexo , oblongo-  
ovato ; operculo demisse conoideo , sub-  
acuto.

DILL. t. 53. f. 79.

*HAB.* in Carolina et Canada.

ROSACEUM. M. caulibus simplicibus ; summitate  
*HEDW.* rosaceo-foliosis , cætero nudis : foliis  
ovali-oblongis , cuspidatis , insigniter  
aculeolato-serratis : sporangiis reflexis.

DILL. t. 52. f. 77. Meum autem foliis copiosius  
densiusque aggregatis etiamque angustioribus.

*HAB.* in Canada et in monte *Grand-Father* Caro-  
linæ.

## NECKERA. Hedw.

*Peristomium* duplex : *exterius* dentibus sedecim, acutis : *interius* ciliis sedecim, distinctis, cum dentibus alternantibus.

*Flores masculi* gemmacei in distincta planta.

PENNATA. N. foliatione distiche compressa : foliis ovali-lanceolatis, acuminatis, nerviis, disco undulatilibus : sporangio intra perigonium immerso ; operculo conoideo, modice cuspidato.

Obs. Mea omnino Hedwigianæ conformis, sequentibus tamen notis aliquantulum discrepans. *Calyptra* sub ipsa Perigonii dehiscentia sporangium immaturum deserens ; superna parte operculum includens, glabra ; inferna rariter breviterque subpilosa. *Peristomium* simplicissimum, exalbido-pallens ; dentibus 16, æqualibus, angusto-subulatis, interdum nonnullis superie geminatim connexis ; interjectis minoribus certo nullis. Nonnunquam etiam vidi dentes omnes, jam excusso operculo, summitatibus connexos.

An ista, una cum nonnullis affinibus, proprium Genus constituere debet !

HAB. in truncis arborum Canadæ et Carolinæ superioris.

CLADORRHIZANS. N. repens, patule et subdistiche  
Hedw. ramulosa ; summitatibus passim radican-  
tibus : foliis subdepresso-imbricatis, ova-

libus, acutis, enerviis : sporangio cylindraceo ; operculo oblonge obtuseque conoideo ; columella exerta.

*Obs.* *Folia* solito nitidule flavicantia etiamque dorso (omissis marginibus et summitate) fulvescentia ; concavula, sensim angustato-acuta, integra. *Perigonium* longiusculum : foliis erectis, interioribus lanceolatis. *Pedunculus* directione ramulorum, rectus, demum insigniter tortilis, 8-10 lin. longus, validus, sanguineus. *Sporangium* erectum, cylindraceo-oblongum, sursum tantisper angustatum : rufescens : *Peristomii externi* denticuli angusto-subulati, purpurascens, summitatem versus flavescentes ; *interni*, longitudine aequalis, cilia pallentia : *Columella* supra *Peristomii* dentes exerta.

*HAB.* in arborum truncis Carolinae.

**SEDUCTRIX.** *N. repenti-subassurgens* ; ramulis erectis, passim fasciculatis : foliis arcte cylindraceo-imbricatis, ovalibus, acuminulatis, enerviis : sporangio cylindraceo ; operculo e basi conica breviuscule cuspidato.

*Obs.* Præcedente, cui affinis, paulo minor. *Folia* absque nitore viridula seu pallescentia ; ramorum inferiora obtusiora. *Pedunculus* vix semiuncialis, saturatius coloratus. *Differentia* a *Peristomio* desumpta fallax.

*HAB.* in Carolina, terrestris et ad imos truncos.

**VITICULOSA** : minor. *N. cauliculis assurgentibus*, simplicibus pauciterque ramosis : foliis

CRYPTOGAMIA. *musci* 309

laxiusculis, e latiore basi ligulato-obtusis: sporangio oblongo; operculo conico, longiuscule cuspidato.

*Obs.* Europæa multo minor nec ita viticulosa, multoties gracilior; adeo ut, primo obtutu, diversa videatur.

*HAB.* in truncis arborum Carolina.

LESKEA. Hedw.

*Peristomium* duplex: *exterius* dentibus sedecim, acutis: *interius* membranaceum, partitum in lacinias æquales.

*Flores masculi* gemmacei in distinctis individuis, axillares.

DENDROIDES. L? caulibus dendroideis: foliis con-  
*Hedw.* fertis, erectis, lanceolatis; subtriner-  
viis, superne serrulatis: sporangio erecto,  
longe cylindraceo; peristomio interno  
demum capillatim dissecto.

*Obs.* *Mœæ pedunculi* breviores, 8-9 lin. longi. *Sporangia* angusto-cylindracea, basi in acutum angustata. *Peristomii exterioris* dentes angusto-subulati, purpurei: *internum* pallens, primum superne 16-fidum, deinde fere ad basim fissum; laciniis demum in 2 seu 3 capillos divisibilibus. *Columella* promissa exserta. Vix hujus generis!

*HAB.* in sylvis Carolina.

310 CRYPTOGAMIA. MUSCI.

ROSTRATA. *L. densissime cæspitosa*, filiformi-ramosa : *HEDW.* foliis cylindrice imbricatis, ovalilanceolatis, subcapillaceo-acuminatis : sporangio erecto, ovato; operculo e basi conica graciliter cuspidato.

*Obs.* Cæspitibus confertissimis, olivaceis seu flavidis crescit, copiosissime erecteque ramulosa; ramulis tereti-filiformibus, superne solito attenuatis. *Folia* minuta, sensim gracillimeque angustato-acuminata; acumine fere piliformi; marginibus asperiusculis, siccitate subrevolutis; nervo prominulo. *Perigonium* longiusculum, pallens; foliis setaceo-subulatis. *Pedunculus* supra cæspitem emergens, 4-5 lin. altus, purpureus, capillaceus. *Sporangium* oblongiuscule ovatum, rufidulum : *operculum* dimidiam illius longitudinem paulum superans. *Calyptra* angusto-longa, hinc fissilis, glabra. *Peristomium* brevissimum : *exterioris* dentes subulati, inferne lutescentes, superne albescentes et angustati : *interioris* membrana alba, profunde 16 partita; lacinulis brevius subulatis, a se invicem distantibus.

*HAB.* terrestria in montibus Carolinæ.

ADNATA. *L. pusilla*; surculis brevibusque ramulis radicato-adnatis : foliis patulis, lanceolatis, enerviis : sporangio erecto, ovato; operculo e basi convexa gracillime longissimeque cuspidato.

*Obs.* Affinis *L. cæspitosæ*. *Caules* brevissimique ramuli vage repentis, quasi cortici arborum adnati. *Folia* flavescentia, veluti laxè imbricata; superne, sensim tenuiterque angustata, integerrima, cana-

Ecclata; siccitate margines paulisper involventia, et ad summities solito fasciculatim incurvo-adscentia. *Pedunculus* capillaris, 3-4 lin. longus, rubellus, *Perigonii* folia vix a caulinis dissimilia. *Sporangium* rufescens; *operculum* fere hujus longitudine. *Peristomii exterioris* dentes ex aurantiaco-lutescentes: *interioris* cilia angustissime subulata, breviter connexa, exalbentia.

Datur varietas omnibus partibus multo gracilior.

*HAB.* in *Quercu aquatica* Carolinae inferioris.

RECURVANS. *L. intricato-repens*: foliatiōe subdistiche depressa, postice squarrosa; foliis ovalibus, angustato-acuminatis, enerviis: sporangio recurvo; operculo e basi convexo-conoidea oblique cuspidato.

*Obs.* Statura *L. variae*. Densis caespitibus intricatim repens; ramis decumbentibus, subdistiche ramalosis. Foliatio quasi velutino-lucidula; madore, praesertim ad utrumque ramorum latus, patula, et ad posticam eorumdem faciem, modice recurvatis foliis, subsquarrosa; siccitate quasi laxiuscule imbricata et manifestius squarrosa: *Folia* ovalia, in acumen longiusculum angustata, concavula, enervia, integra; summitate tamen (armato oculo) plerumque denticulata. *Perigonii* breviusculi folia lanceolata, acumine patentiusculo, caetero caulinis similia.

*Pedunculi* ex caespitibus copiose assurgentes, croceo-sanguinei, capillares, 6-8 lin. longi. *Sporangium* ab ipsius pedunculi summitate recurvo-cernuum, subovoideo-oblongiusculum, rufum. *Operculum*



paulo plus quam tertiam sporangii longitudinis partem constituens; cuspidē, modo sursum, modo deorsum, plus minus inclinata. *Peristomii* dentes lutei seu etiam subcrocei : membrana interior mediocriter 16-fida.

*HAB.* in montibus Carolinæ, ad imos truncos.

**SQUARROSA.** L. repens : foliatione subdistiche depressa, laxē imbricata, postice insigniter uncinato-squarrosa ; foliis lanceolatis, gracillime angustato-acuminatis ; perigonii longiusculi lineari-subulatis : sporangio propria arcuatione cernuo.

*OBS.* Valde affinis præcedenti. In posterum, cognito operculo (a me non viso) rectius distinguenda. *Folia* enervia, summitatē subdenticulata, solito stramineo-flaventia. *Pedunculi* subunciales, madore rutili. *Peristomium* præcedentis.

*HAB.* in humosis humidis Carolinæ.

## ARRHENOPTERUM. HEDW.

*Peristomium* BRYT.*Masculus flos alaris* : *femineus terminans*.

HETEROSTICHUM. A. erectiusculum : foliis laxiusculè imbricatis, ovalibus, cum apiculo obtusis, serrulatis; nervo totali : pedunculo plerumque terminali : sporangio oblongo, arcuato; operculo conico, obtuse acuminato.

Oss. *Caules* indefinite erecto-cæspitosi, subunciales.

*Folia* majuscula, concavula, pellucida, ovalia, cum exilissimo acumine obtusa, superne manifeste serrulata. *Pedunculus* plerumque terminalis; interdum, factis prope summitates innovationibus,

quasi lateralis, 6-9 lin. longus, dilute purpureus.

*Perigonii* folia multo breviora; exteriora sublanceo-lata; interiora pauciora, obtusa, aristato-cuspidata: setulae nonnullae breviusculae intimum occupantes.

*Peristomii exterioris* dentes dilate lutescentes, latiuscule subulati : *interioris* membrana exalbida, laciniata; lacinulis capillisque numero proportionaque variantibus.

HAB. in putridis humosis Carolinis.

## HYPNUM. L. HEDW.

*Peristomium* duplex : *exterius* dentibus sedecim ,  
*latiusculis* : *interius* membranaceum , æquale ,  
 laciniatum : laciniis latiusculis cum interjectis  
 capillaribus.

*Flores masculi* gemmacei in diversis plantis.

CRISTA CASTRENSIS. H. caulibus elongato - simpli-  
 L. HEDW. ciusculis , recurvatim concinneque pen-  
 nato-ramulosis : foliis uno versu cir-  
 rhato-recurvis , subulatis , tenuiter ner-  
 voso-lineatis : sporangio longissime pe-  
 dunculato , oblongo , arcuato ; operculo  
 conico.

OBS. *Caules* 3-6 unciales , incumbenti-procumbentes ,  
 sæpe simplices. *Folia* fere piliformiter desinentia.  
*Perigonii* longi folia lineari-lanceolata , erecto-ap-  
 pressa , conspicue striata. *Pedunculi* sæpius subaggre-  
 gato-approximati ,  $1 \frac{1}{2}$  —  $2 \frac{1}{2}$  uncias longi , firmiter  
 setacei , sanguinei. *Sporangium* crassum , cernuum ,  
 cylindraceo-oblongum , manifeste arcuatum , ru-  
 fum : *Operculum* conicum , apiculatum. *Peristomia*  
*externi* dentes aurantiaci ; *internum* luteolum.

HAB. in humosis Canadæ et Novæ Angliæ.

MOLLUSCUM. H. subpennatim confertimque ra-  
 HEDW. mulosum : foliis sublineatis ramulisque  
 insigniter recurvo-crispis : pedunculo  
 breviusculo : sporangio cernuo , turgide

CRYPTOGAMIA. *MUSCI*. 315

ovato ; operculo conico , acutissime apiculato.

*HAB.* in sylvis Canadæ.

**IMPONENS.** *H. brevissime subpennato-ramulosum*,

*HEDW.* recurvatum : foliis uno versu uncinatis , enerviis : sporangio erecto , cylindraceo-oblongo ; operculo conico , obtuse apiculato.

*OBS.* Statura vetuli *H. mollusci* , caulibus solito brevioribus : ramuli , saltem arefactione , insigniter uncinato-recurvi. Foliatio nitide subfulvo-flava , foliis densis , lanceolatis , graciliter angustatis , integris , ad apicem ( sub microscopio ) obsolete rariterque subdenticulatis. *Pedunculus* mediocris , capillaris. *Sporangium* æquilateraliter erectum ! *Peristomii exterioris* dentes pallido - lutescentes ; *interioris* membrana ciliolaque exalbida.

*HAB.* ad imos truncos Carolinæ montosæ.

**REPTILE.** *H. minutum* , adnato-repens , pinnato-ramulosum : foliis dense imbricatis , uno versu recurvatis , subtilissime denticulatis , nervo tenui subcarinatis : sporangio erectiuscule subarcuato , cylindraceo-oblongo : operculo convexo , breviuscule acuminato.

*OBS.* Valde affine *H. cupressiformi* ; quo multoties omnibus partibus , omisso tamen sporangio , minor. *Caulēs* arcte cortici arborum radicato-adnati : ramuli brevissimi. *Folia* quasi lanceolata , sensim in

acutissimum angustata, subcarinato-canaliculata.

*Sporangium* veluti in *H. cupressiformi*.

*HAB.* in monte *Grand-Father* Carolinae.

TRIQUETRUM. *H. maximum*, erectiusculum : ra-

*L.* mulis subdistichis, recurvatis : foliis patulis, quasi triangulatim ovali-lanceolatis, integris, lineatis : sporangio recurvato, crassius ovoideo ; operculo convexo, minutim acuteque apiculato,

*HAB.* in Canada.

ABDETINUM. HYPNUM.

*L.*

Nº. Fructificans reperire nequivi.

*HAB.* in Canada.

MINUTULUM. *H. gracillimum*, decumbens, pin-

*HEDW.* natim ramulosum : foliis minutissimis, ovalibus, acutis ; marginibus carinaque scabriusculis ; siccitate seorsim incurvis : sporangio recurvo ; operculo longe rostrato.

*OBS.* Habitu *H. delicatulo* quodammodo affine, multo minus. *Caules ramulique* fere capillari tenuitate. *Folia* perexigua, saturate viridia, absque densitate approximata ; concava ; brevi-ovalia : madore, assurgenti-patentiuscula ; arefactione, ita singulatim incurvata, ut cauliculos (sub lente vitrea communi) veluti granulosa efficiant. *Perigonii* conspicui et flaviduli folia appresso-erecta, in acumen sensim

CRYPTOGAMIA. MUSCI. 319

angustata ; exteriora lanceolata ; interiora linearia. *Pedunculus* (ratione plantulae) longus, 9-12 lin. altus, capillaris, purpureus. *Sporangium* recurvum, subturbinato-oblongum ; *operculum* fere illius longitudine, e basi conoideo-convexa rostellatum ; rostellum longiore, gracillimo, solito curvo. *Peristomii* dentes crocei ; inferne aurantiaci ; membrana ciliolaque exalbido-pallentia.

Var. *α. muticum* : foliis simpliciter subacutis.

— *β. acuminatum* : paulo majus, foliis angustoa-cuminatis.

Hoc posterius affinius *H. delicatulo* Hedwien.

*HAB.* terrestre in Carolina.

PROTENSUM. *H.* longius rependo-protensum, brevissime bipinnato-ramosum : foliis exiguis, lanceolato-ovalibus, marginibus dorsoque muricatis : pedunculo longissimo : sporangio oblongo, arcuato ; operculo breviuscule rostrato.

*Obs.* Valde affine *H. tamariscino*, paulo gracilius ; brevitate ramorum et conspicua pedunculorum longitudine primo intuitu distinguendum. *Folia* concava, ovalia, sensim angustato-acuta, manifesto scabra. *Pedunculus* valide setaceus, subbiuncialis. *Sporangium* ejusque partes *H. tamariscini* ; *peristomii* dentibus croceis seu subaurantiacis.

*HAB.* in sylvis Canadae et Carolinae.

SPLENDENS : interruptum. *H.* caulibus elongatis ; ramis complanato-bipinnatis, interrupte con-

*LICHENACEÆ.*

*PATELLARIA.* Ach.

*Crusta* solida contigua et æquabilis aut rimosa areolata etiam granulata, suborbicularis vel irregulariter expansa.

*Receptacula* : *scutellae* sessiles, superficiales, concavæ, planæ et convexiusculæ marginatæ, vel etiam hemisphæricæ et subglobosæ immarginatæ seu margine tenui et vix manifesto ad basin cinctæ.

GRANULOSA. P. crusta indefinita, cinerascens, tota granuloso-pulverulenta : scutellis aterrimis, planiusculis, marginulatis.

*Oss.* Crusta non limitata, obsolete cinerascens aut subterrea; *scutellae* mere aterrimæ, rariusculæ, orbiculatæ, marginulo obtusæ; interdum passim confluentes in glomerulos convexos.

*HAB.* in lignis putrescentibus Carolinæ montosæ.

CRYPTOGAMIA. LICHENACEÆ. 321.

PSOROMA: Ach.

*Crusta* solida irregularis effigurata, foliolis leprosis distinctis squamatim incumbentibus composita aut tecta.

*Receptacula* : *Scutellae* sparsae marginatae planae, demum convexae.

**PALMULATA.** P. crusta foliaceo-frondiculosa, subtereti-laciniosa; lacinulis passim subpalmatis: scutellis concavis, castaneis, crustae laciniato-marginanti immersis.

*Obs.* Crusta obsolete olivacea; quasi e condensatis lacinulis inaequalibus, obtusiusculis, intervalla scutellartum occupantibus easque coronatim marginantibus: ad ambitum sterilem solito excurrit in lacinias laminatas, varie ramosas et dentatas, frondiculas *PHYSCIAE tenellae* veluti mentientes. *Scutellae* concavo-depressae, majusculae, orbiculatae aut saepius pressione difformes.

Propter quamdam habitus affinitatem cum *P. microphylla*, ad hoc genus refero; forsitan potius inter *PLACODIA* recensenda.

*HAB.* in arboribus montis *Grand-Father* Carolinae.



322 CRYPTOGRAMIA. LICHENACEÆ.

IMBRICARIA. Ach.

*Folia* membranacea (rarius subcrustacea fragiliora) plano-depressa imbricatim centrifuga, rosulam formantia, laciniato-incisa atque pinata, lobata; subtus fibrillosa.

*Receptacula* : *Scutellae* primo urceolatæ seu concavæ, dein planiusculæ submembranaceæ, centro tantum suo foliis affixæ, circumcirca liberæ, elevatæ, marginatæ. *Glomeruli* in quibusdam sparsi, centrales et marginales pulverulenti.

CONVEXIUSCULA. I. subtus glabello-atra, supra lucidule subolivacea; laciniis convexis, sinuato-lobatis; extremitatibus dilatatis: scutellis castaneo-fuscis, extus frondi concoloribus.

*Oss.* Species IMBRICARIIS *omphalodi* et *physodi* intermedia.

*HAB.* in saxis Canadæ.

UMBILICARIA. ACH.

*Folia* rigida cartilagineo-membranacea, peltata, rotundata, depressa integra laciniata; subtus nuda aut fibrillosa.

*Receptacula*: *Tricae* sessiles planiusculæ et convexæ immo globosæ, rotundatæ et angulatæ; plicis concentricis et gyrosis, rugosis et glabris ornatæ, semper atræ.

VELLEA. U. amplissima, simplex, concava, ACH. ambitu modice lobata, squalide cinerea, subtus atro-pannosa: tricis atris, disco demum contortuplicatim extuberante.

DILL. t. 82. f. 5. quadrat.

Obs. Amplitudinem interdum pedalem assequens; concavo-peltata, superficie lævis; crassina coriacea; villo subtus densissimo, brevi, composito. *Tricae* atræ, primum discoideo-orbiculatæ, ambitu annulatæ; demum protuberantiis disci varie flexuosis et lineæ exaratis compositæ.

HAB. in saxis Canadae.

PUSTULATA. U. supra cineritia, bullato-pustulosa; subtus subolivaceo-fuliginosa, profunde lacunosa; superficie pulverulento-scaberula: tricis planis, demum gyrosis.

HAB. in saxis Canadae.

## LOBARIA. AGH.

*Folia* membranaceo-succoriacea, lato-lobata, et rotundata, expanso-adsocendentia, vaga, laxa, subtus villosa.

*Receptacula*: *Scutellae* sparsae subsessiles, planiusculae pateriformes et concavae.

PERFORATA. *L. grisea* seu subfulva, laevis; sub-  
 ACH. tus nigrescens; laciniis undulosis, obtuse lobulatis, ambitu nigro-crinitis; scutellis castaneo-fuscis, extus frondis concoloribus laevibusque, centro perforatis, fractura croceo-miniata.

Obs. Statura fere IMBR. *acetabuli*. Supra exalbido-cinereascens aut obsolete fulvescens, superficie laevis; subtus nigricans et subnuda, extremitatibus liberis in varietate *grisea* griseolis et in *fulva* croceis: laciniis quasi lobulato-crenatis, ambitu conspicue crinitis. *Scutellae* majusculae, prominentes, insigniter concavae, margine simpliciter, sub fractura, croceo-miniata.

HAB. in arboribus Canadae et Carolinae.

QUERCIZANS. *L. tenuiuscule* membranacea, laevis; subtus rufa; laciniis oblongis, sinuoso-lobulatis: scutellis parvulis, crebris, subadnatis, concavulo-planiusculis, fuligineo-fuscis.

Obs. Affinis *L. herbaceae*; paulo major crassiorque: supra solito obsolete rufescens laevisque. Laciniis

**CRYPTOGAMIA. ECHINACEAE. 325**

quodammodo, tum colore quam lobatione, folia  
Quercus emortua simulantes.

Datur varietas sterilis, marginibus pannoso-  
crispis.

*HAB.* arboricola in monte *Grand-Peak* Carolinae.

**SUBMARGINALIS.** L. tenui-membranacea; supra  
cinerascens laeviusculaque; subtus ater-  
rima, versus summitates fuscescens: scu-  
tellis submarginalibus, concavis, intus  
fuscis, extus rugelosis, margine sim-  
plici.

*Obs.* Statura *E. herbaceae*: supra (armato oculo) punc-  
ticulis pulveraceis nigris conspersis.

*HAB.* arboricola in Canada et Carolina.

**PULMONARIA.** *H.* subcartilaginea; divaricato-lo-  
*ACH.* bata, sinibus rotundato-obtusis; supra  
conspicue reticulato-lacunosa; subtus  
flavidulo-tomentosa: scutellis submargi-  
nalibus, spadiceis.

*HAB.* arboricola in montibus Carolinae.

## PHYSICIA. АСН.

*Folia* membranacea, vel pro maxima parte cartilaginea, glabra, rigida, angustiora, profunde et quasi ramoso-laciniata, interdum paululum latiora et canaliculata, lobato-crispa, caespitosa, erecta et adscendentia.

*Receptacula*; *Scutellae* sparsae et terminales concavae ac plano-convexiusculae.

*Glomeruli* in quibusdam laterales marginalesque, nec non foveolae in axis farinosae passim superficiales.

FRAXINEA. P. utrinque cinereo-concolor, rugoso-lacunosa; laciniis elongatis; scutellis lateralibus marginalibusque frondi subconcoloribus.

Она. Мел. varietas subuncialis, seu paulo ultra.

Над. in arboribus Canadae.

ISLANDICA. P. erecto-caespitosa, subrufa; frondibus membranaceis, sublineari-ramosissimis, laciniatis, breviter rigiduleque ciliatis; scutellis sessilibus, planiusculis, castaneo-rufis, margine subspinosulis.

Над. in monte Grand-Father Carolinae.

LEUCOMELOS. P. intricatim caespitosa, erecta, exalbida, furcatim lineari-ramosa, marginibus proluxe nigro-crinita: scutellis

CRYPTOGAMIA. LICHENACEÆ. 327

lateralibus, stipitatis, margine plano-  
dentatis.

*Oss.* Frondes angusto-lineares, subdichotome ramosæ; modo nivæ, modo sordide exalbantes; interdum passim subcroceo imbutæ; crinities marginalis in mea multo prelixior quam in icone Swartziana. *Scutellæ* rariores, stipite compresso, colore variabiles, extus sæpe corneæ: margo dentibus planis, nec spinuloso-radiatus.

*HAB.* arboricola in monte *Grand-Father* Carolinæ.

*EXILIS.* P. pusilla, sordide exalbida, divaricatim intricatimque setaceo-ramosissima: scutellis minutis, lateralibus, planiusculis, nitide croceis, margine simplici.

*Oss.* Minuta pulchellaque species, *P. chrysophthalma* (cui affinis) delicatior; intricatim cæspitosa, erectiuscula, compressiuscule setacea, veluti instar *CLADONIÆ pusillæ* ramosissima, ramulis gracillimo capillaceo-attenuatis. *Scutellæ* subsessiles, quasi lenticuliformes, intus crocæ, vix manifeste marginulata.

*HAB.* in arboribus Carolinæ; prius a clariss. Rosc detecta.

## SCYPHOPHORUS. ACH.

*Folia* cartilagineo-membranacea rigida fragilia subimbricata, adscendentia et erecta, flexuosa inaequaliter et profunde ut plurimum laciniata sinuataque, vel etiam sublobato-crenata; supra virescentia subtus albida; *bacilla* tubulosa simplicia (per aetatem aliquando ramosa) superne incrassata dilatata scyphiformia clausa, disco concavo, producentia.

*Receptacula*: *Tubercula* convexa et subglobosa fungiformia, ad margines scyphorum, (rarius in ipsis foliis) subbasalia, (inque bacillorum apicibus ramosis) terminalia.

SULPHURINUS. S. bacillis crassis, submembranaceis, subclavato-elongatis, laevibus, sulphurinis; summitate demum difformiter subdivisis et rimoso-pertusis: scyphorum tuberculis confluentibus, atrofuscis.

*Obs.* Color incano-sulphureus. *Bacilla* simplicia; primum apice simplicissime parvulaque scyphata; demum 2-3-uncialia, sursum sensim crassescencia, ad summitatem difformiter et veluti laciniatim subramosa; *scyphis* fructificantibus parvulis, margine confluenti-tuberculiferis. Interdum Bacilla e latere seu margine scyphi prolifera.

*HAB.* in Canada.

VERTICILLARIS. S. scyphis bacillorum, continuata innovatione proliferorum, verticillifor-

## CRYPTOGAMIA. LYCHENACEAE. 329

mibus, frondiculiferis et margine minutissime tubereulosis; tuberculis fuliginosis.

*Obs.* Species elegans: *Bacilla* crassiuscula filiformia, sordide subincana, simpliciuscula sive inferne subdichotoma, recta,  $2\frac{1}{2}$  ad 3-unc. alta. *Scyphi* centro prolifici, numero 6-7, subaequali distantia, in singulis bacillis veluti totidem conficiunt verticillos; margine extus tuberculiferi intusque et e disco producentes frondiculis varie crenato-lobulatas et radicalibus consimiles: ita ut singula bacillorum innovatio suam, pariter ac bacillum primordiale, habeat frondem basilem.

*HAB.* in Carolina.

## HELOPODIUM. ACH.

*Folia* cartilaginea rigida minuta subimbricata erectiuscula emutato-crenata supra virescentia subtus pallidiora. *Bacillae* subsolidae simpliciusculae, asperae parum dilatatae vel etiam subdivisae apice tuberculiferae.

*Receptacula*: *Tubercula* in bacillis terminalia fungiformia majora, simplicia, glomerata et aggregata, margine subinflexa.

**CAPITATUM.** H. bacillis discretis, simplicibus, obsolete substramineis: tuberculis capitatum subgloboso-concretescentibus, saturate rubiginoso-rufis.

*Obs.* Rarissime apex bacillorum breviter divisus, divisionibus etiam subcompositis tuberculiferis: tuber-



### 330 CRYPTOGAMIA. LICHENACEÆ.

...cepta vero solitarie discreta non vidi. Frondiculæ radicæ  
...dicales supra sordide subolivaceæ, subtus albæ;  
...imam bacillorum partem solito etiam occupantes.

*HAB.* terrestre in Carolina.

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

**UNCIALIS.** C. bacillis exalbido-pallidis, conspi-  
cuis fistulosis; axillis perforatis: ramulis  
brevissimis, divergentibus, rectis; ste-  
rilibus acutis; fertilium tuberculis minu-

...tis, fuscis.

*Obs.* Mea est varietas major, DILL. t. 16. f. 21.

*HAB.* in Canada.

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

...cardini ...

**RANGIFERINA.** C. bacillis aphyllis, exalbido-cine-

...tis, fuscis.

*ACH.* rasantibus, fistulosis, caespitoso-ramo-

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

...tis, fuscis.

# CRYPTOGAMIA. LICHEN-~~APH~~ 331

quasi fasciculatis; tuberculis exiguis, hemisphaericis, atro-fuscis.

*Obs.* Varietates duæ : 1<sup>a</sup>. *major*, cinerascens aut rarius pallescenti-incana : 2<sup>a</sup>. *minor*, capillaceo-ramulosissima, ramulis ita dense intricatis ut cæspes veluti crinitum crispam referat.

*Hab.* 1<sup>a</sup>. in Canada ; 2<sup>a</sup>. in Carolina.

## STEREOCAULON. ACH.

*Caules* erecti, solidi, liguescenti rigidi, saepe inæquales et acutijugati, vel doves, ramosi, raris inordinatis affixis. *Receptacula* : *Scutellae* nigrae, planae, vel etiam rotundatae, hemisphaericae et subglobosae sessiles, sparsae et terminales.

*PASCHALE*. S. bacillis vage ramosis, granulosa ramulorum matris confertim obsitis; scutellis sparsis, convexis, plerumque compositis, fusco-atris.

*Hab.* in Canada.

## SETARIA. ACH.

*Caules* : *Lora* cartilaginea filiformia teretiuscula nuda glabra, diffusa et pendula, ramosa, intus tela bombycina facta.

*Receptacula* : *Scutellae* planiusculae sessiles, demum convexae, sparsae, margine integræ.

*Glomeruli* sparsi pulverulenti.

*TRICHODES*. S. tenuissime capilliformis, proluxa,

332 CRYPTOGRAMIA. LICHENACEAE.

fuliginoso-atra, laevigata: scutellis coloribus, peltatis, convexis.

HAB. in lignis putrescentibus Canada.

U. S. N. E. A. AGH.

**Caules:** Lora cortice crustacea obducta, filo centrali sive medullari hyalino elastico, subarticulata scabriuscula teretia, ad basim crassiora, sensim attenuata fibrillosa.

**Receptacula:** Scutellus jam spinosus saevius marginē integro, demum convexus fangosae inaequales et saturatius coloratae, jam plani radiatae majores subterminales. Glomeruli pulverulenti sparsi.

FLORIDA. U. erectiuscula caespitosa, ramuloso-hispida: scutellis subterminalibus, amplicis, planiusculis, radiatis.

**Obs.** Tres mihi occurrere varietates:

1<sup>a</sup>. *major*; scutellis luteolo-pallidis.

2<sup>a</sup>. *intermedia*; minor, divaricator; scutellis minoribus, cineritiis.

3<sup>a</sup>. *rubiginea*; minus hirta, rubiginoso-rubens; scutellis concoloribus.

HAB. in Carolina, var. 3<sup>a</sup> in Canada.

# INDEX GENERUM.

Romanus numerus partem ; alter paginam indicat.

## A.

|                   |          |                    |     |
|-------------------|----------|--------------------|-----|
| ABIES, II.        | pag. 206 | Anthoxanthum, I.   | 39  |
| Acalypha, II.     | 215      | Antirrhinum, II.   | 20  |
| Acer, II.         | 252      | Anychia, I.        | 112 |
| Acrida, II.       | 253      | Apocynum, I.       | 121 |
| Aconitum, I.      | 315      | Aquilegia, I.      | 316 |
| Acorus, I.        | 194      | Arabis, II.        | 31  |
| Acróstichum, II.  | 272      | Aralia, I.         | 185 |
| Actæa, I.         | 308      | Arbutus, I.        | 249 |
| Adelia, II.       | 223      | Arenaria, I.       | 273 |
| Adiantum, II.     | 263      | Arethusa, II.      | 159 |
| Aeschynomene, II. | 74       | Aristida, I.       | 41  |
| Aesculus, I.      | 219      | Aristolochia, II.  | 161 |
| Agave, I.         | 187      | Arrhenopterum, II. | 53  |
| Agrimonia, I.     | 287      | Artemisia, II.     | 128 |
| Agrostis, I.      | 52       | Arum, II.          | 188 |
| Aira, I.          | 61       | Arundinaria, I.    | 73  |
| Aletris, I.       | 189      | Arundo, I.         | 73  |
| Alisma, I.        | 218      | Asarum, I.         | 279 |
| Alkonia, I.       | 100      | Asclepias, I.      | 114 |
| Alhum, I.         | 194      | Ascyrum, II.       | 77  |
| Alôpecurus, I.    | 43       | Asplenium, II.     | 264 |
| Amaryllis, I.     | 187      | Aster, II.         | 108 |
| Ambrosia, II.     | 182      | Astragalus, II.    | 66  |
| Ammannia, I.      | 99       | Avena, I.          | 72  |
| Ammi, I.          | 164      | Azalea, I.         | 150 |
| Amorpha, II.      | 64       |                    |     |
| Ampelopsis, I.    | 159      |                    |     |
| Amsonia, I.       | 121      |                    |     |
| Andromeda, I.     | 252      |                    |     |
| Andropogon, I.    | 56       |                    |     |
| Anemone, I.       | 319      |                    |     |
| Angelica, I.      | 167      |                    |     |
| Anictangium, II.  | 287      |                    |     |
| Anthoceros, II.   | 280      |                    |     |

## B.

|                |     |
|----------------|-----|
| Baccharis, II. | 125 |
| Bartramia, II. | 300 |
| Bartsia, II.   | 17  |
| Batschia, I.   | 129 |
| Befaria, I.    | 280 |
| Bellis, II.    | 151 |



|                    |     |                   |     |
|--------------------|-----|-------------------|-----|
| Crotonopsis, II.   | 185 | Eleusine, I.      | 64  |
| Cteisium, II.      | 275 | Elodea, I.        | 20  |
| Cucubalus, I.      | 271 | Elymus, I.        | 65  |
| Cunila, I.         | 13  | Elytraria, I.     | 8   |
| Cuphea, I.         | 281 | Empetrum, II.     | 255 |
| Cupressus, II.     | 208 | Epigæa, I.        | 250 |
| Cuscuta, I.        | 175 | Epilobium, I.     | 223 |
| Cynoglossum, I.    | 132 | Equisetum, II.    | 281 |
| Cyperus, I.        | 27  | Erianthus, I.     | 54  |
| Cypripedium, II.   | 161 | Erigeron, II.     | 123 |
| Cyrilla, I.        | 157 | Eriocaulon, II.   | 165 |
|                    |     | Eriogonum, I.     | 240 |
| D.                 |     | Eriophorum, I.    | 32  |
| Dalea, II.         | 56  | Eryngium, I.      | 163 |
| Dalibarda, I.      | 209 | Erysimum, II.     | 31  |
| Daucus, I.         | 164 | Erythrina, II.    | 62  |
| Decumaria, I.      | 282 | Erythronium, I.   | 198 |
| Delphinium, I.     | 314 | Erythrorhiza, II. | 34  |
| Dentaria, II.      | 30  | Evonymus, I.      | 155 |
| Dichondra, I.      | 135 | Eupatorium, II.   | 97  |
| Dicranum, II.      | 297 | Euphorbia, II.    | 209 |
| Dichromena, I.     | 37  | Euphrasia, II.    | 16  |
| Diervilla, I.      | 107 | F.                |     |
| Digitaria, I.      | 45  | Fagus, II.        | 194 |
| Dilepyrum, I.      | 40  | Fedia, I.         | 18  |
| Diodia, I.         | 81  | Festuca, I.       | 66  |
| Dionæa, I.         | 267 | Fissidens, II.    | 299 |
| Dioscorea, II.     | 239 | Fothergilla, I.   | 312 |
| Diospyros, II.     | 258 | Fragaria, I.      | 299 |
| Diphylleia, I.     | 203 | Frasera, I.       | 96  |
| Dirca, I.          | 236 | Fraxinus, II.     | 255 |
| Dodecatheon, I.    | 123 | Fuirena, I.       | 37  |
| Doronicum, II.     | 121 | Fumaria, II.      | 51  |
| Draba, II.         | 28  | Funaria, II.      | 303 |
| Dracocephalum, II. | 10  | G.                |     |
| Drosera, I.        | 186 | Galactia, II.     | 61  |
| E.                 |     | Galardia, II.     | 142 |
| Echites, I.        | 120 | Galega, II.       | 67  |
| Eclipta, II.       | 129 | Galium, I.        | 78  |
| Elephantopus, II.  | 148 |                   |     |

Gaultheria, I.  
 Gaura, I.  
 Gelsemium, I.  
 Gentiana, I.  
 Geranium, II.  
 Gerardia, II.  
 Geum, I.  
 Glaux, I.  
 Gleditsia, II.  
 Glycine, II.  
 Gnaphalium, II.  
 Gonolobus, I.  
 Gordonia, I.  
 Gratiola, I.  
 Grimmia, II.  
 Gymnocladus, II.  
 Gymnostomum, II.

## H.

Halesia, II.  
 Hamamelis, I.  
 Hedysarum, II.  
 Helenium, II.  
 Helianthemum, I.  
 Helianthus, II.  
 Heliotropium, I.  
 Helleborus, I.  
 Helonias, I.  
 Helopodium, II.  
 Heracleum, I.  
 Heritiera, I.  
 Hesperis, II.  
 Heuchera, I.  
 Hibiscus, II.  
 Hieracium, II.  
 Hippophaë, II.  
 Hippuris, I.  
 Holcus, I.  
 Hoepa, II.  
 Houstonia, I.  
 Humulus, II.  
 Hydrangea, I.

249 Hydrastis, I.  
 226 Hydrocotyle, I.  
 120 Hydrolea, I.  
 175 Hydropeltis, I.  
 38 Hydrophyllum, I.  
 19 Hymenopappus, II.  
 300 Hyoseris, II.  
 114 Hypericum, II.  
 257 Hypnum, II.  
 63 Hypoxis, I.  
 127 Hyptis, II.  
 119 Hyssopus, II.

## J.

238 Jatropha, II.  
 216 Jeffersonia, I.  
 228 Jlex, II.  
 34 Illecebrum, II.  
 326 Illicium, I.  
 323 Imbricaria, II.  
 149 Inapatiens, II.  
 68 Indigofera, II.  
 123 Inula, II.  
 140 Ipomoea, I.  
 141 Ipomopsis, I.  
 244 Iresine, II.  
 22 Iris, I.  
 3 Isanthus, II.  
 156 Itea, I.  
 184 Iva, II.  
 191 Juglans, II.  
 190 Juncus, I.  
 277 Jungermannia, II.  
 245 Juniperus, II.  
 267 Jussiaea, I.  
 7 Justicia, I.

## K.

230 Kalmia, I.  
 268 Kyllingia, I.

|                     |     |                   |     |
|---------------------|-----|-------------------|-----|
| <b>L.</b>           |     |                   |     |
| Lactuca, II.        | 85  | Malus, I.         | 292 |
| Lamium, II.         | 4   | Manisuris, I.     | 75  |
| Larix, II.          | 203 | Marchantia, II.   | 376 |
| Lathyrus, II.       | 66  | Mayaca, I.        | 26  |
| Laurus, I.          | 243 | Medeola, I.       | 214 |
| Lechea, I.          | 76  | Medicago, II.     | 60  |
| Ledum, I.           | 259 | Meesia, II.       | 303 |
| Leersia, I.         | 39  | Melampyrum, II.   | 16  |
| Lemna, II.          | 163 | Melananthera, II. | 106 |
| Leontodon, II.      | 88  | Melanthium, II.   | 250 |
| Lepidium, II.       | 27  | Melica, I.        | 62  |
| Leptanthus, I.      | 24  | Melothria, II.    | 217 |
| Leskea, II.         | 309 | Menispermum, II.  | 241 |
| Lespedeza, II.      | 70  | Mentha, II.       | 2   |
| Liatris, II.        | 91  | Menyanthes, I.    | 124 |
| Ligusticum, I.      | 166 | Menziesia, I.     | 235 |
| Ligustrum, I.       | 3   | Mespilus, I.      | 291 |
| Lilium, I.          | 197 | Micranthemum, I.  | 10  |
| Limodorum, II.      | 159 | Mimosa, II.       | 254 |
| Linnæa, I.          | 87  | Mimulus, II.      | 23  |
| Linum, II.          | 36  | Mitchella, I.     | 86  |
| Lippia, II.         | 15  | Mitella, I.       | 270 |
| Liquidambar, II.    | 202 | Mnium, II.        | 305 |
| Liriodendron, I.    | 326 | Mollugo, I.       | 77  |
| Lithospermum, I.    | 130 | Monarda, I.       | 16  |
| Lobaria, II.        | 324 | Monniera, II.     | 22  |
| Lobelia, II.        | 151 | Monotropa, I.     | 266 |
| Ludwigia, I.        | 87  | Morus, II.        | 179 |
| Lupinus, II.        | 55  | Myosotis, I.      | 129 |
| Lycium, I.          | 95  | Myrica, II.       | 227 |
| Lycopodium, II.     | 282 | Myriophyllum, II. | 190 |
| Lycopus, I.         | 14  | Myrrhis, I.       | 170 |
| Lysimachia, I.      | 126 | <b>N.</b>         |     |
| Lythrum, I.         | 280 | Najas, II.        | 220 |
| <b>M.</b>           |     | Narthecium, I.    | 209 |
| Magnolia, I.        | 327 | Neckera, II.      | 307 |
| Malachodendron, II. | 43  | Nelumbium, I.     | 317 |
| Malaxis, II.        | 157 | Nepeta, II.       | 2   |
| Malva, II.          | 44  | Nephrodium, II.   | 266 |
|                     |     | Nolina, I.        | 207 |
|                     |     | Nymphæa, I.       | 311 |
|                     |     | Nyssa, II.        | 258 |



## O.

|                   |     |                     |     |
|-------------------|-----|---------------------|-----|
| Oenothera, I.     | 224 | Physcia, II.        | 326 |
| Oldenlandia, I.   | 83  | Phytolacca, I.      | 278 |
| Olea, II.         | 222 | Pinckneya, I.       | 103 |
| Onoclea, II.      | 272 | Pinguicula, I.      | 11  |
| Onosmodium, I.    | 132 | Pinus, II.          | 202 |
| Ophioglossum, II. | 275 | Pistia, II.         | 162 |
| Ophiorrhiza, I.   | 148 | Planera, II.        | 247 |
| Ophrys, II.       | 157 | Plantago, I.        | 94  |
| Orchidocarpum, I. | 329 | Platanus, II.       | 163 |
| Orchis, II.       | 155 | Pleca, I.           | 247 |
| Orchidocarpum, I. | 329 | Poa, I.             | 67  |
| Orchis, II.       | 155 | Podalyria, I.       | 263 |
| Orchidocarpum, I. | 329 | Podophyllum, I.     | 309 |
| Orchidocarpum, I. | 329 | Podostemum, II.     | 164 |
| Orchidocarpum, I. | 329 | Polemonium, I.      | 142 |
| Orchidocarpum, I. | 329 | Polygala, II.       | 52  |
| Orchidocarpum, I. | 329 | Polygonella, II.    | 240 |
| Orchidocarpum, I. | 329 | Polygonum, I.       | 257 |
| Orchidocarpum, I. | 329 | Polymnia, II.       | 147 |
| Orchidocarpum, I. | 329 | Polypodium, II.     | 271 |
| Orchidocarpum, I. | 329 | Polyprenum, I.      | 82  |
| Orchidocarpum, I. | 329 | Polytrichum, II.    | 295 |
| Orchidocarpum, I. | 329 | Pontederia, I.      | 195 |
| Orchidocarpum, I. | 329 | Populus, II.        | 245 |
| Orchidocarpum, I. | 329 | Potamogeton, I.     | 101 |
| Orchidocarpum, I. | 329 | Potentilla, I.      | 302 |
| Orchidocarpum, I. | 329 | Pothos, II.         | 186 |
| Orchidocarpum, I. | 329 | Prenanthes, II.     | 83  |
| Orchidocarpum, I. | 329 | Primula, I.         | 124 |
| Orchidocarpum, I. | 329 | Prinos, II.         | 236 |
| Orchidocarpum, I. | 329 | Proserpinaca, I.    | 76  |
| Orchidocarpum, I. | 329 | Prunus, I.          | 284 |
| Orchidocarpum, I. | 329 | Psilotum, II.       | 281 |
| Orchidocarpum, I. | 329 | Psoralea, II.       | 57  |
| Orchidocarpum, I. | 329 | Psoroma, II.        | 321 |
| Orchidocarpum, I. | 329 | Ptelea, I.          | 99  |
| Orchidocarpum, I. | 329 | Pterigynandrum, II. | 291 |
| Orchidocarpum, I. | 329 | Pteris, II.         | 261 |
| Orchidocarpum, I. | 329 | Pulmonaria, I.      | 131 |
| Orchidocarpum, I. | 329 | Pycnanthemum, II.   | 7   |
| Orchidocarpum, I. | 329 | Pyrola, I.          | 250 |
| Orchidocarpum, I. | 329 | Pyrolaria, II.      | 231 |
| Orchidocarpum, I. | 329 | Pyxidanthera, I.    | 152 |

## P.

|                   |     |                  |     |
|-------------------|-----|------------------|-----|
| Pachysandra, II.  | 177 | Physcia, II.     | 326 |
| Panax, II.        | 256 | Phytolacca, I.   | 278 |
| Pancratium, I.    | 188 | Pinckneya, I.    | 103 |
| Panicum, I.       | 46  | Pinguicula, I.   | 11  |
| Parnassia, I.     | 184 | Pinus, II.       | 202 |
| Parthenium, II.   | 147 | Pistia, II.      | 162 |
| Paspalum, I.      | 43  | Planera, II.     | 247 |
| Passiflora, II.   | 57  | Plantago, I.     | 94  |
| Patellaria, II.   | 320 | Platanus, II.    | 163 |
| Pedicularis, II.  | 18  | Pleca, I.        | 247 |
| Penthorum, I.     | 278 | Poa, I.          | 67  |
| Pentstemon, II.   | 21  | Podalyria, I.    | 263 |
| Persoonia, II.    | 104 | Podophyllum, I.  | 309 |
| Petalostemum, II. | 48  | Podostemum, II.  | 164 |
| Phacelia, I.      | 134 | Polemonium, I.   | 142 |
| Phalangium, I.    | 196 | Polygala, II.    | 52  |
| Phalaris, I.      | 43  | Polygonella, II. | 240 |
| Phaseolus, II.    | 60  | Polygonum, I.    | 257 |
| Philadelphus, I.  | 283 | Polymnia, II.    | 147 |
| Phlox, I.         | 142 | Polypodium, II.  | 271 |
| Phryma, II.       | 15  | Polyprenum, I.   | 82  |
| Phyllanthus, II.  | 209 | Polytrichum, II. | 295 |
| Physalis, I.      | 149 | Pontederia, I.   | 195 |

**CRYPTOGAMIA. LICHENACEAE. 331**

quasi fasciculatis; tuberculis exiguis, hemisphaericis, atro-fuscis.

**Obs.** Varietates duæ: 1<sup>a</sup>. *major*, cinerascens aut rarius pallescenti-incana: 2<sup>a</sup>. *minor*, capillaceo-ramulosissima, ramulis ita dense intricatis ut cæspes veluti crinitum erispam referat.

**Hab.** 1<sup>a</sup>. in Canada; 2<sup>a</sup>. in Carolina.

**STEREOCAULON. ACH.**

**Caules** erecti, solidi, liguescenti rigidi sæpe inæquales et scabridi, vel laeves, ramosi, ramis inordinatis effusis. **Receptacula** *Scutellae* supra planæ, vel etiam rotundata, hemisphaerica et subglobosæ sessiles, sparsæ et terminales.

**PASCHALE.** S. bacillis vage ramosis, granulosa ramulorum materia confertim obsitis; scutellis sparsis, convexis, plerumque compositis, fusco-atris.

**Hab.** in Canada.

**SETARIA. ACH.**

**Caules**: *Lora* cartilaginea filiformia teretiuscula nuda glabra, diffusa et pendula, ramosa, intus tela bombycina farta.

**Receptacula**: *Scutellae* planiusculæ sessiles, demum convexæ, sparsæ, margine integræ.

**Glomeruli** sparsi pulverulenti.

**TRICHODES.** S. tenuissime capilliformis, prolixa,

|                    |     |                  |     |
|--------------------|-----|------------------|-----|
| Stipulicida, I.    | 26  | Urtica, II.      | 178 |
| Streptopus, I.     | 200 | Usnea, II.       | 352 |
| Stylosanthes, II.  | 75  | Utricularia, I.  | 12  |
| Styrax, II.        | 41  | Uvularia, I.     | 198 |
| Swertia, I.        | 97  |                  |     |
| Symphoricarpos, I. | 106 | V.               |     |
|                    |     | Vaccinium, I.    | 227 |
| T.                 |     | Valeriana, I.    | 18  |
| Tagetes, II.       | 132 | Vallisneria, II. | 220 |
| Taxus, II.         | 245 | Veratrum, II.    | 249 |
| Tetraphis, II.     | 287 | Verbascum, I.    | 148 |
| Teucrium, II.      | 1   | Verbena, II.     | 13  |
| Thalictrum, I.     | 322 | Verbesina, II.   | 134 |
| Thesium, I.        | 112 | Vernonia, II.    | 94  |
| Thuja, II.         | 209 | Veronica, I.     | 4   |
| Thymus, II.        | 9   | Viburnum, I.     | 178 |
| Tiarella, I.       | 270 | Vicia, II.       | 69  |
| Tilia, I.          | 306 | Viola, II.       | 149 |
| Tillandsia, I.     | 195 | Vitis, II.       | 230 |
| Timmia, II.        | 304 | Vittaria, II.    | 261 |
| Tortula, II.       | 295 | W.               |     |
| Trachynotia, I.    | 63  | Webera, II.      | 302 |
| Tradescantia, I.   | 193 | Weissia, II.     | 288 |
| Tragia, II.        | 175 | Woodwardia, II.  | 263 |
| Trematodon, II.    | 289 | X.               |     |
| Trichodium, I.     | 41  | Xanthium, II.    | 182 |
| Trichostema, II.   | 10  | Xerophyllum, I.  | 210 |
| Trichostomum, II.  | 296 | Xylosteum, I.    | 106 |
| Trientalis, I.     | 220 | Xyris, I.        | 23  |
| Trifolium, II.     | 58  | Y.               |     |
| Triglochin, I.     | 208 | Yucca, I.        | 196 |
| Trillium, I.       | 215 | Z.               |     |
| Triosticum, I.     | 107 | Zamia, II.       | 242 |
| Tripsacum, I.      | 60  | Zanthorhiza, I.  | 186 |
| Tripterella, I.    | 19  | Zanthoxylum, II. | 235 |
| Tussilago, II.     | 121 | Zigadenus, I.    | 213 |
| U.                 |     | Zizania, I.      | 74  |
| Ulmus, I.          | 172 | Zornia, II.      | 76  |
| Umbilicaria, II.   | 323 |                  |     |
| Uniola, I.         | 70  |                  |     |

# CRYPTOGAMIA. LICHENACEAE. 331

quasi fasciculatis; tuberculis exiguis, hemisphaericis, atro-fuscis.

**Obs.** Varietates duæ : 1<sup>a</sup>. *major*, cinerascens aut rarius pallescenti-incana : 2<sup>a</sup>. *minor*, capillaceo-ramulosissima, ramulis ita dense intricatis ut cæspes veluti crinitum eripam referat.

**Hab.** 1<sup>a</sup>. in Canada; 2<sup>a</sup>. in Carolina.

## STEREOCAULON. ACH.

**Caules** erecti, solidi, liguescentes, rigidi sæpe inæquales et scabridi, vel læves, ramosi, ramis inordinatis affixis. **Receptacula** : **Scutellae** magna, planæ, vel etiam rotundata, hemisphaerica et subglobosæ sessiles, sparsæ et terminales.

**PASCHALE.** S. bacillis vix ramosis, granulosa ramulorum matris confertim obsitis; scutellis sparsis, convexis, plerumque compositis, fusco-atris.

**Hab.** in Canada.

## SETEARIA. ACH.

**Caules** : **Lora** cartilaginea filiformi tereti uscula nuda glabra, diffusa et pendula, ramosa, intus tela bombycina farcta.

**Receptacula** : **Scutellae** planiusculæ sessiles, demum convexæ, sparsæ, margine integræ.

**Glomeruli** sparsi pulverulenti.

**TRICHODES.** S. tenuissime capilliformis, proluxa,



